

# Clinical trials of 12 months dual antiplatelet

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 stent

Trial	Treatments	Patients	Trials design and methods
<b>12 months dual antiplatelet vs 6 months dual antiplatelet</b>			
<a href="#">ISAR-SAFE , 2009</a> [NCT00661206] n=NA follow-up: 12 months	additional 6 month period of clopidogrel versus placebo	Patients on clopidogrel therapy at least 6 months after DES implantation, who do not require a reintervention	Parallel groups double blind

More details and results :

- antithrombotics for stent in all type of patients at <http://www.trialresultscenter.org/go-Q151>
- dual antiplatelet therapy for stent in all type of patients at <http://www.trialresultscenter.org/go-Q578>

## References

### ISAR-SAFE, 2009:

Byrne RA, Schulz S, Mehilli J, Iijima R, Massberg S, Neumann FJ, ten Berg JM, Schmig A, Kastrati A Rationale and design of a randomized, double-blind, placebo-controlled trial of 6 versus 12 months clopidogrel therapy after implantation of a drug-eluting stent: The Intracoronary Stenting and Antithrombotic Regimen: Safety And Efficacy of 6 Months Dual Antiplatelet Therapy After Drug-Eluting Stenting (ISAR-SAFE) study. Am Heart J 2009 Apr;157:620-4.e2 [19332187] [10.1016/j.ahj.2008.12.019](https://doi.org/10.1016/j.ahj.2008.12.019)

## 2 percutaneous coronary intervention

Trial	Treatments	Patients	Trials design and methods
<b>12 months dual antiplatelet vs 6 months dual antiplatelet</b>			
<a href="#">ISAR-SAFE , 2009</a> [NCT00661206] n=NA follow-up: 12 months	additional 6 month period of clopidogrel versus placebo	Patients on clopidogrel therapy at least 6 months after DES implantation, who do not require a reintervention	Parallel groups double blind

More details and results :

- antithrombotics for percutaneous coronary intervention in PCI with stent placement at <http://www.trialresultscenter.org/go-Q152>
- antiplatelets drug for percutaneous coronary intervention in all type of patients at <http://www.trialresultscenter.org/go-Q389>

## References

### **ISAR-SAFE, 2009:**

Byrne RA, Schulz S, Mehilli J, Iijima R, Massberg S, Neumann FJ, ten Berg JM, Schmig A, Kastrati A Rationale and design of a randomized, double-blind, placebo-controlled trial of 6 versus 12 months clopidogrel therapy after implantation of a drug-eluting stent: The Intracoronary Stenting and Antithrombotic Regimen: Safety And Efficacy of 6 Months Dual Antiplatelet Therapy After Drug-Eluting Stenting (ISAR-SAFE) study. Am Heart J 2009 Apr;157:620-4.e2 [19332187] 10.1016/j.ahj.2008.12.019