

Clinical trials of 3-6 months

TrialResults-center www.trialresultscenter.org

1 venous thrombosis

Trial	Treatments	Patients	Trials design and methods
3-6 months vs 1.5-3 months			
Pinede , 2001 n=NA follow-up: 15 months after randomization	Long course of therapy (6 months for proximal DVT and/or PE; 12 weeks for calf DVT) by fluindione adjusted for INR range of 2.0 to 3.0 versus Short oral anticoagulant course (3 months for proximal DVT and/or PE; 6 weeks for isolated calf DVT) by fluindione adjusted for INR range of 2.0 to 3.0	-	Parallel groups open France

More details and results :

- antithrombotics for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q101>
- heparin (UFH or LMWH) for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q204>

References

Pinede, 2001:

Pinede L, Ninet J, Duhaut P, Chabaud S, Demolombe-Rague S, Durieu I, Nony P, Sanson C, Boissel JP Comparison of 3 and 6 months of oral anticoagulant therapy after a first episode of proximal deep vein thrombosis or pulmonary embolism and comparison of 6 and 12 weeks of therapy after isolated calf deep vein thrombosis. *Circulation* 2001;103:2453-60 [[11369685](#)]

Entry terms: warfarin