

# Clinical trials of BIAsp 30)twice-daily + metformin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
<b>BIAsp 30)twice-daily + metformin vs once-daily glargine + metformin + secretagogues</b>			
<a href="#">Ligthelm , 2011</a> n=NA follow-up: 24 weeks	biphasic insulin aspart 70/30 (BIAsp 30)twice-daily + metformin versus once-daily glargine + metformin + secretagogues	type 2 diabetic patients who were not maintaining glycemic control on basal insulin and oral antidiabetic drugs	Parallel groups open-label

More details and results :

- insulin therapy for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q548>

↑

## References

### Ligthelm, 2011:

Ligthelm RJ, Gylvin T, DeLuzio T, Raskin P A comparison of twice-daily biphasic insulin aspart 70/30 and once-daily insulin glargine in persons with type 2 diabetes mellitus inadequately controlled on basal insulin and oral therapy: a randomized, open-label study. *Endocr Pract* 2011;17:41-50 [[20713345](#)] [10.4158/EP10079.OR](#)