

Clinical trials of GCS + LMWH

TrialResults-center www.trialresultscenter.org

1 DVT prophylaxis

Trial	Treatments	Patients	Trials design and methods
GCS + LMWH vs LMWH			
Kalodiki (GCS+LMWH vs LMWH) , 1996 n=NA follow-up:	enoxaparin (40 mg once daily) plus graduated elastic compression (TEDR stockings) for 8-12 days versus low molecular weight heparin: (enoxaparin 40 mg once daily)	patients having elective total hip replacement	Parallel groups

More details and results :

- graduated compression stockings for DVT prophylaxis in all type of patients at <http://www.trialresultscenter.org/go-Q158>
- mechanical devices for thromboprophylaxis for DVT prophylaxis in all type of patients at <http://www.trialresultscenter.org/go-Q402>
- mechanical devices for thromboprophylaxis for DVT prophylaxis in orthopaedic surgery at <http://www.trialresultscenter.org/go-Q465>

References

Kalodiki (GCS+LMWH vs LMWH), 1996:

Kalodiki EP, Hoppensteadt DA, Nicolaidis AN, Fareed J, Gill K, Regan F, al-Kutoubi A, Cunningham DA, Birch R, Harris N, Hunt D, Johnson J, Marx C Deep venous thrombosis prophylaxis with low molecular weight heparin and elastic compression in patients having total hip replacement. A randomised controlled trial. Int Angiol 1996;15:162-8 [[8803642](#)]