

Clinical trials of HNF

TrialResults-center www.trialresultscenter.org

1 peripheral vascular diseases

Trial	Treatments	Patients	Trials design and methods
HNF vs placebo			
Tesi , 1989 n=10/10 follow-up: 6 mois	HBPM 8000 U / j versus Placebo	AOMI stade II	Parallel groups Double aveugle
Antonicelli , 1999 n=101/100 follow-up: 3 mois	Hparine calcique 12500 IU / j versus Placebo	AOMI stade II	Parallel groups Double aveugle

More details and results :

- antithrombotics for peripheral vascular diseases in all type of patients at <http://www.trialresultscenter.org/go-Q50>

References

Tesi, 1989:

Tesi M, Bronchi GF, Carini A, Morfini M, Cinotti S, Filiberti E. Efficacy and safety of a new low molecular weight heparin in the medium- term treatment of atherosclerotic arteriopathy of the lower limbs. J Drug Dev 1989; 2(2):73-82.

Antonicelli, 1999:

Randomized trial of the effects of low-dose calcium-heparin in patients with peripheral arterial disease and claudication. Italian CAP Study Group. Antonicelli R, Sardina M, Scotti A, Bonizzoni E, Paciaroni E Am J Med 1999 Sep;107:234-9 [[10492316](#)]

Entry terms: UFH