

Clinical trials of IPC

TrialResults-center www.trialresultscenter.org

1 thrombosis prevention

Trial	Treatments	Patients	Trials design and methods
IPC vs aspirin			
HAAS I , 1990 n=NA	-	elective knee replacement	open
HAAS II , 1990 n=NA	-	elective knee replacement	open
McKenna (vs aspirin) , 1980 n=NA	-	elective knee replacement	open
IPC + aspirin vs aspirin			
Hull 2 (+asp) , 1979 n=NA follow-up:	-	patients undergoing elective knee surgery	open
Hull (+asp) , 1979 n=NA follow-up:	-	patients undergoing elective knee surgery	Parallel groups open
65279;Lieberman (A) , 1994 n=130/130 follow-up:	hypotensive epidural anesthesia, external pneumatic-compression boots, and aspirin versus hypotensive epidural anesthesia and aspirin	primary unilateral or bilateral total hip arthroplasty with use of hypotensive epidural anesthesia	Parallel groups open
thigh Length IPCD vs calf Length IPCD			
Soderdahl , 1997 n=NA follow-up: in hospital	Thigh Length IPCD versus Calf Length IPCD	patients undergoing major urological surgery	Parallel groups open
IPC + GCS vs control			
Turpie , 1989 n=NA follow-up: 14 days	graduated compression stockings plus intermittent pneumatic compression versus untreated control	potential neurosurgical patients	Parallel groups open
IPC + UFH vs control			

continued...

Trial	Treatments	Patients	Trials design and methods
Killewich , 1997 n=NA follow-up:	low-dose heparin sodium therapy (5000 U every 12 hours) and calf-length intermittent mechanical compression devices versus no prophylaxis	patients undergoing aortic reconstruction for aneurysmal or occlusive disease	open
IPC sequential compression vs control			
65279;Blackshear exclude , 1987 n=20/20 follow-up:	Sequential external pneumatic compression versus control	abdominal or thoracic surgery	Cross over open
Hull II , 1990 n=152/158 follow-up:	sequential intermittent calf and thigh compression versus no prophylaxis	patients undergoing total hip replacement	open
Fisher , 1995 n=145/159 follow-up:	pneumatic sequential leg compression devices versus no specific form of prophylaxis	orthopaedic trauma patients with hip and pelvic fractures	Parallel groups open
Turple II , 1979 n=112/106 follow-up:	-	patients with intracranial disease	open
Knudson , 1994 n=26/39 follow-up:	sequential gradient pneumatic leg compression versus control	trauma patients	open
Kosir , 1996 n=25/45 follow-up: 30 days	sequential pneumatic compression devices during surgery and 2 days postoperatively versus control	Patients undergoing procedures of at least 1 hr duration (abdominal, thoracic, head and neck, inguinal) requiring general or spinal anesthetic	Parallel groups open
IPC single compression vs control			
Prasad , 1982 n=NA follow-up:	pneumatic compression of the calf versus No Prophylaxis	elderly hemiplegics	Parallel groups open
Butson , 1981 n=62/57 follow-up:	intermittent pneumatic calf compression, begun after the induction of anesthesia and continued until the patient was walking versus control	patients undergoing major abdominal general surgical procedures	open

continued...

Trial	Treatments	Patients	Trials design and methods
Clark , 1974 n=36/36 follow-up:	-	abdominal or thoracic surgery	open
Clarke-Pearson I , 1984 n=59/57 follow-up:	-	-	open
Clarke-Pearson II , 1984 n=104/105 follow-up:	-	abdominal or thoracic surgery	open
Coe , 1978 n=31/24 follow-up:	low-dose heparin and external pneumatic compression (EPC) of the calves versus no treatment	patients undergoing open urological operations	open
Hills , 1972 n=70/70 follow-up:	-	abdominal or thoracic surgery	open
Gallus , 1983 n=43/47 follow-up: in hospital	preventive intermittent calf compression versus no prophylaxis	elective hip replacement	Parallel groups open
65279;Bachmann , 1976 n=26/28 follow-up: until patients fully mobilized	-	patients undergoing joint-replacement operations	Parallel groups open
Bynke , 1987 n=31/31 follow-up:	-	neurosurgery	open
Skillman , 1978 n=47/48 follow-up: NA	external pneumatic compression (EPC) of the calves versus no specific form of prophylaxis	patients undergoing neurosurgical operative procedures	Parallel groups open
Turpie I , 1977 n=82/79 follow-up:	-	patients with intracranial disease	open
Hull I , 1979 n=32/29 follow-up:	-	patients undergoing elective knee surgery	Parallel groups open
IPC unknown type vs control			

continued...

Trial	Treatments	Patients	Trials design and methods
Weitz , 1986 n=5/9 follow-up:	-	-	Parallel groups open
IPCD vs control			
McKenna , 1980 n=10/12 follow-up:	IPCD versus No Prophylaxis	Elective knee replacement	Parallel groups open
IPC + dextran vs dextran			
Smith (D) , 1978 n=97/97 follow-up:	dextran 70 and pneumatic calf compression versus dextran 70	-	open
IPC vs DHE + low dose heparin			
Mellbring , 1986 n=54/54 follow-up:	peroperative intermittent pneumatic calf compression versus dihydroergotamine (DHE) combined with low-dose heparin (LDH)	patients undergoing Abdominal surgery	Parallel groups open
IPC sequential compression vs Footpump			
Elliott , 1999 n=149 follow-up: NA	Calf-thigh sequential pneumatic compression versus foot pump (plantar venous pneumatic compression)	Trauma patients >13 years old	Parallel groups open (blind assesement) United States
IPC + GCS vs GCS			
Goldhaber , 1995 n=164/166 follow-up: hospital stay	intermittent pneumatic compression (IPC) plus graduated compression stockings (GCS) versus standard compression stockings alone	patients undergoing coronary artery bypass without concomitant valve surgery or coronary endarterectomy	open
Fordyce , 1992 n=NA follow-up:	venous foot pump (A-V Impulse System) versus control	elective hip replacement	open

continued...

Trial	Treatments	Patients	Trials design and methods
Rokito , 1996 n=NA follow-up:	TED stockings and thigh-length cuffs that provided sequential pneumatic compression to the calf and thigh versus bilateral thigh-high thrombosis embolic deterrent (TED) compression stockings (Kendall Company, MA).	neurosurgery	Parallel groups open
Turpie , 1989 n=NA follow-up: 14 days	graduated compression stockings plus intermittent pneumatic compression versus graduated compression stockings alone	neurosurgery	open
Wautrecht , 1996 n=NA	-	neurosurgery	open
Caprini , 1983 n=NA	-	general surgery	open
Lacut , 2005 n=151 follow-up: 1 days	elastic stockings combined with intermittent pneumatic compression versus elastic stockings alone	patients with a documented intracerebral hemorrhage	Parallel groups open
Pambianco , 1995 n=NA	-	stroke	open
IPC + GCS +LMWH vs GCS +LMWH			
Dickinson , 1998 n=23/21 follow-up: 1 month	sequential compression device +enoxaparin (+ GCS) versus enoxaparin (+GCS)	neurosurgery, patients with brain tumors	open
IPC/FID vs GES			
Hansberry , 1991 n=NA	-	-	open
IPC vs LMWH			
Kurtoglu , 2004 n=60/60 follow-up: 1 week after discharge	intermittent pneumatic compression versus LMWH	head/spinal traumatized patients	Parallel groups open Turkey
Warwick excludure , 2002 n=NA	IPCD/FID versus LMWH	-	open

continued...

Trial	Treatments	Patients	Trials design and methods
Ginzburg , 2003 n=224/218 follow-up: 30 days	intermittent pneumatic compression versus LMWH	Trauma patients >18 years old	Parallel groups open United States
IPC + GCS vs LMWH			
Norgren , 1998 n=NA follow-up:	IPCD/FID + GCS versus LMWH	elective knee replacement	open
IPC vs UFH			
Clarke Pearson , 1993 n=NA	-	-	open
Coe (vs UFH) , 1978 n=NA	-	-	open
Hansberry (vs UFH) , 1991 n=NA	-	-	open
Kosir , 1998 n=NA	-	-	open
IPC + GCS vs UFH			
Niolaides , 1983 n=NA follow-up:	IPCD + GCS versus UFH	general surgery	open
Santori , 1994 n=67/65 follow-up:	IPC + GCS versus UFH	elective hip replacement	open
IPC + UFH vs UFH			
Ramos , 1996 n=1355/1196 follow-up:	prophylactic regimen of Pneumatic compression stocking and subcutaneous heparin versus with 5,000 U of subcutaneous heparin every 12 h	cardiac surgery	Parallel groups open
Siragusa (H) , 1994 n=35/35 follow-up:	-	-	Parallel groups open
IPC vs vitamin K antagonists			
Chandhoke , 1992 n=NA follow-up:	-	major urological surgery	open

continued...

Trial	Treatments	Patients	Trials design and methods
paieiment , 1987 n=NA	-	elective hip replacement	open
Kaempffe , 1991 n=NA	-	elective hip & knee replacement	open
GCS +IPC vs IPC			
Mellbring , 1986 n=NA follow-up:	graduated static compression + peroperative intermittent pneumatic calf compression versus peroperative intermittent pneumatic calf compression	patients undergoing major abdominal surgery	Factorial plan
Scurr (GCS+IPC vs IPC) , 1987 n=78/78 follow-up:	simultaneous use of graduated compression stockings and intermittent sequential pneumatic compression versus intermittent sequential pneumatic compression	patients undergoing Abdominal surgery	parallel groups

More details and results :

- graduated compression stockings for thrombosis prevention in all type of patients at <http://www.trialresultscenter.org/go-Q158>
- mechanical devices for thromboprophylaxis for thrombosis prevention in all type of patients at <http://www.trialresultscenter.org/go-Q402>
- mechanical devices for thromboprophylaxis for thrombosis prevention in general non orthopaedic surgery at <http://www.trialresultscenter.org/go-Q466>

References

HAAS I, 1990:

Haas SB, Insall JN, Scuderi GR, Windsor RE, Ghelman B Pneumatic sequential-compression boots compared with aspirin prophylaxis of deep-vein thrombosis after total knee arthroplasty. J Bone Joint Surg Am 1990;72:27-31 [2404020]

HAAS II, 1990:

Haas SB, Insall JN, Scuderi GR, Windsor RE, Ghelman B Pneumatic sequential-compression boots compared with aspirin prophylaxis of deep-vein thrombosis after total knee arthroplasty. J Bone Joint Surg Am 1990;72:27-31 [2404020]

McKenna (vs aspirin), 1980:

McKenna R, Galante J, Bachmann F, Wallace DL, Kaushal PS, Meredith P Prevention of venous thromboembolism after total knee replacement by high-dose aspirin or intermittent calf and thigh compression. Br Med J 1980;280:514-7 [6989432]

Hull 2 (+asp), 1979:

Hull R, Delmore TJ, Hirsh J, Gent M, Armstrong P, Lofthouse R, MacMillan A, Blackstone I, Reed-Davis R, Detwiler RC Effectiveness of intermittent pulsatile elastic stockings for the prevention of calf and thigh vein thrombosis in patients undergoing elective knee surgery. *Thromb Res* 1979;16:37-45 [[505427](#)]

Hull (+asp), 1979:

Hull R, Delmore TJ, Hirsh J, Gent M, Armstrong P, Lofthouse R, MacMillan A, Blackstone I, Reed-Davis R, Detwiler RC Effectiveness of intermittent pulsatile elastic stockings for the prevention of calf and thigh vein thrombosis in patients undergoing elective knee surgery. *Thromb Res* 1979;16:37-45 [[505427](#)]

65279;Lieberman (A), 1994:

Lieberman JR, Huo MM, Hanway J, Salvati EA, Sculco TP, Sharrock NE The prevalence of deep venous thrombosis after total hip arthroplasty with hypotensive epidural anesthesia. *J Bone Joint Surg Am* 1994;76:341-8 [[8126039](#)]

Soderdahl, 1997:

Soderdahl DW, Henderson SR, Hansberry KL A comparison of intermittent pneumatic compression of the calf and whole leg in preventing deep venous thrombosis in urological surgery. *J Urol* 1997;157:1774-6 [[9112525](#)]

Turpie, 1989:

Turpie AG, Hirsh J, Gent M, Julian D, Johnson J Prevention of deep vein thrombosis in potential neurosurgical patients. A randomized trial comparing graduated compression stockings alone or graduated compression stockings plus intermittent pneumatic compression with control. *Arch Intern Med* 1989;149:679-81 [[2645846](#)]

Killewich, 1997:

Killewich LA, Aswad MA, Sandager GP, Lilly MP, Flinn WR A randomized, prospective trial of deep venous thrombosis prophylaxis in aortic surgery. *Arch Surg* 1997;132:499-504 [[9161392](#)]

65279;Blackshear exclure, 1987:

Blackshear WM Jr, Prescott C, LePain F, Benoit S, Dickstein R, Seifert KB Influence of sequential pneumatic compression on postoperative venous function. *J Vasc Surg* 1987;5:432-6 [[3509596](#)]

Hull II, 1990:

Hull RD, Raskob GE, Gent M, McLoughlin D, Julian D, Smith FC, Dale NI, Reed-Davis R, Lofthouse RN, Anderson C Effectiveness of intermittent pneumatic leg compression for preventing deep vein thrombosis after total hip replacement. *JAMA* 1990;263:2313-7 [[2182917](#)]

Fisher, 1995:

Fisher CG, Blachut PA, Salvian AJ, Meek RN, O'Brien PJ Effectiveness of pneumatic leg compression devices for the prevention of thromboembolic disease in orthopaedic trauma patients: a prospective, randomized study of compression alone versus no prophylaxis. *J Orthop Trauma* 1995;9:1-7 [[7714648](#)]

Turple II, 1979:

Turpie AG, Delmore T, Hirsh J, Hull R, Genton E, Hiscoe C, Gent M Prevention of venous thrombosis by intermittent sequential calf compression in patients with intracranial disease. *Thromb Res* 1979;15:611-6 [[494166](#)]

Knudson, 1994:

Knudson MM, Lewis FR, Clinton A, Atkinson K, Megerman J Prevention of venous thromboembolism in trauma patients. *J Trauma* 1994;37:480-7 [[8083913](#)]

Kosir, 1996:

Kosir MA, Kozol RA, Perales A, McGee K, Beleski K, Lange P, Dahn M Is DVT prophylaxis overemphasized? A randomized prospective study. *J Surg Res* 1996;60:289-92 [[8598656](#)] [10.1006/jsre.1996.0045](#)

Prasad, 1982:

Prasad BK, Banerjee AK, Howard H Incidence of deep vein thrombosis and the effect of pneumatic compression of the calf in elderly hemiplegics. *Age Ageing* 1982;11:42-4 [7072559]

Butson, 1981:

Butson AR Intermittent pneumatic calf compression for prevention of deep venous thrombosis in general abdominal surgery. *Am J Surg* 1981;142:525-7 [7025675]

Clark, 1974:

Clark WB, MacGregor AB, Prescott RJ, Ruckley CV Pneumatic compression of the calf and postoperative deep-vein thrombosis. *Lancet* 1974;2:5-7 [4134404]

Clarke-Pearson I, 1984:

Clarke-Pearson DL, Synan IS, Hinshaw WM, Coleman RE, Creasman WT Prevention of postoperative venous thromboembolism by external pneumatic calf compression in patients with gynecologic malignancy. *Obstet Gynecol* 1984 Jan;63:92-8 [6691021]

Clarke-Pearson II, 1984:

Clarke-Pearson DL, Creasman WT, Coleman RE, Synan IS, Hinshaw WM Perioperative external pneumatic calf compression as thromboembolism prophylaxis in gynecologic oncology: report of a randomized controlled trial. *Gynecol Oncol* 1984 Jun;18:226-32 [6376294]

Coe, 1978:

Coe NP, Collins RE, Klein LA, Bettmann MA, Skillman JJ, Shapiro RM, Salzman EW Prevention of deep vein thrombosis in urological patients: a controlled, randomized trial of low-dose heparin and external pneumatic compression boots. *Surgery* 1978;83:230-4 [622696]

Hills, 1972:

Hills NH, Pflug JJ, Jeyasingh K, Boardman L, Calnan JS Prevention of deep vein thrombosis by intermittent pneumatic compression of calf. *Br Med J* 1972;1:131-5 [5007837]

Gallus, 1983:

Gallus A, Raman K, Darby T Venous thrombosis after elective hip replacement—the influence of preventive intermittent calf compression and of surgical technique. *Br J Surg* 1983;70:17-9 [6337668]

65279;Bachmann, 1976:

Bachmann F, McKenna R, Meredith P, Carta S [Intermittent pneumatic compression of leg and thigh: a new successful method for the prevention of postoperative thrombosis] *Schweiz Med Wochenschr* 1976;106:1819-21 [1013707]

Bynke, 1987:

Bynke O, Hillman J, Lassvik C Does peroperative external pneumatic leg muscle compression prevent post-operative venous thrombosis in neurosurgery? *Acta Neurochir (Wien)* 1987;88:46-8 [3425413]

Skillman, 1978:

Skillman JJ, Collins RE, Coe NP, Goldstein BS, Shapiro RM, Zervas NT, Bettmann MA, Salzman EW Prevention of deep vein thrombosis in neurosurgical patients: a controlled, randomized trial of external pneumatic compression boots. *Surgery* 1978;83:354-8 [628896]

Turpie I, 1977:

Turpie AG, Gallus A, Beattie WS, Hirsh J Prevention of venous thrombosis in patients with intracranial disease by intermittent pneumatic compression of the calf. *Neurology* 1977;27:435-8 [558547]

Hull I, 1979:

Hull R, Delmore TJ, Hirsh J, Gent M, Armstrong P, Lofthouse R, MacMillan A, Blackstone I, Reed-Davis R, Detwiler RC Effectiveness of intermittent pulsatile elastic stockings for the prevention of calf and thigh vein thrombosis in patients undergoing elective knee surgery. *Thromb Res* 1979;16:37-45 [505427]

Weitz, 1986:

Weitz J, Michelsen J, Gold K, Owen J, Carpenter D Effects of intermittent pneumatic calf compression on postoperative thrombin and plasmin activity. *Thromb Haemost* 1986;56:198-201 [2949390]

McKenna, 1980:

McKenna R, Galante J, Bachmann F, Wallace DL, Kaushal PS, Meredith P Prevention of venous thromboembolism after total knee replacement by high-dose aspirin or intermittent calf and thigh compression. *Br Med J* 1980 Feb 23;280:514-7 [6989432]

Smith (D), 1978:

Smith RC, Elton RA, Orr JD, Hart AJ, Graham IF, Fuller GA, Rundle JS, Macpherson AI, Ruckley CV Dextran and intermittent pneumatic compression in prevention of postoperative deep vein thrombosis: multiunit trial. *Br Med J* 1978;1:952-4 [638545]

Mellbring, 1986:

Mellbring G, Palmr K Prophylaxis of deep vein thrombosis after major abdominal surgery. Comparison between dihydroergotamine-heparin and intermittent pneumatic calf compression and evaluation of added graduated static compression. *Acta Chir Scand* 1986;152:597-600 [3544626]

Elliott, 1999:

Elliott CG, Dudney TM, Egger M, Orme JF, Clemmer TP, Horn SD, Weaver L, Handrahan D, Thomas F, Merrell S, Kitterman N, Yeates S Calf-thigh sequential pneumatic compression compared with plantar venous pneumatic compression to prevent deep-vein thrombosis after non-lower extremity trauma. *J Trauma* 1999;47:25-32 [10421182]

Goldhaber, 1995:

Goldhaber SZ, Hirsch DR, MacDougall RC, Polak JF, Creager MA, Cohn LH Prevention of venous thrombosis after coronary artery bypass surgery (a randomized trial comparing two mechanical prophylaxis strategies). *Am J Cardiol* 1995 Nov 15;76:993-6 [7484878]

Fordyce, 1992:

Fordyce MJ, Ling RS A venous foot pump reduces thrombosis after total hip replacement. *J Bone Joint Surg Br* 1992 Jan;74:45-9 [1732264]

Rokito, 1996:

Rokito SE, Schwartz MC, Neuwirth MG Deep vein thrombosis after major reconstructive spinal surgery. *Spine (Phila Pa 1976)* 1996 Apr 1;21:853-8; discussion 859 [8779018]

Turpie, 1989:

Turpie AG, Hirsh J, Gent M, Julian D, Johnson J Prevention of deep vein thrombosis in potential neurosurgical patients. A randomized trial comparing graduated compression stockings alone or graduated compression stockings plus intermittent pneumatic compression with control. *Arch Intern Med* 1989 Mar;149:679-81 [2645846]

Wautrecht, 1996:

Wautrecht JC, Macquaire V, Vandesteene A, Daoud N, Golzarian J, Capel P. Prevention of deep vein thrombosis in neurosurgical patients with brain tumors: a controlled, randomized study comparing graded compression stockings alone and intermittent sequential compression. Correlation with pre- and postoperative fibrinolysis: preliminary results. *International Angiology* 1996, 15:5-10

Caprini, 1983:

Caprini JA, Chucker JL, Zuckerman L, Vagher JP, Franck CA, Cullen JE Thrombosis prophylaxis using external compression. *Surg Gynecol Obstet* 1983 May;156:599-604 [6845123]

Lacut, 2005:

Lacut K, Bressollette L, Le Gal G, Etienne E, De Tinteniak A, Renault A, Rouhart F, Besson G, Garcia JF, Mottier D, Oger E Prevention of venous thrombosis in patients with acute intracerebral hemorrhage. *Neurology* 2005 Sep 27;65:865-9 [[16186525](#)] [10.1212/01.wnl.0000176073.80532.a2](#)

Pambianco, 1995:

Pambianco G, Orchard T, Landau P Deep vein thrombosis: prevention in stroke patients during rehabilitation. *Arch Phys Med Rehabil* 1995 Apr;76:324-30 [[7717832](#)]

Dickinson, 1998:

Dickinson LD, Miller LD, Patel CP, Gupta SK Enoxaparin increases the incidence of postoperative intracranial hemorrhage when initiated preoperatively for deep venous thrombosis prophylaxis in patients with brain tumors. *Neurosurgery* 1998 Nov;43:1074-81 [[9802851](#)]

Hansberry, 1991:

Hansberry KL, Thompson IM Jr, Bauman J, Deppe S, Rodriguez FR A prospective comparison of thromboembolic stockings, external sequential pneumatic compression stockings and heparin sodium/dihydroergotamine mesylate for the prevention of thromboembolic complications in urological surgery. *J Urol* 1991;145:1205-8 [[2033693](#)]

Kurtoglu, 2004:

Kurtoglu M, Yanar H, Bilsel Y, Guloglu R, Kizilirmak S, Buyukkurt D, Granit V Venous thromboembolism prophylaxis after head and spinal trauma: intermittent pneumatic compression devices versus low molecular weight heparin. *World J Surg* 2004;28:807-11 [[15457363](#)]

Warwick exlure, 2002:

Warwick DJ, Pandit H, Shewale S, Sulkin T Venous impulse foot pumps: should graduated compression stockings be used? *J Arthroplasty* 2002 Jun;17:446-8 [[12066274](#)]

Ginzburg, 2003:

Ginzburg E, Cohn SM, Lopez J, Jackowski J, Brown M, Hameed SM Randomized clinical trial of intermittent pneumatic compression and low molecular weight heparin in trauma. *Br J Surg* 2003;90:1338-44 [[14598411](#)]

Norgren, 1998:

Norgren L, Toksvig-Larsen S, Magyar G, Lindstrand A, Albrechtsson U Prevention of deep vein thrombosis in knee arthroplasty. Preliminary results from a randomized controlled study of low molecular weight heparin vs foot pump compression. *Int Angiol* 1998 Jun;17:93-6 [[9754896](#)]

Clarke Pearson, 1993:

Clarke-Pearson DL, Synan IS, Dodge R, Soper JT, Berchuck A, Coleman RE A randomized trial of low-dose heparin and intermittent pneumatic calf compression for the prevention of deep venous thrombosis after gynecologic oncology surgery. *Am J Obstet Gynecol* 1993 Apr;168:1146-53; discussion 1153-4 [[8475960](#)]

Coe (vs UFH), 1978:

Hansberry (vs UFH), 1991:

Hansberry KL, Thompson IM Jr, Bauman J, Deppe S, Rodriguez FR A prospective comparison of thromboembolic stockings, external sequential pneumatic compression stockings and heparin sodium/dihydroergotamine mesylate for the prevention of thromboembolic complications in urological surgery. *J Urol* 1991 Jun;145:1205-8 [[2033693](#)]

Kosir, 1998:

Kosir MA, Schmittinger L, Barno-Winarski L, Duddella P, Pone M, Perales A, Lange P, Brish LK, McGee K, Beleski K, Pawlak J, Mammen E, Sajahan NP, Kozol RA Prospective double-arm study of fibrinolysis in surgical patients. *J Surg Res* 1998;74:96-101 [[9536981](#)] [10.1006/jsre.1997.5233](#)

Niolaides, 1983:

Nicolaides AN, Miles C, Hoare M, Jury P, Helms E, Venniker R Intermittent sequential pneumatic compression of the legs and thromboembolism-deterrent stockings in the prevention of postoperative deep venous thrombosis. *Surgery* 1983 Jul;94:21-5 [6857507]

Santori, 1994:

Santori FS, Vitullo A, Stopponi M, Santori N, Ghera S Prophylaxis against deep-vein thrombosis in total hip replacement. Comparison of heparin and foot impulse pump. *J Bone Joint Surg Br* 1994 Jul;76:579-83 [8027144]

Ramos, 1996:

Ramos R, Salem BI, De Pawlikowski MP, Coordes C, Eisenberg S, Leidenfrost R The efficacy of pneumatic compression stockings in the prevention of pulmonary embolism after cardiac surgery. *Chest* 1996 Jan;109:82-5 [8549224]

Siragusa (H), 1994:

Siragusa S, Vicentini L, Carbone S, Barone M, Beltrametti C, Piovella F. Intermittent pneumatic leg compression (IPLC) and unfractionated heparin (UFH) in the prevention of post-operative deep vein thrombosis in hip surgery: a randomized clinical trial *Br J Haematol* 1994; 87:186.

Chandhoke, 1992:

Chandhoke PS, Gooding GA, Narayan P Prospective randomized trial of warfarin and intermittent pneumatic leg compression as prophylaxis for postoperative deep venous thrombosis in major urological surgery. *J Urol* 1992 Apr;147:1056-9 [1552582]

paient, 1987:

Paient G, Wessinger SJ, Waltman AC, Harris WH Low-dose warfarin versus external pneumatic compression for prophylaxis against venous thromboembolism following total hip replacement. *J Arthroplasty* 1987;2:23-6 [3572408]

Kaempffe, 1991:

Kaempffe FA, Lifeso RM, Meinking C Intermittent pneumatic compression versus coumadin. Prevention of deep vein thrombosis in lower-extremity total joint arthroplasty. *Clin Orthop Relat Res* 1991 Aug;:89-97 [1864061]

Mellbring, 1986:

Mellbring G, Palmr K Prophylaxis of deep vein thrombosis after major abdominal surgery. Comparison between dihydroergotamine-heparin and intermittent pneumatic calf compression and evaluation of added graduated static compression. *Acta Chir Scand* 1986;152:597-600 [3544626]

Scurr (GCS+IPC vs IPC), 1987:

Scurr JH, Coleridge-Smith PD, Hasty JH Regimen for improved effectiveness of intermittent pneumatic compression in deep venous thrombosis prophylaxis. *Surgery* 1987;102:816-20 [3672322]

Entry terms: aspirin, UFH