

Clinical trials of Minoctoparine

TrialResults-center www.trialresultscenter.org

1 venous thrombosis

| Trial | Treatments | Patients | Trials design and methods |
|----------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------|----------|---------------------------|
| Minoctoparine vs unfractionated heparin | | | |
| Faivre et al , 1988 n=33/37 follow-up: 10 Days | Minoctoparine (CY222) Subcutaneous twice daily for 10 Days,155 U/kg BID versus unfractionated heparin subcutaneous twice daily APPTx2-3 | - | |

More details and results :

- antithrombotics for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q101>
- LMWH for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q203>
- heparin (UFH or LMWH) for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q204>

References

Faivre et al , 1988:

Faivre R, Neuhart Y, Kieffer Y, Apfel F, Magnin D, Didier D, Toulemonde F, Bassand JP, Maurat JP [A new treatment of deep venous thrombosis: low molecular weight heparin fractions. Randomized study] Presse Med 1988 Feb 13;17:197-200 [[2965375](#)]