

Clinical trials of Omacor

TrialResults-center www.trialresultscenter.org

1 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
Omacor vs control			
Eritsland , 1996 n=317/293 follow-up: 12 months	Omacor capsules, 4/d (3.3g EPA + DHA daily) versus no treatment	people admitted for coronary bypass grafting	Parallel groups open Norway
GISSI-P , 1999 n=5665/5668 follow-up: median 40 months	Omacor gelatine capsules, 1/d (0.9g/d EPA + DHA daily) versus no treatment	people with recent myocardial infarction	Parallel groups open Italy
Omacor vs placebo			
Johansen , 1999 n=250/250 follow-up: 6.5 months	Omacor capsules, 6/d (5g EPA + DHA daily) versus placebo (corn oil capsules, 6/d)	people about to undergo elective coronary angioplasty	Parallel groups double blind Norway
Nilsen , 2001 n=150/150 follow-up: 24 months	Omacor capsules 4/d (3.5g EPA + DHA) versus placebo (corn oil capsules, 4/d)	people with acute myocardial infarction 4-8 days agoe/pj	Parallel groups double-blind Norway

More details and results :

- omega-3 fatty acids for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q121>
- omega-3 fatty acids for cardiovascular prevention in patients at high risk at <http://www.trialresultscenter.org/go-Q123>

References

Eritsland, 1996:

Eritsland J, Arnesen H, Gronseth K, Fjeld NB, Abdelnoor M Effect of dietary supplementation with n-3 fatty acids on coronary artery bypass graft patency. Am J Cardiol 1996 Jan 1;77:31-6 [8540453]

Eritsland J, Arnesen H, Seljeflot I, Hostmark AT Long-term metabolic effects of n-3 polyunsaturated fatty acids in patients with coronary artery disease. Am J Clin Nutr 1995 Apr;61:831-6 [7702027]

GISSI-P, 1999:

Dietary supplementation with n-3 polyunsaturated fatty acids and vitamin E after myocardial infarction: results of the GISSI-Prevenzione trial. Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto miocardico. *Lancet* 1999 Aug 7;354:447-55 [[10465168](#)]

Marchioli R, Barzi F, Bomba E, Chieffo C, Di Gregorio D, Di Mascio R, Franzosi MG, Geraci E, Levantesi G, Maggioni AP, Mantini L, Marfisi RM, Mastrogiuseppe G, Mininni N, Nicolosi GL, Santini M, Schweiger C, Tavazzi L, Tognoni G, Tucci C, Valagussa F Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation* 2002 Apr 23;105:1897-903 [[11997274](#)]

Johansen, 1999:

Johansen O, Brekke M, Seljeflot I, Abdelnoor M, Arnesen H N-3 fatty acids do not prevent restenosis after coronary angioplasty: results from the CART study. Coronary Angioplasty Restenosis Trial. *J Am Coll Cardiol* 1999;33:1619-26 [[10334433](#)]

Nilsen, 2001:

Nilsen DW, Albrektsen G, Landmark K, Moen S, Aarsland T, Woie L Effects of a high-dose concentrate of n-3 fatty acids or corn oil introduced early after an acute myocardial infarction on serum triacylglycerol and HDL cholesterol. *Am J Clin Nutr* 2001 Jul;74:50-6 [[11451717](#)]

Entry terms: omacor, Omacor, Lovaza, omega-3 ethyl ester 90, P-OM3 adjunct,