

Clinical trials of Oxprenolol

TrialResults-center www.trialresultscenter.org

1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
Oxprenolol vs placebo			
CPRG , 1981 n=177/136 follow-up: 8 weeks	-	-	Double
EIS , 1984 n=858/883 follow-up: 1 year	Oxprenolol slow release 160 mg b.i.d. versus placebo	patients 35 to 69 years who has survived acute myocardial infarction	Parallel groups Double blind Europe
Fuccella , 1968 n=106/114 follow-up: 21 days	-	-	Unclear
Lombardo , 1979 n=133/127 follow-up: 20 days	-	-	Double
Schwartz (high risk and low risk) , 1992 n=485/488 follow-up: 22 months (at least 6mo)	Oxprenolol 160mg daily versus placebo	patients surviving MI with or without complication by either ventricular tachycardia or fibrillation	Parallel groups Double blind Italy
Taylor , 1982 n=632/471 follow-up: 48 months	Oxprenolol 40mg twice daily versus placebo	Men 35 to 65 years old who had an acute myocardial infarction between 1 and 90 months reviously	Parallel groups Double blind UK
Wilcox , 1980 n=157/158 follow-up: 6 weeks	-	-	Double

More details and results :

- beta-blockers for acute myocardial infarction in immediate beta-blockers at <http://www.trialresultscenter.org/go-Q41>
- beta-blockers for acute myocardial infarction in long term beta-blockers at <http://www.trialresultscenter.org/go-Q42>

References

CPRG , 1981:

An early intervention secondary prevention study with oxprenolol following myocardial infarction. Eur Heart J 1981 Oct;2:389-93 [7037414]

EIS , 1984:

European Infarction Study (E.I.S.). A secondary prevention study with slow release oxprenolol after myocardial infarction: morbidity and mortality Eur Heart J 1984;5:189-202 [6373264]

Bethge KP, Andresen D, Boissel JP, von Leitner ER, Peyrieux JC, Schroder R, Tietze U Effect of oxprenolol on ventricular arrhythmias: the European Infarction Study experience J Am Coll Cardiol 1985;6:963-72 [2413097]

Fuccella , 1968:

Fuccella LM. Report on the double blind trial with compound CIBA 39089 (trasicor) in myocardial infarction

Lombardo , 1979:

Lombardo M, Selvini A, Belli C, Motolese M, Pedroni P/pj beta-blocking therapy in 440 cases of acute myocardial infarction: a double-blind trial with oxprenolol./1 Proceedings of the Florence international meeting on myocardial infarction 1979;2:803-7

Schwartz (high risk and low risk) , 1992:

Schwartz PJ, Motolese M, Pollavini G, Lotto A, Ruberti U, Trazzi R, et al. Prevention of sudden cardiac death after a first myocardial infarction by pharmacologic or surgical antiadrenergic interventions. J Cardiovasc Electrophysiol 1992;3:2-16.

Taylor , 1982:

Taylor SH, Silke B, Ebbutt A, Sutton GC, Prout BJ, Burley DM A long-term prevention study with oxprenolol in coronary heart disease N Engl J Med 1982;307:1293-301 [6752712]

Wilcox , 1980:

Wilcox RG, Rowley JM, Hampton JR, Mitchell JR, Roland JM, Banks DC Randomised placebo-controlled trial comparing oxprenolol with disopyramide phosphate in immediate treatment of suspected myocardial infarction. Lancet 1980 Oct 11;2:765-9 [6107452]

2 hypertension

Trial	Treatments	Patients	Trials design and methods
oxprenolol vs placebo			
IPPPSH , 1985 n=3185/3172 follow-up: 40y	Oxprenolol versus Placebo	men and women aged 40-64 years with uncomplicated essential hypertension (diastolic blood pressures 100-125 mmHg)	Parallel groups Double blind

More details and results :

- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in uncomplicated hypertension at <http://www.trialresultscenter.org/go-Q685>

References

IPPPSH, 1985:

Cardiovascular risk and risk factors in a randomized trial of treatment based on the beta-blocker oxprenolol: the International Prospective Primary Prevention Study in Hypertension (IPPPSH). The IPPPSH Collaborative Group. *J Hypertens* 1985 Aug;3:379-92 [[2864374](#)]

Entry terms: oxprenolol