

# Clinical trials of UFH + aspirin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 acute coronary syndrome

Trial	Treatments	Patients	Trials design and methods
<b>UFH + aspirin vs placebo</b>			
<b>RISC (ASP+ heparin vs PBO) , 1990</b> n=210/199 follow-up: 1y (5,30 and 90 days)	oral aspirin 75mg/d + intermittent IV heparin 10000UI/d followed by 7500 UI 6-hourly for 4 days versus placebo	men with unstable coronary artery disease (unstable angina or non-Q-wave myocardial infarction)	Sweden
<b>Theroux (heparin+aspirin vs PBO) , 1988</b> n=122/118 follow-up: 3-9 days	heparin (1000 units per hour by intravenous infusion)+ aspirin (325 mg twice daily) versus aspirin (325 mg twice daily)	-	double blind

More details and results :

- antithrombotics for acute coronary syndrome in all type of patients at <http://www.trialresultscenter.org/go-Q24>
- heparin (UFH or LMWH) for acute coronary syndrome in all type of patients at <http://www.trialresultscenter.org/go-Q171>

## References

### **RISC (ASP+ heparin vs PBO), 1990:**

Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary artery disease. The RISC Group. Lancet 1990 Oct 6;336:827-30 [[1976875](#)]

### **Theroux (heparin+aspirin vs PBO), 1988:**

Throux P, Ouimet H, McCans J, Latour JG, Joly P, Lvy G, Pelletier E, Juneau M, Stasiak J, deGuise P Aspirin, heparin, or both to treat acute unstable angina. N Engl J Med 1988;319:1105-11 [[3050522](#)]

Entry terms: UFH