

Clinical trials of UFH

TrialResults-center www.trialresultscenter.org

1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
UFH vs no heparin			
ISIS-2 Pilot , 1987 n=106/103 follow-up: 65279;In-hospital, 1 y (deat	UFH nNo bolus, 1000 IU/h for 48 h versus 65279;No heparin	patients with s65279;uspected MI <=24 h	Parallel groups open
DUCCS , 1994 n=128/122 follow-up: 14 d	UFH no bolus, 15 IU/kg per h for 4 d; target aPTT 5090 s versus No heparin	patients with acute myocardial infarction four hours after APSAC administration, age <=85 y STEMI <=12 h	Parallel groups open
UFH vs placebo			
ECSG , 1992 n=324/320 follow-up: In-hospital	UFH 5000 IU bolus, UFH 1000 IU/h for 48120 h versus Placebo	patients treated with alteplase thrombolysis for acute myocardial infarction, Age 2170 y STEMI <=6h	Parallel groups Double-blind
OSIRIS , 1992 n=64/64 follow-up: In-hospital	UFH 10 000 IU bolus, 1000 IU/h for 24 h versus Placebo	STEMI w=6 h	Parallel groups Double-blind

More details and results :

- antithrombotics for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q36>
- heparin (UFH or LMWH) for acute myocardial infarction in patients eligible to receive fibrinolytic therapy at <http://www.trialresultscenter.org/go-Q308>

References

ISIS-2 Pilot, 1987:

Randomized factorial trial of high-dose intravenous streptokinase, of oral aspirin and of intravenous heparin in acute myocardial infarction. ISIS (International Studies of Infarct Survival) pilot study. Eur Heart J 1987;8:634-42 [2887430]

DUCCS, 1994:

O'Connor CM, Meese R, Carney R, Smith J, Conn E, Burks J, Hartman C, Roark S, Shadoff N, Heard M 3rd A randomized trial of intravenous heparin in conjunction with anistreplase (anisoylated plasminogen streptokinase activator complex) in acute myocardial infarction: the Duke University Clinical Cardiology Study (DUCCS) 1. J Am Coll Cardiol 1994;23:11-8 [8277068]

ECSG, 1992:

de Bono DP, Simoons ML, Tijssen J, Arnold AE, Betriu A, Burgersdijk C, Lopez Bescos L, Mueller E, Pfisterer M, Van de Werf F Effect of early intravenous heparin on coronary patency, infarct size, and bleeding complications after alteplase thrombolysis: results of a randomised double blind European Cooperative Study Group trial. Br Heart J 1992;67:122-8 [1540431]

OSIRIS, 1992:

Col J, Decoster O, Hanique G, Delligne B, Boland J, Pirenne B, Cheron Infusion of heparin conjunct to streptokinase accelerates reperfusion of acute myocardial infarction: results of a double blind randomized study (OSIRIS).6891 Circulation. 1992;86(suppl 1):259a.

2 acute coronary syndrome

2

Trial	Treatments	Patients	Trials design and methods
UFH vs control (on top of aspirin)			
Holdright , 1994 n=154/131 follow-up: hospital stay	intravenous heparin plus oral aspirin (150 mg once daily) versus aspirin alone 150 mg/d	unstable angina	Parallel groups single blind
RISC (heparin+aspirin vs ASP) , 1990 n=210/189 follow-up: 90 days	5 days of intermittent intravenous heparin + oral aspirin 75 mg/day versus oral aspirin 75 mg/day	unstable angina or non-Q-wave myocardial infarction	Parallel groups open
Theroux (heparin+ASP vs ASP) , 1988 n=122/121 follow-up: 3-9 days	aspirin 325 mg/d + heparin 1000 UI/hr IV versus aspirin 325 mg/d		double blind
UFH, warfarin vs control (on top of aspirin)			
ATACS (Cohen) , 1994 n=105/109 follow-up: 12 weeks	aspirin 162.5 mg daily plus heparin (activated partial thromboplastin time, two times control) followed by aspirin 162.5 mg daily plus warfarin (international normalized ratio, 2 to 3) for 12 weeks. versus aspirin alone (162.5 mg daily) for 12 weeks.	patients with unstable rest angina or non-Q-wave myocardial infarction with last episode of pain within 48 hours of randomization and who were nonprior aspirin users	Parallel groups single blind

continued...

Trial	Treatments	Patients	Trials design and methods
Cohen (ATACS pilot) (heparin+aspirin vs asp) , 1990 n=37/32 follow-up: 12 weeks	aspirin (80 mg/day) plus heparin and then warfarin versus aspirin (325 mg/day)	Patients between 21 and 75 years with unstable angina or non-Q-wave MI with last episode of pain within 48 hours of screening.	Parallel groups open
UFH vs placebo			
RISC (heparin vs PBO) , 1990 n=198/199 follow-up: 1y (5,30 and 90 days)	5 days of intermittent intravenous heparin versus placebo	men with unstable coronary artery disease (unstable angina or non-Q-wave myocardial infarction)	Factorial plan Sweden
Theroux (heparin vs PBO) , 1988 n=118/118 follow-up: 3-9 days	heparin (1000 units per hour by intravenous infusion) versus placebo	patients with acute unstable angina pectoris	double blind
UFH + aspirin vs placebo			
RISC (ASP+ heparin vs PBO) , 1990 n=210/199 follow-up: 1y (5,30 and 90 days)	oral aspirin 75mg/d + intermittent IV heparin 10000UI/d followed by 7500 UI 6-hourly for 4 days versus placebo	men with unstable coronary artery disease (unstable angina or non-Q-wave myocardial infarction)	Sweden
Theroux (heparin+aspirin vs PBO) , 1988 n=122/118 follow-up: 3-9 days	heparin (1000 units per hour by intravenous infusion)+ aspirin (325 mg twice daily) versus aspirin (325 mg twice daily)	-	double blind
UFH vs placebo (on top of aspirin)			
Gurfinkel (UFH+aspirin vs aspirin) , 1995 n=70/73 follow-up: 5-7 days	aspirin plus UFH 5000 IU iv then 400 IU/kg body weight per day intravenously and titered by activated partial thromboplastin time versus aspirin 200 mg/day	patients greater than 21 years with unstable angina within 24 hours of randomization	Parallel groups double blind
UFH, warfarin vs aspirin			
Cohen (ATACS pilot) (heparin vs asp) , 1990 n=24/32 follow-up: 12 weeks	heparin followed by warfarin (without aspirin) versus aspirin 325 mg/day	Patients between 21 and 75 years with unstable angina or non-Q-wave MI with last episode of pain within 48 hours of screening	Parallel groups open

More details and results :

- antithrombotics for acute coronary syndrome in all type of patients at <http://www.trialresultscenter.org/go-Q24>
- heparin (UFH or LMWH) for acute coronary syndrome in all type of patients at <http://www.trialresultscenter.org/go-Q171>

References

Holdright, 1994:

Holdright D, Patel D, Cunningham D, Thomas R, Hubbard W, Hendry G, Sutton G, Fox K Comparison of the effect of heparin and aspirin versus aspirin alone on transient myocardial ischemia and in-hospital prognosis in patients with unstable angina. J Am Coll Cardiol 1994;24:39-45 [8006281]

RISC (heparin+aspirin vs ASP), 1990:

Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary artery disease. The RISC Group. Lancet 1990;336:827-30 [1976875]

Theroux (heparin+ASP vs ASP), 1988:

ATACS (Cohen), 1994:

Cohen M, Adams PC, Parry G, Xiong J, Chamberlain D, Wiecek I, Fox KA, Chesebro JH, Strain J, Keller C Combination antithrombotic therapy in unstable rest angina and non-Q-wave infarction in nonprior aspirin users. Primary end points analysis from the ATACS trial. Antithrombotic Therapy in Acute Coronary Syndromes Research Group. Circulation 1994;89:81-8 [8281698]

Cohen (ATACS pilot) (heparin+aspirin vs asp), 1990:

Cohen M, Adams PC, Hawkins L, Bach M, Fuster V Usefulness of antithrombotic therapy in resting angina pectoris or non-Q-wave myocardial infarction in preventing death and myocardial infarction (a pilot study from the Antithrombotic Therapy in Acute Coronary Syndromes Study Group). Am J Cardiol 1990;66:1287-92 [2244556]

RISC (heparin vs PBO), 1990:

Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary artery disease. The RISC Group. Lancet 1990;336:827-30 [1976875]

Theroux (heparin vs PBO), 1988:

Throux P, Ouimet H, McCans J, Latour JG, Joly P, Lvy G, Pelletier E, Juneau M, Stasiak J, deGuise P Aspirin, heparin, or both to treat acute unstable angina. N Engl J Med 1988;319:1105-11 [3050522]

RISC (ASP+ heparin vs PBO), 1990:

Risk of myocardial infarction and death during treatment with low dose aspirin and intravenous heparin in men with unstable coronary artery disease. The RISC Group. Lancet 1990 Oct 6;336:827-30 [1976875]

Theroux (heparin+aspirin vs PBO), 1988:

Throux P, Ouimet H, McCans J, Latour JG, Joly P, Lvy G, Pelletier E, Juneau M, Stasiak J, deGuise P Aspirin, heparin, or both to treat acute unstable angina. N Engl J Med 1988;319:1105-11 [3050522]

Gurfinkel (UFH+aspirin vs aspirin), 1995:

Gurfinkel EP, Manos EJ, Mejia RI, Cerd MA, Duronto EA, Garca CN, Daroca AM, Mautner B Low molecular weight heparin versus regular heparin or aspirin in the treatment of unstable angina and silent ischemia. J Am Coll Cardiol 1995 Aug;26:313-8 [7608429]

Cohen (ATACS pilot) (heparin vs asp), 1990:

Cohen M, Adams PC, Hawkins L, Bach M, Fuster V Usefulness of antithrombotic therapy in resting angina pectoris or non-Q-wave myocardial infarction in preventing death and myocardial infarction (a pilot study from the Antithrombotic Therapy in Acute Coronary Syndromes Study Group). Am J Cardiol 1990;66:1287-92 [2244556]

3 thrombosis prevention

Trial	Treatments	Patients	Trials design and methods
IPC + UFH vs control			
Killewich , 1997 n=NA follow-up:	low-dose heparin sodium therapy (5000 U every 12 hours) and calf-length intermittent mechanical compression devices versus no prophylaxis	patients undergoing aortic reconstruction for aneurysmal or occlusive disease	open
UFH vs control			
Handley , 1972 n=30/30 follow-up: 14 d	Heparin 5000 U IV (loading dose) followed by 20 000 U IV twice daily versus no heparin	Myocardial infarction	Parallel groups
Handley (low dose) , 1972 n=26/24 follow-up: 7 d	Heparin 5000 U IV and 7500 U SC as soon as possible, then 7500 U SC twice daily versus no heparin	Myocardial infarction	Parallel groups
Blech , 1981 n=50/50 follow-up: <=14 days	Unfractionated heparin, 5000 U trice daily, until mobilized versus control	Heart failure, chest infection	Parallel groups open
Gallus , 1973 n=38/40 follow-up: until mobile	heparin 5000 U SC 3 times daily versus no heparin	Myocardial infarction	Parallel groups
Cade n=NA follow-up: <=10 days	-	Age >40, complete bed rest, cardiac failure, obesity, previous VTE, cancer or recent surgery	Parallel groups
Warlow , 1973 n=73/73 follow-up: 10 d	Heparin sodium 5000 U SC twice dailyly versus no heparin	Myocardial infarction	Parallel groups double blind

continued...

Trial	Treatments	Patients	Trials design and methods
Gardlund , 1996 n=5776/5917 follow-up: <=60 days	Unfractionated heparin, 5000 U twice daily, until hospital discharge, <=21 days versus control	Age >55, infectious disease Immobilization	Parallel groups open
Emerson , 1973 n=37/41 follow-up: NA	Heparin Low dose SCium9 versus no heparin	Myocardial infarction	Parallel groups
Zawilska , 1989 n=50/53 follow-up: 14-21 d	Heparin sodium 5000 IU SC twice daily for 14 to 21 days versus no heparin	patients with acute myocardial infarction, disqualified for thrombolytic treatment	Parallel groups
UFH vs no treatment			
Abernethy , 1974 n=63/62	UFH twice daily for 7 days versus no treatment	all other general surgery	
Abraham-Inpijn , 1979 n=20/20	UFH twice daily for 7 days versus no treatment	all other general surgery	
Allen , 1978 n=30/30	UFH twice daily for D versus no treatment	urologic surgery	
Bergqvist , 1979 n=84/77	UFH twice daily for 5 days versus no treatment	elective orthopedic surgery	
Bergqvist , 1979 n=32/23	UFH twice daily for 5 days versus no treatment	traumatic orthopedic surgery	
Bergqvist , 1980 n=53/58	UFH twice daily for 5 days versus no treatment	all other general surgery	
Caloghera , 1984 n=40/40	UFH 3 times daily for 7 days versus no treatment	all other general surgery	
Clarke-Pearson , 1983 n=95/105	UFH twice daily for 7 days versus no treatment	all other general surgery	
Coe , 1978 n=28/24	UFH twice daily for D versus no treatment	urologic surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
Dechavanne , 1974 n=29/29	UFH 3 times daily for 10 days versus no treatment	elective orthopedic surgery	
Dechavanne , 1975 n=20/21	UFH 3 times daily for 10 days versus no treatment	elective orthopedic surgery	
Galasko , 1976 n=50/50	UFH twice daily (duration unknown) versus no treatment	traumatic orthopedic surgery	
Gallus , 1973 n=23/23	UFH 3 times daily for A versus no treatment	traumatic orthopedic surgery	
Gallus , 1973 n=8/9	UFH twice daily for A days versus no treatment	elective orthopedic surgery	
Gallus , 1973 n=100/109	UFH 3 times daily for A versus no treatment	all other general surgery	
Gallus , 1976 n=408/412	UFH 3 times daily for 7 days versus no treatment	all other general surgery	
Gordon-Smith , 1972 n=NA	UFH twice daily for 5 days versus no treatment	all other general surgery	
Groote Schuur , 1979 n=323/323	UFH 3 times daily for 6 days versus no treatment	all other general surgery	
Gruber n=119/113	UFH 3 times daily for 7 days versus no treatment	All other general surgery	
Kakkar (IMT) n=2111/2137	UFH 3 times daily for 7 days versus no treatment	All other general surgery	
Kettunen , 1974 n=83/117	UFH 3 times daily for 7 days versus no treatment	all other general surgery	
Kopenhagen , 1982 n=162/50	UFH twice daily versus no treatment	all other general surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
Kraytman , 1976 n=21/28	UFH 3 times daily for 6 days versus no treatment	all other general surgery	
Kraytman , 1977 n=23/27	UFH 3 times daily for 6 days versus no treatment	all other general surgery	
Mannucci , 1976 n=23/24	UFH 3 times daily for 7 days versus no treatment	elective orthopedic surgery	
Marchelli , 1983 n=30/30	UFH 3 times daily for ? versus no treatment	all other general surgery	
Morris , 1974 n=36/36	UFH twice daily for 10 days versus no treatment	elective orthopedic surgery	
Morris , 1977 n=24/24	UFH 3 times daily for 10 days versus no treatment	traumatic orthopedic surgery	
Multiunit , 1974 n=128/128	UFH twice daily for 7 days versus no treatment	all other general surgery	
Nicolaides , 1972 n=128/122	UFH twice daily for 7 days versus no treatment	all other general surgery	
Plante , 1979 n=42/66	UFH twice daily for 8 days versus no treatment	all other general surgery	
Sagar , 1974 n=264/236	UFH twice daily for 5 days versus no treatment	all other general surgery	
Sebeseri , 1975 n=34/31	UFH twice daily for D versus no treatment	urologic surgery	
Spebar , 1981 n=24/19	UFH twice daily for 5-7 days versus no treatment	all other general surgery	
VTCSG , 1975 n=34/30	UFH twice daily for 10 days versus no treatment	elective orthopedic surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
Welin-Berger , 1982 n=20/20	UFH twice daily for 7 days versus no treatment	elective orthopedic surgery	
Ziemski , 1979 n=30/20	UFH 3 times daily for 2 days versus no treatment	all other general surgery	
Cerrato , 1978 n=50/50 follow-up: NA	Unfractionated heparin, 5000 IU subcutaneously 3 times a day, starting 2 hours preoperatively for +7 days versus no treatment	Elective neurosurgery for tumors, 40 years or older	
UFH vs placebo			
Abraham-Inpijn , 1975 n=12/13	UFH twice daily for 8 days versus placebo	elective orthopedic surgery	
Bejjani , 1983 n=17/17	UFH twice daily for 2 days versus placebo	urologic surgery	
Belch , 1980 n=24/25	UFH twice daily for 7 days versus placebo	all other general surgery	
Pitt , 1980 n=36/37 follow-up: 2-3 d	Heparin 500 U IV twice daily versus placebo	patients with acute myocardial infarction of less than 48 hours duration	Parallel groups open
Covey , 1975 n=53/52	UFH twice daily for 8 days versus placebo	all other general surgery	
Hampson , 1974 n=48/52	UFH 3 times daily for 7- 10 days versus placebo	elective orthopedic surgery	
Hedlund , 1979 n=30/29	UFH 3 times daily for 5 days versus placebo	urologic surgery	
Jourdan , 1984 n=21/21	UFH twice daily for 5 days versus placebo	all other general surgery	
Kakkar , 1972 n=39/39	UFH twice daily for 7 days versus placebo	All other general surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
Kiil , 1978 n=650/663	UFH twice daily for 7 days versus placebo	All other general surgery	
Kraytman , 1977 n=25/22	UFH 3 times daily for 6 days versus placebo	all other general surgery	
Kruse-Blinkenberg , 1980 n=29/33	UFH 3 times daily for 8 days versus placebo	all other general surgery	
Kutnowski , 1977 n=6/13	UFH 3 times daily for 6 days versus placebo	urologic surgery	
Lahnborg , 1975 n=58/54	UFH twice daily for 5 days versus placebo	all other general surgery	
Lahnborg , 1976 n=24/24	UFH twice daily for 5 days versus placebo	all other general surgery	
Lahnborg , 1980 n=70/69	UFH twice daily for 10 days versus placebo	traumatic orthopedic surgery	
Lawrence , 1977 n=133/129	UFH twice daily for 5 days versus placebo	all other general surgery	
Lowe , 1981 n=51/49	UFH twice daily for 16 days versus placebo	elective orthopedic surgery	
Marchetti , 1982 n=16/13	UFH 3 times daily versus placebo	all other general surgery	
Moskovitz , 1978 n=29/23	UFH 3 times daily for 7 days versus placebo	traumatic orthopedic surgery	
Moskovitz , 1978 n=35/32	UFH 3 times daily for 7 days versus placebo	elective orthopedic surgery	
Muiticentre , 1984 n=222/108	UFH twice daily for 5-7 days versus placebo	all other general surgery	

continued...

Trial	Treatments	Patients	Trials design and methods
Ribaudo , 1975 n=75/75	UFH twice daily for A days versus placebo	all other general surgery	
Roberts , 1975 n=39/45	UFH twice daily for 7 days versus placebo	all other general surgery	
Strand , 1975 n=55/55	UFH twice daily for 7 days versus placebo	all other general surgery	
Svend-Hansen , 1981 n=65/65	UFH 3 times daily for 14 days versus placebo	traumatic orthopedic surgery	
Taberner , 1978 n=50/50	UFH twice daily for 7 days versus placebo	all other general surgery	
Torngren n=47/62	UFH twice daily for 6-8 days versus placebo	all other general surgery	
Torngren , 1978 n=10/10	UFH 3 times daily for 6-8 days versus placebo	all other general surgery	
Torngren , 1978 n=66/62	UFH twice daily for 6-8 days versus placebo	all other general surgery	
Vandendris , 1980 n=32/33	UFH 3 times daily for 6 days versus placebo	urologic surgery	
Williams , 1978 n=106/106	UFH twice daily for 14 days versus placebo	elective orthopedic surgery	
Wu , 1977 n=44/44	UFH twice daily for 7 days versus placebo	all other general surgery	
Xabregas , 1977 n=27/26	UFH twice daily for 14 days versus placebo	traumatic orthopedic surgery	
IPC + UFH vs UFH			

continued...

Trial	Treatments	Patients	Trials design and methods
Ramos , 1996 n=1355/1196 follow-up:	prophylactic regimen of Pneumatic compression stocking and subcutaneous heparin versus with 5,000 U of subcutaneous heparin every 12 h	cardiac surgery	Parallel groups open
Siragusa (H) , 1994 n=35/35 follow-up:	-	-	Parallel groups open
UFH vs aspirin			
Loew , 1977 n=60/67	UFH twice daily for 7 days versus aspirin	all other general surgery	
Vinazzer , 1980 n=402/404	UFH twice daily for A days versus aspirin	all other general surgery	
UFH vs DHE			
Muiticentre , 1984 n=226/110	UFH twice daily for 5-7 days versus DHE	all other general surgery	
Muiticentre 53,S4 , 1984 n=214/110	UFH twice daily for 5-7 days versus DHE	all other general surgery	
UFH vs oral anticoagulant			
van Geloven , 1977 n=74/80	UFH twice daily for 4 days versus Oral anticoagulant	all other general surgery	
out of hospital UFH vs standard prophylaxis			
Manganelli , 1998 n=79/80	65279;in hospital thromboprophylaxis followed by out of hospital UFH 5000 IU three times a day for a total duration of 30 days versus UFH 5000 IU three times a day for 15 days	THR	
GCS + UFH vs UFH			

continued...

Trial	Treatments	Patients	Trials design and methods
*Rasmussen (adj) , 1988 n=NA follow-up:	subcutaneous heparin and graduated compression stockings to the knee (TED stockings), versus subcutaneous heparin	patients (age more than 40 yrs) admitted for major abdominal surgery	Parallel groups open
Torngren , 1980 n=NA	-	-	
Wille-Jorgensen , 1985 n=86/90 follow-up:	low-dose heparin treatment with graded compression stockings versus low-dose heparin treatment (5000 units twice daily subcutaneously)	patients undergoing Abdominal surgery	parallel groups
Wille-Jorgensen , 1991 n=83/83 follow-up:	low dose heparin and graded compression stockings versus low dose heparin	patients undergoing Abdominal surgery	parallel groups

More details and results :

- antithrombotics for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q37>
- antithrombotics for thrombosis prevention in medical patients at <http://www.trialresultscenter.org/go-Q87>
- antithrombotics for thrombosis prevention in general surgery at <http://www.trialresultscenter.org/go-Q92>
- antithrombotics for thrombosis prevention in urologic surgery at <http://www.trialresultscenter.org/go-Q93>
- antithrombotics for thrombosis prevention in neurosurgery at <http://www.trialresultscenter.org/go-Q99>
- graduated compression stockings for thrombosis prevention in all type of patients at <http://www.trialresultscenter.org/go-Q158>
- antiplatelets drug for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q186>
- UFH for thrombosis prevention in general surgery at <http://www.trialresultscenter.org/go-Q191>
- UFH for thrombosis prevention in all type of surgery at <http://www.trialresultscenter.org/go-Q192>
- UFH for thrombosis prevention in urologic surgery at <http://www.trialresultscenter.org/go-Q194>
- heparin (UFH or LMWH) for thrombosis prevention in neurosurgery at <http://www.trialresultscenter.org/go-Q197>
- LMWH for thrombosis prevention in neurosurgery at <http://www.trialresultscenter.org/go-Q198>

- mechanical devices for thromboprophylaxis for thrombosis prevention in all type of patients at <http://www.trialresultscenter.org/go-Q402>
- antithrombotics for thrombosis prevention in patients with myocardial infarction at <http://www.trialresultscenter.org/go-Q406>
- mechanical devices for thromboprophylaxis for thrombosis prevention in general non orthopaedic surgery at <http://www.trialresultscenter.org/go-Q466>

References

Killewich, 1997:

Killewich LA, Aswad MA, Sandager GP, Lilly MP, Flinn WR A randomized, prospective trial of deep venous thrombosis prophylaxis in aortic surgery. Arch Surg 1997;132:499-504 [9161392]

Handley, 1972:

Handley AJ, Emerson PA, Fleming PR Heparin in the prevention of deep vein thrombosis after myocardial infarction. Br Med J 1972;2:436-8 [4555651]

Handley (low dose), 1972:

Handley AJ Low-dose heparin after myocardial infarction. Lancet 1972;2:623-4 [4116776]

Blech, 1981:

Belch JJ, Lowe GD, Ward AG, Forbes CD, Prentice CR Prevention of deep vein thrombosis in medical patients by low-dose heparin. Scott Med J 1981;26:115-7 [7291971]

Gallus, 1973:

Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carroll JJ, Minden JH, Hudecki SM Small subcutaneous doses of heparin in prevention of venous thrombosis. N Engl J Med 1973;288:545-51 [4568221]

Cade, 0:

Cade JF, Andrews JT, Stubbs AE Comparison of sodium and calcium heparin in prevention of venous thromboembolism. Aust N Z J Med 1982;12:501-4 [6758747]

Warlow, 1973:

Warlow C, Terry G, Kenmure AC, Beattie AG, Ogston D, Douglas AS A double-blind trial of low doses of subcutaneous heparin in the prevention of deep-vein thrombosis after myocardial infarction. Lancet 1973;2:934-6 [4126560]

Gardlund, 1996:

Grdlund B Randomised, controlled trial of low-dose heparin for prevention of fatal pulmonary embolism in patients with infectious diseases. The Heparin Prophylaxis Study Group. Lancet 1996;347:1357-61 [8637340]

Emerson, 1973:

Emerson PA, Marks P Preventing thromboembolism after myocardial infarction: effect of low-dose heparin or smoking. Br Med J 1977;1:18-20 [831965]

Zawilska, 1989:

Zawilska K, Psuja P, Lewandowski K, Wrz M Low-dose heparin in the prevention of thrombotic complications following acute myocardial infarction. Cor Vasa 1989;31:179-85 [2670447]

Abernethy, 1974:

Abernethy EA, Hartsuck JM Postoperative pulmonary embolism. A prospective study utilizing low dose heparin. Am J Surg 1974;128:739-42 [4432994]

Abraham-Inpijn, 1979:

Abraham-Inpijn L Critical evaluation of low-dose heparin in laryngectomy. Arch Chir Neerl 1979;31:9-15 [443830]

Allen, 1978:

Allen NH, Jenkins JD, Smart CJ Surgical haemorrhage in patients given subcutaneous heparin as prophylaxis against thromboembolism. Br Med J 1978;1:1326 [647262]

Bergqvist , 1979:

Bergqvist D, Efsing HO, Hallbk T, Hedlund T Thromboembolism after elective and post-traumatic hip surgery—a controlled prophylactic trial with dextran 70 and low-dose heparin. Acta Chir Scand 1979;145:213-8 [386676]

Bergqvist, 1979:

Bergqvist D, Efsing HO, Hallbk T, Hedlund T Thromboembolism after elective and post-traumatic hip surgery—a controlled prophylactic trial with dextran 70 and low-dose heparin. Acta Chir Scand 1979;145:213-8 [386676]

Bergqvist, 1980:

Bergqvist D, Hallbk T Prophylaxis of postoperative venous thrombosis in a controlled trial comparing dextran 70 and low-dose heparin. World J Surg 1980;4:239-43 [6157256]

Caloghera, 1984:

Caloghera C, Bordos D, Miculit F, Aboubakr W, Teodorescu C, Vancea D [Prevention of postoperative thromboembolism with small doses of heparin] Rev Chir Oncol Radiol O R L Oftalmol Stomatol Chir 1984;33:161-7 [6238351]

Clarke-Pearson, 1983:

Clarke-Pearson DL, Coleman RE, Synan IS, Hinshaw W, Creasman WT Venous thromboembolism prophylaxis in gynecologic oncology: a prospective, controlled trial of low-dose heparin. Am J Obstet Gynecol 1983;145:606-13 [6338722]

Coe, 1978:

Coe NP, Collins RE, Klein LA, Bettmann MA, Skillman JJ, Shapiro RM, Salzman EW Prevention of deep vein thrombosis in urological patients: a controlled, randomized trial of low-dose heparin and external pneumatic compression boots. Surgery 1978;83:230-4 [622696]

Dechavanne, 1974:

Dechavanne M, Saudin F, Viala JJ, Kher A, Bertrix L, de Mourgues G [Prevention of venous thrombosis. Success of high doses of heparin during total hip replacement for osteoarthritis] Nouv Presse Med 1974;3:1317-9 [4843085]

Dechavanne, 1975:

Dechavanne M, Ville D, Viala JJ, Kher A, Faivre J, Pousset MB, Dejour H Controlled trial of platelet anti-aggregating agents and subcutaneous heparin in prevention of postoperative deep vein thrombosis in high risk patients. Haemostasis 1975;4:94-100 [1205340]

Galasko , 1976:

Galasko CS, Edwards DH, Fearn CB, Barber HM The value of low dosage heparin for the prophylaxis of thromboembolism in patients with transcervical and intertrochanteric femoral fractures. Acta Orthop Scand 1976;47:276-82 [782146]

Gallus , 1973:

Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carroll JJ, Minden JH, Hudecki SM Small subcutaneous doses of heparin in prevention of venous thrombosis. N Engl J Med 1973;288:545-51 [4568221]

Gallus , 1973:

Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carroll JJ, Minden JH, Hudecki SM Small subcutaneous doses of heparin in prevention of venous thrombosis. N Engl J Med 1973;288:545-51 [4568221]

Gallus, 1973:

Gallus AS, Hirsh J, Tuttle RJ, Trebilcock R, O'Brien SE, Carroll JJ, Minden JH, Hudecki SM Small subcutaneous doses of heparin in prevention of venous thrombosis. N Engl J Med 1973;288:545-51 [4568221]

Gallus, 1976:

Gallus AS, Hirsh J, O'Brien SE, McBride JA, Tuttle RJ, Gent M Prevention of venous thrombosis with small, subcutaneous doses of heparin. JAMA 1976;235:1980-2 [946507]

Gordon-Smith, 1972:

Gordon-Smith IC, Le Quesne LP, Grundy DJ, Newcombe JF, Bramble FJ Controlled trial of two regimens of subcutaneous heparin in prevention of postoperative deep-vein thrombosis. Lancet 1972;1:1133-5 [4113052]

Groote Schuur, 1979:

Failure of low-dose heparin to prevent significant thromboembolic complications in high-risk surgical patients: interim report of prospective trial. Groote Schuur Hospital Thromboembolus Study Group. Br Med J 1979;1:1447-50 [380742]

Gruber, 0:

Gruber UF, Duckert F, Fridrich R, Torhorst J, Rem J Prevention of postoperative thromboembolism by dextran 40, low doses of heparin, or xantinol nicotinate. Lancet 1977;1:207-10 [64746]

Kakkar (IMT), 0:

Prevention of fatal postoperative pulmonary embolism by low doses of heparin. An international multicentre trial. Lancet 1975;2:45-51 [49649]

Kakkar VV, Corrigan TP, Fossard DP, Sutherland I, Thirwell J Prevention of Fatal Postoperative pulmonary embolism by low doses of heparin. Reappraisal of results of international multicentre trial. Lancet 1977;1:567-9 [65660]

Kettunen, 1974:

Kettunen K, Poikolainen E, Karjalainen P, Oksala I, Alhava E, Rehnberg V, Huttunen H, Mattila M [Prevention of postoperative deep vein thrombosis with small doses of heparin] Duodecim 1974;90:834-8 [4426267]

Kopenhagen, 1982:

Kopenhagen K, Matthes M [Heparin-dihydergot or heparin alone in thrombosis prophylaxis?] Med Welt 1982;33:216-23 [7070283]

Kraytman, 1976:

Kraytman M, Kutnowski M, Ansay J, Fastrez R [Prophylaxis of postoperative deep vein thromboses by means of weak doses of subcutaneous heparin (author's transl)] Acta Chir Belg 1976;75:519-29 [1015181]

Kraytman, 1977:

Kraytman M, Kutnowski M, Ansay J, Vandendris M, Fastrez R, Steinberger R [Prevention of postoperative venous thrombosis with small doses of subcutaneous heparin] Acta Clin Belg 1977;32:422-7 [358713]

Mannucci , 1976:

Mannucci PM, Citterio LE, Panajotopoulos N Low-dose heparin and deep-vein thrombosis after total hip replacement. Thromb Haemost 1976;36:157-64 [1036806]

Marchelli, 1983:

Marchetti V, Beati C, Pogliani EM, Vincere G [Low-dose calcium-heparin prophylaxis in thoracic surgery. Bleeding, changes in coagulation and fibrinolysis] *Minerva Med* 1983;74:1745-8 [6866306]

Morris, 1974:

Morris GK, Henry AP, Preston BJ Prevention of deep-vein thrombosis by low-dose heparin in patients undergoing total hip replacement. *Lancet* 1974;2:797-800 [4138248]

Morris , 1977:

Morris GK, Mitchell JR Preventing venous thromboembolism in elderly patients with hip fractures: studies of low-dose heparin, dipyridamole, aspirin, and flurbiprofen. *Br Med J* 1977;1:535-7 [843794]

Multiunit, 1974:

Heparin versus dextran in the prevention of deep-vein thrombosis. A multi-unit controlled trial. *Lancet* 1974;2:118-20 [4135556]

Nicolaides, 1972:

Nicolaides AN, Dupont PA, Desai S, Lewis JD, Douglas JN, Dodsworth H, Fourides G, Luck RJ, Jamieson CW Small doses of subcutaneous sodium heparin in preventing deep venous thrombosis after major surgery. *Lancet* 1972;2:890-3 [4116600]

Plante, 1979:

Plante J, Boneu B, Vaysse C, Barret A, Gouzi M, Bierme R Dipyridamole-aspirin versus low doses of heparin in the prophylaxis of deep venous thrombosis in abdominal surgery. *Thromb Res* 1979;14:399-403 [442014]

Sagar, 1974:

Sagar S Heparin prophylaxis against fatal postoperative pulmonary embolism. *Br Med J* 1974;2:153-5 [4825114]

Plante J, Boneu B, Vaysse C, Barret A, Gouzi M, Bierme R Dipyridamole-aspirin versus low doses of heparin in the prophylaxis of deep venous thrombosis in abdominal surgery. *Thromb Res* 1979;14:399-403 [442014]

Sebeseri, 1975:

Sebeseri O, Kummer H, Zingg E Controlled prevention of post-operative thrombosis in urological diseases with depot heparin. *Eur Urol* 1975;1:229-30 [1233188]

Spebar, 1981:

Spebar MJ, Collins GJ Jr, Rich NM, Kang IY, Clagett GP, Salander JM Perioperative heparin prophylaxis of deep venous thrombosis in patients with peripheral vascular disease. *Am J Surg* 1981;142:649-50 [7316026]

VTCSG, 1975:

Small doses of subcutaneous sodium heparin in the prevention of deep vein thrombosis after elective hip operations. *Br J Surg* 1975;62:348-50 [1139128]

Welin-Berger, 1982:

Welin-Berger T, Bygdeman S, Mebius C Deep vein thrombosis following hip surgery. Relation to activated factor X inhibitor activity: effect of heparin and dextran. *Acta Orthop Scand* 1982;53:937-45 [6184938]

Ziemski, 1979:

Ziemski JM, Kostrzewska E, Marchlewski S, Wiczorek K, Rudowski W, Michalski R, Lopaciuk S, Kolakowski L, Sawicki F [Efficacy of small doses of heparin given during 2 to 6 days in the prevention of postoperative deep vein thrombosis] *Pol Tyg Lek* 1979;34:161-4 [432128]

Cerrato, 1978:

Cerrato D, Ariano C, Fiacchino F Deep vein thrombosis and low-dose heparin prophylaxis in neurosurgical patients. *J Neurosurg* 1978;49:378-81 [681999]

Abraham-Inpijn , 1975:

Abraham-Inpijn L, Vreeken J Effect of low-dose heparin on incidence of postoperative thrombosis in orthopaedic patients. Arch Chir Neerl 1975;27:63-8 [1098580]

Bejjani, 1983:

Bejjani BB, Chen DC, Nolan NG, Edson M Minidose heparin in transurethral prostatectomy. Urology 1983;22:251-4 [6623770]

Belch, 1980:

Belch JJ, Lowe GD, Pollock JG, Forbes CD, Prentice CR Low dose heparin in the prevention of deep-vein thrombosis after aortic bifurcation graft surgery. Thromb Haemost 1980;42:1429-33 [7368149]

Pitt, 1980:

Pitt A, Anderson ST, Habersberger PG, Rosengarten DS Low dose heparin in the prevention of deep-vein thromboses in patients with acute myocardial infarction. Am Heart J 1980;99:574-8 [7369096]

Covey, 1975:

Covey TH, Sherman L, Baue AE Low-dose heparin in postoperative patients: a prospective, coded study. Arch Surg 1975;110:1021-6 [1100014]

Hampson, 1974:

Hampson WG, Harris FC, Lucas HK, Roberts PH, McCall IW, Jackson PC, Powell NL, Staddon GE Failure of low-dose heparin to prevent deep-vein thrombosis after hip-replacement arthroplasty. Lancet 1974;2:795-7 [4138905]

Hedlund, 1979:

Hedlund PO, Blombck M The effect of prophylaxis with low dose heparin on blood coagulation parameters. A double blind study in connection with transvesical prostatectomy. Thromb Haemost 1979;41:337-45 [382433]

Hedlund PO, Blombck M The effects of low-dose heparin treatment on patients undergoing transvesical prostatectomy. Urol Res 1981;9:147-52 [7257029]

Jourdan, 1984:

Jourdan M, McColl I The use of prophylactic subcutaneous heparin in patients undergoing hernia repairs. Br J Clin Pract 1984;38:298-300 [6477809]

Kakkar, 1972:

Kakkar VV, Corrigan T, Spindler J, Fossard DP, Flute PT, Crellin RQ, Wessler S, Yin ET Efficacy of low doses of heparin in prevention of deep-vein thrombosis after major surgery. A double-blind, randomised trial. Lancet 1972;2:101-6 [4113891]

Kiil, 1978:

Kiil J, Kiil J, Axelsen F, Andersen D Prophylaxis against postoperative pulmonary embolism and deep-vein thrombosis by low-dose heparin. Lancet 1978;1:1115-6 [77413]

Kraytman, 1977:

Kraytman M, Kutnowski M, Ansay J, Vandendris M, Fastrez R, Steinberger R [Prevention of postoperative venous thrombosis with small doses of subcutaneous heparin] Acta Clin Belg 1977;32:422-7 [358713]

Kruse-Blinkenberg, 1980:

Kruse-Blinkenberg HO, Gormsen J The influence of low dose heparin in elective surgery on blood coagulation, fibrinolysis, platelet function, antithrombin III and antiplasmin. Acta Chir Scand 1980;146:375-82 [7008475]

Kutnowski, 1977:

Kraytman M, Kutnowski M, Ansay J, Vandendris M, Fastrez R, Steinberger R [Prevention of postoperative venous thrombosis with small doses of subcutaneous heparin] Acta Clin Belg 1977;32:422-7 [358713]

Lahnborg, 1975:

Leter: The consultants' contract. *Lancet* 1975;1:329-30 [46465]

Lahnborg G, Bergström K Clinical and haemostatic parameters related to thromboembolism and low-dose heparin prophylaxis in major surgery. *Acta Chir Scand* 1975;141:590-5 [55029]

Lahnborg, 1976:

Lahnborg G, Lagergren H, Hedenstierna G Effect of low-dose heparin prophylaxis on arterial oxygen tension after high laparotomy. *Lancet* 1976;1:54-6 [54579]

Lahnborg, 1980:

Moreno-Gonzalez E, Sanmartin JH, Pascual MH, Moreno-Azcoita M, Selas PR Total esophagectomy by right anterior thoracotomy and immediate esophageal reconstruction for carcinoma of the esophagus. *Acta Chir Scand* 1980;146:19-23 [7376779]

Lahnborg G Effect of low-dose heparin and dihydroergotamine on frequency of postoperative deep-vein thrombosis in patients undergoing post-traumatic hip surgery. *Acta Chir Scand* 1980;146:319-22 [7468062]

Lawrence, 1977:

Lawrence JC, Xabregas A, Gray L, Ham JM Seasonal variation in the incidence of deep vein thrombosis. *Br J Surg* 1977;64:777-80 [338089]

Lowe , 1981:

Lowe LW Venous thrombosis and embolism. *J Bone Joint Surg Br* 1981;63-B:155-67 [6163785]

Marchetti, 1982:

Kopenhagen K, Matthes M [Heparin-dihydroergot or heparin alone in thrombosis prophylaxis?] *Med Welt* 1982;33:216-23 [7070283]

Moskovitz, 1978:

Moskovitz PA, Ellenberg SS, Feffer HL, Kenmore PI, Neviasser RJ, Rubin BE, Varma VM Low-dose heparin for prevention of venous thromboembolism in total hip arthroplasty and surgical repair of hip fractures. *J Bone Joint Surg Am* 1978;60:1065-70 [363722]

Moskovitz, 1978:

Moskovitz PA, Ellenberg SS, Feffer HL, Kenmore PI, Neviasser RJ, Rubin BE, Varma VM Low-dose heparin for prevention of venous thromboembolism in total hip arthroplasty and surgical repair of hip fractures. *J Bone Joint Surg Am* 1978;60:1065-70 [363722]

Muiticentre, 1984:

Dihydroergotamine-heparin prophylaxis of postoperative deep vein thrombosis. A multicenter trial. The Multicenter Trial Committee. *JAMA* 1984;251:2960-6 [6371278]

DiSerio FJ, Sasahara AA United States trial of dihydroergotamine and heparin prophylaxis of deep vein thrombosis. *Am J Surg* 1985;150:25-32 [3901790]

Ribaudo, 1975:

Ribaudo JM, Hoellrich RG, McKinnon WM, Shuler SE Evaluation of mini-dose heparin administration as a prophylaxis against postoperative pulmonary embolism: a prospective double-blind study. *Am Surg* 1975;41:289-95 [1093461]

Ribaudo JM, Hoellrich RG, McKinnon WM, Shuler SE Evaluation of mini-dose heparin administration as a prophylaxis against postoperative pulmonary embolization: a prospective double-blind study. *Rev Surg* 1975;32:297-9 [1198038]

Roberts, 1975:

Roberts VC, Cotton LT Failure of low-dose heparin to improve efficacy of peroperative intermittent calf compression in preventing postoperative deep vein thrombosis. *Br Med J* 1975;3:458-60 [1098726]

Strand, 1975:

Strand L, Bank-Mikkelsen OK, Lindewald H Small heparin doses as prophylaxis against deep-vein thrombosis in major surgery. *Acta Chir Scand* 1975;141:624-7 [1108565]

Svend-Hansen, 1981:

Svend-Hansen H, Bremerskov V, Gtrik J, Ostri P Low-dose heparin in proximal femoral fractures. Failure to prevent deep-vein thrombosis. *Acta Orthop Scand* 1981;52:77-80 [7010893]

Taberner, 1978:

Taberner DA, Poller L, Burslem RW, Jones JB Oral anticoagulants controlled by the British comparative thromboplastin versus low-dose heparin in prophylaxis of deep vein thrombosis. *Br Med J* 1978;1:272-4 [340005]

Torngren, 0:

Torngren, 1978:

Trngren S, Forsberg K Concentrated or diluted heparin prophylaxis of postoperative deep venous thrombosis. *Acta Chir Scand* 1978;144:283-8 [367024]

Torngren, 1978:

Trngren S, Forsberg K Concentrated or diluted heparin prophylaxis of postoperative deep venous thrombosis. *Acta Chir Scand* 1978;144:283-8 [367024]

Vandendris, 1980:

Vandendris M, Kutnowski M, Futeral B, Gianakopoulos X, Kraytman M, Gregoir W Prevention of postoperative deep-vein thrombosis by low-dose heparin in open prostatectomy. *Urol Res* 1980;8:219-21 [7013228]

Kraytman M, Kutnowski M, Ansay J, Vandendris M, Fastrez R, Steinberger R [Prevention of postoperative venous thrombosis with small doses of subcutaneous heparin] *Acta Clin Belg* 1977;32:422-7 [358713]

Williams , 1978:

Williams JW, Eikman EA, Greenberg SH, Hewitt JC, Lopez-Cuenca E, Jones GP, Madden JA Failure of low dose heparin to prevent pulmonary embolism after hip surgery or above the knee amputation. *Ann Surg* 1978;188:468-74 [697431]

Wu, 1977:

Wu TK, Tsapogas MJ, Jordan FR Prophylaxis of deep venous thrombosis by hydroxychloroquine sulfate and heparin. *Surg Gynecol Obstet* 1977;145:714-8 [333643]

Xabregas , 1977:

Xabregas A, Gray L, Ham JM Heparin prophylaxis of deep vein thrombosis in patients with a fractured neck of the femur. *Med J Aust* 1978;1:620-2 [355810]

Lawrence JC, Xabregas A, Gray L, Ham JM Seasonal variation in the incidence of deep vein thrombosis. *Br J Surg* 1977;64:777-80 [338089]

Ramos, 1996:

Ramos R, Salem BI, De Pawlikowski MP, Coords C, Eisenberg S, Leidenfrost R The efficacy of pneumatic compression stockings in the prevention of pulmonary embolism after cardiac surgery. *Chest* 1996 Jan;109:82-5 [8549224]

Siragusa (H), 1994:

Siragusa S, Vicentini L, Carbone S, Barone M, Beltrametti C, Piovella F. Intermittent pneumatic leg compression (IPLC) and unfractionated heparin (UFH) in the prevention of post-operative deep vein thrombosis in hip surgery: a randomized clinical trial *Br J Haematol* 1994; 87:186.

Loew, 1977:

Loew D, Brcke P, Simma W, Vinazzer H, Dienstl E, Boehme K Acetylsalicylic acid, low dose heparin, and a combination of both substances in the prevention of postoperative thromboembolism. A double blind study. *Thromb Res* 1977;11:81-6 [329468]

Vinazzer, 1980:

Vinazzer H, Loew D, Simma W, Brcke P Prophylaxis of postoperative thromboembolism by low dose heparin and by acetylsalicylic acid given simultaneously. A double blind study. *Thromb Res* 1980;17:177-84 [7376128]

Muiticentre, 1984:

Dihydroergotamine-heparin prophylaxis of postoperative deep vein thrombosis. A multicenter trial. The Multicenter Trial Committee. *JAMA* 1984;251:2960-6 [6371278]

DiSerio FJ, Sasahara AA United States trial of dihydroergotamine and heparin prophylaxis of deep vein thrombosis. *Am J Surg* 1985;150:25-32 [3901790]

Muiticentre 53,S4, 1984:

Dihydroergotamine-heparin prophylaxis of postoperative deep vein thrombosis. A multicenter trial. The Multicenter Trial Committee. *JAMA* 1984;251:2960-6 [6371278]

DiSerio FJ, Sasahara AA United States trial of dihydroergotamine and heparin prophylaxis of deep vein thrombosis. *Am J Surg* 1985;150:25-32 [3901790]

van Geloven, 1977:

van Geloven F, Wittebol P, Sixma JJ Comparison of postoperative coumarin, dextran 40 and subcutaneous heparin in the prevention of postoperative deep vein thrombosis. *Acta Med Scand* 1977;202:367-72 [335793]

Manganelli, 1998:

Manganelli D, Pazzagli M, Mazzantini D, Punzi G, Manca M, Vignali C, Palla A, Troiani R, Rossi G Prolonged prophylaxis with unfractionated heparin is effective to reduce delayed deep vein thrombosis in total hip replacement. *Respiration* 1998;65:369-74 [9782219]

***Rasmussen (adj), 1988:**

Rasmussen A, Hansen PT, Lindholt J, Poulsen TD, Toftdahl DB, Gram J, Toftgaard C, Jespersen J Venous thrombosis after abdominal surgery. A comparison between subcutaneous heparin and antithrombotic stockings, or both. *J Med* 1988;19:193-201 [2972790]

Torngren, 1980:

Wille-Jorgensen, 1985:

Wille-Jrgensen P, Thorup J, Fischer A, Holst-Christensen J, Flamsholt R Heparin with and without graded compression stockings in the prevention of thromboembolic complications of major abdominal surgery: a randomized trial. *Br J Surg* 1985;72:579-81 [4016545]

Wille-Jorgensen, 1991:

Wille-Jrgensen P, Hauch O, Dimo B, Christensen SW, Jensen R, Hansen B Prophylaxis of deep venous thrombosis after acute abdominal operation. *Surg Gynecol Obstet* 1991;172:44-8 [1702235]

4 superficial thrombophlebitis

Trial	Treatments	Patients	Trials design and methods
Prophylactic UFH vs no treatment			
Belcaro (prophylactic UFH vs no) , 1999 n=NA	-	-	

More details and results :

- antithrombotics for superficial thrombophlebitis in superficial thrombophlebitis of the leg at <http://www.trialresultscenter.org/go-Q218>

References

Belcaro (prophylactic UFH vs no), 1999: