

Clinical trials of amiloride

TrialResults-center www.trialresultscenter.org

1 hypertension

Trial	Treatments	Patients	Trials design and methods
hydrochlorothiazide + amiloride vs placebo			
MRC old , 1992 n=1081/2213 follow-up:	-	hypertensive patients aged 64-75	

More details and results :

- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in uncomplicated hypertension at <http://www.trialresultscenter.org/go-Q685>

References

MRC old, 1992:

Medical Research Council trial of treatment of hypertension in older adults: principal results. MRC Working Party. BMJ 1992;304:405-12 [[1445513](#)]

2 heart failure

Trial	Treatments	Patients	Trials design and methods
amiloride vs placebo			
Cheitlin , 1991 n=12/12 follow-up: 12 weeks	amiloride versus placebo	men with a history of congestive heart failure	Cross over double blind
frusemide plus amiloride vs captopril			
Richardson , 1987 n=14/14 follow-up: 8 weeks	frusemide plus amiloride versus captopril	patients who had previously been treated with diuretics	Cross over double blind

More details and results :

- diuretics for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q75>

References

Cheitlin, 1991:

Cheitlin MD, Byrd R, Benowitz N, Liu E, Modin G Amiloride improves hemodynamics in patients with chronic congestive heart failure treated with chronic digoxin and diuretics. *Cardiovasc Drugs Ther* 1991;5:719-25 [[1888694](#)]

Richardson, 1987:

Richardson A, Bayliss J, Scriven AJ, Parameshwar J, Poole-Wilson PA, Sutton GC Double-blind comparison of captopril alone against frusemide plus amiloride in mild heart failure. *Lancet* 1987;2:709-11 [[2888942](#)]