

# Clinical trials of amlodipine plus benazepril

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 hypertension

Trial	Treatments	Patients	Trials design and methods
<b>amlodipine plus benazepril vs hydrochlorothiazide plus benazepril</b>			
<b>ACCOMPLISH , 2008</b> [NCT00170950] n=5744/5762 follow-up: 36 months	benazepril 40mg plus amlodipine 5mg daily versus benazepril 40mg plus hydrochlorothiazide 12.5mg daily	patients with hypertension who were at high risk for cardiovascular events	Parallel groups double blind US, Sweden, Norway, Denmark, Finland

More details and results :

- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in patients with additional risk factor at <http://www.trialresultscenter.org/go-Q686>

## References

### ACCOMPLISH, 2008:

Jamerson K, Weber MA, Bakris GL, Dahlf B, Pitt B, Shi V, Hester A, Gupte J, Gatlin M, Velazquez EJ Benazepril plus amlodipine or hydrochlorothiazide for hypertension in high-risk patients. N Engl J Med 2008 Dec 4;359:2417-28 [19052124]

Jamerson KA, Bakris GL, Wun CC, Dahlf B, Lefkowitz M, Manfreda S, Pitt B, Velazquez EJ, Weber MA Rationale and design of the avoiding cardiovascular events through combination therapy in patients living with systolic hypertension (ACCOMPLISH) trial: the first randomized controlled trial to compare the clinical outcome effects of first-line combination therapies in hypertension. Am J Hypertens 2004;17:793-801 [15363822] [10.1016/j.amjhyper.2004.05.004](https://doi.org/10.1016/j.amjhyper.2004.05.004)

Entry terms: amlodipine