

Clinical trials of ardeparin

TrialResults-center www.trialresultscenter.org

1 thrombosis prevention

| Trial | Treatments | Patients | Trials design and methods |
|----------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|---------------------------|
| ardeparin vs placebo | | | |
| Levine , 1996 n=122/124 follow-up: 14 days | ardeparin 50/kgx2 +elastic stockings versus Placebo+elastic stockings | Knee | double blind |
| out of hospital Ardeparin vs standard prophylaxis | | | |
| Heit , 2000 n=589/572 | 65279;in hospital thromboprophylaxis followed by out of hospital Ardeparin 100 IU/kg once a day for a total duration of 42 days versus Ardeparin 50 IU/kg twice a day for 4-10 days | THR or TKR | |
| ardeparin vs unfractionated heparin | | | |
| Godwin , 1993 n=595/309 | Ardeparin 90 and 50 units/kg b.i.d versus UFH 10 000 units | Abdominopelvic surgery | Blind |

More details and results :

- antithrombotics for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q37>
- antithrombotics for thrombosis prevention in elective major knee surgery at <http://www.trialresultscenter.org/go-Q38>
- antithrombotics for thrombosis prevention in general surgery at <http://www.trialresultscenter.org/go-Q92>
- heparin (UFH or LMWH) for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q189>
- LMWH for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q190>
- heparin (UFH or LMWH) for thrombosis prevention in general surgery at <http://www.trialresultscenter.org/go-Q195>
- LMWH for thrombosis prevention in general surgery at <http://www.trialresultscenter.org/go-Q196>

References

Levine, 1996:

Levine MN, Gent M, Hirsh J, Weitz J, Turpie AG, Powers P, Neemeh J, Willan A, Skingley P Ardeparin (low-molecular-weight heparin) vs graduated compression stockings for the prevention of venous thromboembolism. A randomized trial in patients undergoing knee surgery. Arch Intern Med 1996 Apr 22;156:851-6 [[8774203](#)]

Heit, 2000:

Heit JA, Elliott CG, Trowbridge AA, Morrey BF, Gent M, Hirsh J Ardeparin sodium for extended out-of-hospital prophylaxis against venous thromboembolism after total hip or knee replacement. A randomized, double-blind, placebo-controlled trial. Ann Intern Med 2000;132:853-61 [[10836911](#)]

Godwin, 1993: