

Clinical trials of balloon angioplasty

TrialResults-center www.trialresultscenter.org

1 stable angina

Trial	Treatments	Patients	Trials design and methods
balloon angioplasty vs medical treatment			
RITA 2 , 1997 n=504/514 follow-up: 7y	PTCA within 3 mo of the randomisation versus medical treatment	Angina leading to admission within 90days, previous Q wave MI, no previousPTCA, no left main stem disease	Parallel groups open UK
ACME , 1992 n=105/107 follow-up: 5y	PTCA within 3 days of randomization versus medical treatment (nitrates, beta-blockers, calcium blockers)	Stable angina, history of angina, MIwithin 3 months, exercise test with STdepression >3 mm, no previous PTCA; Single or serial stenosis within sameartery 70% to 99% proximal twothirds	Parallel groups open US
ACME 2 (Folland) , 1997 n=51/50 follow-up: 5y	PTCA versus medical therapy	Stable angina, history of angina, MIwithin 3 months, exercise test with STdepression >3 mm, no previous PTCA; Stenosis >70% proximal two thirds,no main artery stenosis >50% , no 3vessel disease	Parallel groups open
ACIP , 1997 n=192/366 follow-up: 24 months	revascularization by angioplasty or bypass surgery versus angina-guided drug therapy or angina plus ischemia-guided drug therapy	clinically stable patients with angiographically documented coronary disease (50% stenosis in 1 major vessel or branch) suitable for revascularization	Parallel groups open
INSPIRE , 2006 n=104/101 follow-up: 60 months	coronary revascularization for suppressing scintigraphic ischemia versus intensive medical therapy strategy	Stable survivors of MI, total perfusion defect size 20% , ischemic defect size 10% (by adenosine SPECT), EF 35% t	Parallel groups open
SWISSI II , 2007 [NCT00387231] n=96/105 follow-up: 10.2y	Percutaneous coronary intervention aimed at full revascularization versus intensive anti-ischemic drug therapy	patients with a recent MI, silent myocardial ischemia verified by stress imaging, and 1- or 2-vessel coronary artery disease	Parallel groups open Switzerland
MASS , 1995 n=72/72 follow-up: 5y	PTCA versus medical treatment (aspirin, nitrates, beta-blockers and calcium channel blocking	Stable angina, no Q wave MI, no leftventricular dysfunction	Parallel groups open Brazil

continued...

Trial	Treatments	Patients	Trials design and methods
Sievers , 1993 n=44/44 follow-up: 2y	PTCA versus medical treatment	Previous nonQ wave MI, no angina indaily life, no previous Q wave MI	Parallel groups open Germany
balloon angioplasty vs CABG			
EAST , 1994 [NCT00000465] n=198/194 follow-up: 3 y	transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with multivessels coronary artery disease	open USA
GABI , 1994 n=182/177 follow-up: 1 y	Percutaneous transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with symptomatic multivessel coronary disease	open Germany
BARI , 1996 [NCT00000462] n=915/914 follow-up: 5.4 y	PTCA versus CABG	Patients with multivessel disease	open USA, Canada
RITA , 1993 n=510/501 follow-up: 2.5 y (6.5y)	percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery	patients with one, two, or three diseased coronary arteries	open UK
ERACI , 1992 n=63/64 follow-up: 3.8 y	Percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with multivessel disease and lesions suitable for either form of therapy	open Argentina
MASS , 1995 n=72/70 follow-up: 3.2 y	percutaneous transluminal coronaryangioplasty versus mammary bypass surgery	patients with stable angina,normal ventricular function and a proximal stenosis of the leftanterior descending coronary artery >80%	open Brazil
Toulouse , 1992 n=76/76 follow-up: 2.8 y	PTCA versus CABG	patients with multivessels coronary artery disease	open France
Lausanne , 1994 n=68/66 follow-up: 3.2 y	transluminal coronary angioplasty versus Coronary artery bypass grafting	patients with isolated proximal left anterior descending artery stenosis, conserved left ventricular function, and documented ischaemia	open Switzerland
CABRI , 1995 n=541/513 follow-up: 1 y	percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with symptomatic multivessel coronary disease	open Europe

More details and results :

- myocardial revascularization for stable angina in all type of patient at <http://www.trialresultscenter.org/go-Q25>
- myocardial revascularization for stable angina in single vessel disease at <http://www.trialresultscenter.org/go-Q27>
- myocardial revascularization for stable angina in multivessels disease at <http://www.trialresultscenter.org/go-Q28>

References

RITA 2, 1997:

Coronary angioplasty versus medical therapy for angina: the second Randomised Intervention Treatment of Angina (RITA-2) trial. RITA-2 trial participants Lancet 1997;350:461-8 [[9274581](#)]

Henderson RA, Pocock SJ, Clayton TC, Knight R, Fox KA, Julian DG, Chamberlain DA Seven-year outcome in the RITA-2 trial: coronary angioplasty versus medical therapy J Am Coll Cardiol 2003;42:1161-70 [[14522473](#)]

ACME, 1992:

Parisi AF, Folland ED, Hartigan P A comparison of angioplasty with medical therapy in the treatment of single-vessel coronary artery disease. Veterans Affairs ACME Investigators N Engl J Med 1992;326:10-6 [[1345754](#)] [10.1056/NEJM199201023260102](#)

Hartigan PM, Giacomini JC, Folland ED, Parisi AF Two- to three-year follow-up of patients with single-vessel coronary artery disease randomized to PTCA or medical therapy (results of a VA cooperative study). Veterans Affairs Cooperative Studies Program ACME Investigators. Angioplasty Compared to Medicine Am J Cardiol 1998;82:1445-50 [[9874045](#)]

ACME 2 (Folland), 1997:

Folland ED, Hartigan PM, Parisi AF Percutaneous transluminal coronary angioplasty versus medical therapy for stable angina pectoris: outcomes for patients with double-vessel versus single-vessel coronary artery disease in a Veterans Affairs Cooperative randomized trial. Veterans Affairs ACME InvestigatorS J Am Coll Cardiol 1997;29:1505-11 [[9180111](#)]

Henderson RA, Pocock SJ, Clayton TC, Knight R, Fox KA, Julian DG, Chamberlain DA Seven-year outcome in the RITA-2 trial: coronary angioplasty versus medical therapy J Am Coll Cardiol 2003;42:1161-70 [[14522473](#)]

ACIP, 1997:

Davies RF, Goldberg AD, Forman S, Pepine CJ, Knatterud GL, Geller N, Sopko G, Pratt C, Deanfield J, Conti CR Asymptomatic Cardiac Ischemia Pilot (ACIP) study two-year follow-up: outcomes of patients randomized to initial strategies of medical therapy versus revascularization. Circulation 1997;95:2037-43 [[9133513](#)]

INSPIRE, 2006:

Mahmarian JJ, Dakik HA, Filipchuk NG, Shaw LJ, Iskander SS, Ruddy TD, Keng F, Henzlova MJ, Allam A, Moy LA, Pratt CM An initial strategy of intensive medical therapy is comparable to that of coronary revascularization for suppression of scintigraphic ischemia in high-risk but stable survivors of acute myocardial infarction. J Am Coll Cardiol 2006;48:2458-67 [[17174182](#)]

SWISSI II, 2007:

Erne P, Schoenenberger AW, Burckhardt D, Zuber M, Kiowski W, Buser PT, Dubach P, Resink TJ, Pfisterer M Effects of percutaneous coronary interventions in silent ischemia after myocardial infarction: the SWISSI II randomized controlled trial. JAMA 2007;297:1985-91 [[17488963](#)]

MASS, 1995:

Hueb WA, Bellotti G, de Oliveira SA, Arie S, de Albuquerque CP, Jatene AD, Pileggi F The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses *J Am Coll Cardiol* 1995;26:1600-5 [7594092] [10.1016/0735-1097\(95\)00384-3](https://doi.org/10.1016/0735-1097(95)00384-3)

Sievers, 1993:

Sievers N, Hamm CW, Herzner A, Kuck KH Medical therapy versus PTCA: a prospective, randomized trial in patients with asymptomatic coronary single-vessel disease. Abstract. *Circulation*. 1993;88(suppl I):I-297 [0]

EAST, 1994:

King SB 3rd, Lembo NJ, Weintraub WS, Kosinski AS, Barnhart HX, Kutner MH, Alazraki NP, Guyton RA, Zhao XQ A randomized trial comparing coronary angioplasty with coronary bypass surgery. Emory Angioplasty versus Surgery Trial (EAST) *N Engl J Med* 1994 Oct 20;331:1044-50 [8090163]

GABI, 1994:

Hamm CW, Reimers J, Ischinger T, Rupprecht HJ, Berger J, Bleifeld W A randomized study of coronary angioplasty compared with bypass surgery in patients with symptomatic multivessel coronary disease. German Angioplasty Bypass Surgery Investigation (GABI) *N Engl J Med* 1994 Oct 20;331:1037-43 [8090162]

BARI, 1996:

Comparison of coronary bypass surgery with angioplasty in patients with multivessel disease. The Bypass Angioplasty Revascularization Investigation (BARI) Investigators. *N Engl J Med* 1996 Jul 25;335:217-25 [8657237]

RITA, 1993:

Coronary angioplasty versus coronary artery bypass surgery: the Randomized Intervention Treatment of Angina (RITA) trial. *Lancet* 1993 Mar 6;341:573-80 [8094826]

ERACI, 1992:

Rodriguez A, Bouillon F, Perez-Balino N, Paviotti C, Liprandi MI, Palacios IF Argentine randomized trial of percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery in multivessel disease (ERACI): in-hospital results and 1-year follow-up. ERACI Group. *J Am Coll Cardiol* 1993 Oct;22:1060-7 [8409041]

MASS, 1995:

Hueb WA, Bellotti G, de Oliveira SA, Arie S, de Albuquerque CP, Jatene AD, Pileggi F The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses. *J Am Coll Cardiol* 1995 Dec;26:1600-5 [7594092]

Toulouse, 1992:

Lausanne, 1994:

Goy JJ, Eeckhout E, Burnand B, Vogt P, Stauffer JC, Hurni M, Stumpe F, Ruchat P, Sadeghi H, Kappenberger L Coronary angioplasty versus left internal mammary artery grafting for isolated proximal left anterior descending artery stenosis. *Lancet* 1994 Jun 11;343:1449-53 [7911175]

CABRI, 1995:

First-year results of CABRI (Coronary Angioplasty versus Bypass Revascularisation Investigation). CABRI Trial Participants. *Lancet* 1995 Nov 4;346:1179-84 [7475656]

Martuscelli E, Clementi F, Gallagher MM, D'Eliseo A, Chiricolo G, Nigri A, Marino B, Romeo F Revascularization strategy in patients with multivessel disease and a major vessel chronically occluded; data from the CABRI trial. *Eur J Cardiothorac Surg* 2008;33:4-8 [17988889]

2 coronary artery disease

Trial	Treatments	Patients	Trials design and methods
balloon angioplasty vs medical treatment			
RITA 2 , 1997 n=504/514 follow-up: 7y	PTCA within 3 mo of the randomisation versus medical treatment	Angina leading to admission within 90days, previous Q wave MI, no previousPTCA, no left main stem disease	Parallel groups open UK
ACME , 1992 n=105/107 follow-up: 5y	PTCA within 3 days of randomization versus medical treatment (nitrates, beta-blockers, calcium blockers)	Stable angina, history of angina, MIwithin 3 months, exercise test with STdepression >3 mm, no previous PTCA; Single or serial stenosis within sameartery 70% to 99% proximal twothirds	Parallel groups open US
ACME 2 (Folland) , 1997 n=51/50 follow-up: 5y	PTCA versus medical therapy	Stable angina, history of angina, MIwithin 3 months, exercise test with STdepression >3 mm, no previous PTCA; Stenosis >70% proximal two thirds,no main artery stenosis >50% , no 3vessel disease	Parallel groups open
ACIP , 1997 n=192/366 follow-up: 24 months	revascularization by angioplasty or bypass surgery versus angina-guided drug therapy or angina plus ischemia-guided drug therapy	clinically stable patients with angiographically documented coronary disease (50% stenosis in 1 major vessel or branch) suitable for revascularization	Parallel groups open
INSPIRE , 2006 n=104/101 follow-up: 60 months	coronary revascularization for suppressing scintigraphic ischemia versus intensive medical therapy strategy	Stable survivors of MI, total perfusion defect size 20% , ischemic defect size 10% (by adenosine SPECT), EF 35% t	Parallel groups open
SWISSI II , 2007 [NCT00387231] n=96/105 follow-up: 10.2y	Percutaneous coronary intervention aimed at full revascularization versus intensive anti-ischemic drug therapy	patients with a recent MI, silent myocardial ischemia verified by stress imaging, and 1- or 2-vessel coronary artery disease	Parallel groups open Switzerland
MASS , 1995 n=72/72 follow-up: 5y	PTCA versus medical treatment (aspirin, nitrates, beta-blockers and calcium channel blocking	Stable angina, no Q wave MI, no leftventricular dysfunction	Parallel groups open Brazil
Sievers , 1993 n=44/44 follow-up: 2y	PTCA versus medical treatment	Previous nonQ wave MI, no angina indaily life, no previous Q wave MI	Parallel groups open Germany
balloon angioplasty vs CABG			

continued...

Trial	Treatments	Patients	Trials design and methods
EAST , 1994 [NCT00000465] n=198/194 follow-up: 3 y	transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with multivessels coronary artery disease	open USA
GABI , 1994 n=182/177 follow-up: 1 y	Percutaneous transluminal coronary angioplasty versus coronary-artery bypass grafting	patients with symptomatic multivessel coronary disease	open Germany
BARI , 1996 [NCT00000462] n=915/914 follow-up: 5.4 y	PTCA versus CABG	Patients with multivessel disease	open USA, Canada
RITA , 1993 n=510/501 follow-up: 2.5 y (6.5y)	percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery	patients with one, two, or three diseased coronary arteries	open UK
ERACI , 1992 n=63/64 follow-up: 3.8 y	Percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with multivessel disease and lesions suitable for either form of therapy	open Argentina
MASS , 1995 n=72/70 follow-up: 3.2 y	percutaneous transluminal coronaryangioplasty versus mammary bypass surgery	patients with stable angina,normal ventricular function and a proximal stenosis of the leftanterior descending coronary artery >80%	open Brazil
Toulouse , 1992 n=76/76 follow-up: 2.8 y	PTCA versus CABG	patients with multivessels coronary artery disease	open France
Lausanne , 1994 n=68/66 follow-up: 3.2 y	transluminal coronary angioplasty versus Coronary artery bypass grafting	patients with isolated proximal left anterior descending artery stenosis, conserved left ventricular function, and documented ischaemia	open Switzerland
CABRI , 1995 n=541/513 follow-up: 1 y	percutaneous transluminal coronary angioplasty versus coronary artery bypass grafting	patients with symptomatic multivessel coronary disease	open Europe

More details and results :

- myocardial revascularization for coronary artery disease in all type of patient at <http://www.trialresultscenter.org/go-Q26>
- myocardial revascularization for coronary artery disease in multivessels disease at <http://www.trialresultscenter.org/go-Q31>

- myocardial revascularization for coronary artery disease in single vessel disease at <http://www.trialresultscenter.org/go-Q32>

References

RITA 2, 1997:

Coronary angioplasty versus medical therapy for angina: the second Randomised Intervention Treatment of Angina (RITA-2) trial. RITA-2 trial participants Lancet 1997;350:461-8 [[9274581](#)]

Henderson RA, Pocock SJ, Clayton TC, Knight R, Fox KA, Julian DG, Chamberlain DA Seven-year outcome in the RITA-2 trial: coronary angioplasty versus medical therapy J Am Coll Cardiol 2003;42:1161-70 [[14522473](#)]

ACME, 1992:

Parisi AF, Folland ED, Hartigan P A comparison of angioplasty with medical therapy in the treatment of single-vessel coronary artery disease. Veterans Affairs ACME Investigators N Engl J Med 1992;326:10-6 [[1345754](#)] [10.1056/NEJM199201023260102](#)

Hartigan PM, Giacomini JC, Folland ED, Parisi AF Two- to three-year follow-up of patients with single-vessel coronary artery disease randomized to PTCA or medical therapy (results of a VA cooperative study). Veterans Affairs Cooperative Studies Program ACME Investigators. Angioplasty Compared to Medicine Am J Cardiol 1998;82:1445-50 [[9874045](#)]

ACME 2 (Folland), 1997:

Folland ED, Hartigan PM, Parisi AF Percutaneous transluminal coronary angioplasty versus medical therapy for stable angina pectoris: outcomes for patients with double-vessel versus single-vessel coronary artery disease in a Veterans Affairs Cooperative randomized trial. Veterans Affairs ACME InvestigatorS J Am Coll Cardiol 1997;29:1505-11 [[9180111](#)]

Henderson RA, Pocock SJ, Clayton TC, Knight R, Fox KA, Julian DG, Chamberlain DA Seven-year outcome in the RITA-2 trial: coronary angioplasty versus medical therapy J Am Coll Cardiol 2003;42:1161-70 [[14522473](#)]

ACIP, 1997:

Davies RF, Goldberg AD, Forman S, Pepine CJ, Knatterud GL, Geller N, Sopko G, Pratt C, Deanfield J, Conti CR Asymptomatic Cardiac Ischemia Pilot (ACIP) study two-year follow-up: outcomes of patients randomized to initial strategies of medical therapy versus revascularization. Circulation 1997;95:2037-43 [[9133513](#)]

INSPIRE, 2006:

Mahmorian JJ, Dakik HA, Filipchuk NG, Shaw LJ, Iskander SS, Ruddy TD, Keng F, Henzlova MJ, Allam A, Moy LA, Pratt CM An initial strategy of intensive medical therapy is comparable to that of coronary revascularization for suppression of scintigraphic ischemia in high-risk but stable survivors of acute myocardial infarction. J Am Coll Cardiol 2006;48:2458-67 [[17174182](#)]

SWISSI II, 2007:

Erne P, Schoenenberger AW, Burckhardt D, Zuber M, Kiowski W, Buser PT, Dubach P, Resink TJ, Pfisterer M Effects of percutaneous coronary interventions in silent ischemia after myocardial infarction: the SWISSI II randomized controlled trial. JAMA 2007;297:1985-91 [[17488963](#)]

MASS, 1995:

Hueb WA, Bellotti G, de Oliveira SA, Arie S, de Albuquerque CP, Jatene AD, Pileggi F The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses J Am Coll Cardiol 1995;26:1600-5 [[7594092](#)] [10.1016/0735-1097\(95\)00384-3](#)

Sievers, 1993:

Sievers N, Hamm CW, Herzner A, Kuck KH Medical therapy versus PTCA: a prospective, randomized trial in patients with asymptomatic coronary single-vessel disease. Abstract. Circulation. 1993;88(suppl I):I-297 [0]

EAST, 1994:

King SB 3rd, Lembo NJ, Weintraub WS, Kosinski AS, Barnhart HX, Kutner MH, Alazraki NP, Guyton RA, Zhao XQ A randomized trial comparing coronary angioplasty with coronary bypass surgery. Emory Angioplasty versus Surgery Trial (EAST) N Engl J Med 1994 Oct 20;331:1044-50 [8090163]

GABI, 1994:

Hamm CW, Reimers J, Ischinger T, Rupprecht HJ, Berger J, Bleifeld W A randomized study of coronary angioplasty compared with bypass surgery in patients with symptomatic multivessel coronary disease. German Angioplasty Bypass Surgery Investigation (GABI) N Engl J Med 1994 Oct 20;331:1037-43 [8090162]

BARI, 1996:

Comparison of coronary bypass surgery with angioplasty in patients with multivessel disease. The Bypass Angioplasty Revascularization Investigation (BARI) Investigators. N Engl J Med 1996 Jul 25;335:217-25 [8657237]

RITA, 1993:

Coronary angioplasty versus coronary artery bypass surgery: the Randomized Intervention Treatment of Angina (RITA) trial. Lancet 1993 Mar 6;341:573-80 [8094826]

ERACI, 1992:

Rodriguez A, Bouillon F, Perez-Balino N, Paviotti C, Liprandi MI, Palacios IF Argentine randomized trial of percutaneous transluminal coronary angioplasty versus coronary artery bypass surgery in multivessel disease (ERACI): in-hospital results and 1-year follow-up. ERACI Group. J Am Coll Cardiol 1993 Oct;22:1060-7 [8409041]

MASS, 1995:

Hueb WA, Bellotti G, de Oliveira SA, Arie S, de Albuquerque CP, Jatene AD, Pileggi F The Medicine, Angioplasty or Surgery Study (MASS): a prospective, randomized trial of medical therapy, balloon angioplasty or bypass surgery for single proximal left anterior descending artery stenoses. J Am Coll Cardiol 1995 Dec;26:1600-5 [7594092]

Toulouse, 1992:

Lausanne, 1994:

Goy JJ, Eeckhout E, Burnand B, Vogt P, Stauffer JC, Hurni M, Stumpe F, Ruchat P, Sadeghi H, Kappenberger L Coronary angioplasty versus left internal mammary artery grafting for isolated proximal left anterior descending artery stenosis. Lancet 1994 Jun 11;343:1449-53 [7911175]

CABRI, 1995:

First-year results of CABRI (Coronary Angioplasty versus Bypass Revascularisation Investigation). CABRI Trial Participants. Lancet 1995 Nov 4;346:1179-84 [7475656]

Martuscelli E, Clementi F, Gallagher MM, D'Eliseo A, Chiricolo G, Nigri A, Marino B, Romeo F Revascularization strategy in patients with multivessel disease and a major vessel chronically occluded; data from the CABRI trial. Eur J Cardiothorac Surg 2008;33:4-8 [17988889]

Entry terms: PTCA