

Clinical trials of clopidogrel+aspirin

TrialResults-center www.trialresultscenter.org

1 stent

Trial	Treatments	Patients	Trials design and methods
clopidogrel+aspirin vs aspirin			
REAL-LATE, ZEST-LATE , 2010 [NCT00484926] n=1357/1344 follow-up: 19.2 months	clopidogrel plus aspirin versus aspirin alone	patients who had received drugeluting stents and had been free of major adverse cardiac or cerebrovascular events and major bleeding for a period of at least 12 months	Parallel groups open South Korea

More details and results :

- antithrombotics for stent in all type of patients at <http://www.trialresultscenter.org/go-Q151>
- dual antiplatelet therapy for stent in all type of patients at <http://www.trialresultscenter.org/go-Q578>

References

REAL-LATE, ZEST-LATE, 2010:

Park SJ, Park DW, Kim YH, Kang SJ, Lee SW, Lee CW, Han KH, Park SW, Yun SC, Lee SG, Rha SW, Seong IW, Jeong MH, Hur SH, Lee NH, Yoon J, Yang JY, Lee BK, Choi YJ, Chung WS, Lim DS, Cheong SS, Kim KS, Chae JK, Nah DY, Jeon DS, Seung KB, Jang JS, Park HS, Le Duration of dual antiplatelet therapy after implantation of drug-eluting stents. N Engl J Med 2010;362:1374-82 [20231231] 10.1056/NEJMoa1001266

2 CABG surgery

Trial	Treatments	Patients	Trials design and methods
clopidogrel+aspirin vs aspirin			
CASCADE , 2009 [NCT00228423] n=56/57 follow-up: 1 y	aspirin 162 mg plus clopidogrel 75 mg daily for 1 year versus aspirin 162 mg plus placebo daily	patients after CABG involving at least two saphenous vein grafts	Parallel groups double blind

More details and results :

- antiplatelets drug for CABG surgery in all type of patients at <http://www.trialresultscenter.org/go-Q225>

References

CASCADE, 2009:

Kulik A, Le May M, Wells GA, Mesana TG, Ruel M The clopidogrel after surgery for coronary artery disease (CASCADE) randomized controlled trial: clopidogrel and aspirin versus aspirin alone after coronary bypass surgery [NCT00228423]. Curr Control Trials Cardiovasc Med 2005 Oct 11;6:15 [16219100]

Kulik A, Le May MR, Voisine P, Tardif JC, Delarochelliere R, Naidoo S, Wells GA, Mesana TG, Ruel M Aspirin plus clopidogrel versus aspirin alone after coronary artery bypass grafting: the clopidogrel after surgery for coronary artery disease (CASCADE) Trial. Circulation 2010 Dec 21;122:2680-7 [21135365] 10.1161/CIRCULATIONAHA.110.978007

3 percutaneous coronary intervention

Trial	Treatments	Patients	Trials design and methods
clopidogrel+aspirin vs aspirin			
REAL-LATE, ZEST-LATE , 2010 [NCT00484926] n=1357/1344 follow-up: 19.2 months	clopidogrel plus aspirin versus aspirin alone	patients who had received drugeluting stents and had been free of major adverse cardiac or cerebrovascular events and major bleeding for a period of at least 12 months	Parallel groups open South Korea

More details and results :

- antiplatelets drug for percutaneous coronary intervention in all type of patients at <http://www.trialresultscenter.org/go-Q389>

References

REAL-LATE, ZEST-LATE, 2010:

Park SJ, Park DW, Kim YH, Kang SJ, Lee SW, Lee CW, Han KH, Park SW, Yun SC, Lee SG, Rha SW, Seong IW, Jeong MH, Hur SH, Lee NH, Yoon J, Yang JY, Lee BK, Choi YJ, Chung WS, Lim DS, Cheong SS, Kim KS, Chae JK, Nah DY, Jeon DS, Seung KB, Jang JS, Park HS, Le Duration of dual antiplatelet therapy after implantation of drug-eluting stents. N Engl J Med 2010;362:1374-82 [20231231] 10.1056/NEJMoa1001266

Entry terms: clopidogrel, Plavix, Iscover