

Clinical trials of clopidogrel+aspirin

TrialResults-center www.trialresultscenter.org

1 stent

| Trial | Treatments | Patients | Trials design and methods |
|--|---|---|--|
| clopidogrel+aspirin vs aspirin | | | |
| REAL-LATE, ZEST-LATE, 2010 [NCT00484926] n=1357/1344 follow-up: 19.2 months | clopidogrel plus aspirin versus aspirin alone | patients who had received drugeluting stents and had been free of major adverse cardiac or cerebrovascular events and major bleeding for a period of at least 12 months | Parallel groups open South Korea |

More details and results :

- antithrombotics for stent in all type of patients at <http://www.trialresultscenter.org/go-Q151>
- dual antiplatelet therapy for stent in all type of patients at <http://www.trialresultscenter.org/go-Q578>

References

REAL-LATE, ZEST-LATE, 2010:

Park SJ, Park DW, Kim YH, Kang SJ, Lee SW, Lee CW, Han KH, Park SW, Yun SC, Lee SG, Rha SW, Seong IW, Jeong MH, Hur SH, Lee NH, Yoon J, Yang JY, Lee BK, Choi YJ, Chung WS, Lim DS, Cheong SS, Kim KS, Chae JK, Nah DY, Jeon DS, Seung KB, Jang JS, Park HS, Le Duration of dual antiplatelet therapy after implantation of drug-eluting stents. N Engl J Med 2010;362:1374-82 [[20231231](#)] [10.1056/NEJMoa1001266](https://doi.org/10.1056/NEJMoa1001266)

2 CABG surgery

| Trial | Treatments | Patients | Trials design and methods |
|---|---|--|---------------------------------|
| clopidogrel+aspirin vs aspirin | | | |
| CASCADE, 2009 [NCT00228423] n=56/57 follow-up: 1 y | aspirin 162 mg plus clopidogrel 75 mg daily for 1 year versus aspirin 162 mg plus placebo daily | patients after CABG involving at least two saphenous vein grafts | Parallel groups double blind |

More details and results :

- antiplatelets drug for CABG surgery in all type of patients at <http://www.trialresultscenter.org/go-Q225>

References

CASCADE, 2009:

Kulik A, Le May M, Wells GA, Mesana TG, Ruel M The clopidogrel after surgery for coronary artery disease (CASCADE) randomized controlled trial: clopidogrel and aspirin versus aspirin alone after coronary bypass surgery [NCT00228423]. *Curr Control Trials Cardiovasc Med* 2005 Oct 11;6:15 [[16219100](#)]

Kulik A, Le May MR, Voisine P, Tardif JC, Delarochelliere R, Naidoo S, Wells GA, Mesana TG, Ruel M Aspirin plus clopidogrel versus aspirin alone after coronary artery bypass grafting: the clopidogrel after surgery for coronary artery disease (CASCADE) Trial. *Circulation* 2010 Dec 21;122:2680-7 [[21135365](#)] [10.1161/CIRCULATIONAHA.110.978007](#)

3 percutaneous coronary intervention

| Trial | Treatments | Patients | Trials design and methods |
|---|---|---|--|
| clopidogrel+aspirin vs aspirin | | | |
| REAL-LATE, ZEST-LATE , 2010 [NCT00484926] n=1357/1344 follow-up: 19.2 months | clopidogrel plus aspirin versus aspirin alone | patients who had received drugeluting stents and had been free of major adverse cardiac or cerebrovascular events and major bleeding for a period of at least 12 months | Parallel groups open South Korea |

More details and results :

- antiplatelets drug for percutaneous coronary intervention in all type of patients at <http://www.trialresultscenter.org/go-Q389>

References

REAL-LATE, ZEST-LATE, 2010:

Park SJ, Park DW, Kim YH, Kang SJ, Lee SW, Lee CW, Han KH, Park SW, Yun SC, Lee SG, Rha SW, Seong IW, Jeong MH, Hur SH, Lee NH, Yoon J, Yang JY, Lee BK, Choi YJ, Chung WS, Lim DS, Cheong SS, Kim KS, Chae JK, Nah DY, Jeon DS, Seung KB, Jang JS, Park HS, Le Duration of dual antiplatelet therapy after implantation of drug-eluting stents. *N Engl J Med* 2010;362:1374-82 [[20231231](#)] [10.1056/NEJMoa1001266](#)

Entry terms: clopidogrel, Plavix, Iscover