

Clinical trials of dapagliflozin + Glimepiride

TrialResults-center www.trialresultscenter.org

1 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
dapagliflozin + Glimepiride vs glimepiride			
Strojek , 2011 [NCT00680745] n=NA follow-up:	Dapagliflozin in Comb.With Glimepiride versus glimepiride alone	Type2 Diab.Who Have Inadeq. Glycaemic Control on Glimepiride Therapy Alone	

More details and results :

- SGLT2 inhibitors for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q479>

References

Strojek, 2011:

Strojek K, Yoon KH, Hrubá V, Elze M, Langkilde AM, Parikh S Effect of dapagliflozin in patients with type 2 diabetes who have inadequate glycaemic control with glimepiride: a randomized, 24-week, double-blind, placebo-controlled trial. *Diabetes Obes Metab* 2011;13:928-38 [21672123] [10.1111/j.1463-1326.2011.01434.x](https://doi.org/10.1111/j.1463-1326.2011.01434.x)

Kohan DE, Fioretto P, Johnsson K, Parikh S, Ptaszynska A, Ying L The effect of dapagliflozin on renal function in patients with type 2 diabetes. *J Nephrol* 2016;: [26894924]

Strojek K, Yoon KH, Hrubá V, Elze M, Langkilde AM, Parikh S [Effect of dapagliflozin in patients with type 2 diabetes who have inadequate glycaemic control with glimepiride]. *Dtsch Med Wochenschr* 2013;138 Suppl 1:S16-26 [23529567]

Entry terms: dapagliflozin, dapagliflozin, forxiga, BMS 512148, BMS512148, BMS-512148,