

Clinical trials of deserpidine +methylclothiazide

TrialResults-center www.trialresultscenter.org

1 hypertension

Trial	Treatments	Patients	Trials design and methods
deserpidine +methylclothiazide vs placebo			
HSCS , 1974 n=233/219 follow-up: 2.3y	deserpidine 1mg/d + methylclothiazide 10mg/d versus placebo	stroke	Parallel groups Double blind USA

More details and results :

- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in post stroke at <http://www.trialresultscenter.org/go-Q20>

References

HSCS, 1974:

Effect of antihypertensive treatment on stroke recurrence. Hypertension-Stroke Cooperative Study Group. JAMA 1974 Jul 22;229:409-18 [[4599980](#)]

Effect of antihypertensive treatment on stroke recurrence. Hypertension-Stroke Cooperative Study Group. JAMA 1974;229:409-18 [[4599980](#)]

Entry terms: candesartan, candesartan cilexetil, 1-(cyclohexylcarbonyloxy)ethyl-2-ethoxy-1-(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)-1H-benzimidazole-7-carboxylate, TCV 116, TCV-116, Atacand, Astra brand of candesartan cilexetil, Takeda brand of candesartan cilexetil, Blopress, Kenzen, Promed brand of candesartan cilexetil, Amias, AstraZeneca brand of candesartan cilexetil, Parapres, Almirall brand of candesartan cilexetil, , candesartan, 2-ethoxy-7-carboxy-1-(2'-(1H-tetrazol-5-yl)biphenyl-4-yl)methylbenzimidazole, CV 11974, CV11974, CV-11974,