

Clinical trials of diet

TrialResults-center www.trialresultscenter.org

1 post myocardial infarction

| Trial | Treatments | Patients | Trials design and methods |
|---|--|--------------------|---------------------------|
| low fat diet vs mediterranean-style diet | | | |
| Tuttle , 2008 n=NA follow-up: 24 months | low-fat versus Mediterranean-style diets | First MI survivors | Parallel groups open |

More details and results :

- cholesterol lowering intervention for post myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q45>

References

Tuttle, 2008:

Tuttle KR, Shuler LA, Packard DP, Milton JE, Daratha KB, Bibus DM, Short RA Comparison of low-fat versus Mediterranean-style dietary intervention after first myocardial infarction (from The Heart Institute of Spokane Diet Intervention and Evaluation Trial). Am J Cardiol 2008;101:1523-30 [[18489927](#)]

2 cardiovascular prevention

| Trial | Treatments | Patients | Trials design and methods |
|--|--|---------------------------------------|---------------------------|
| diet vs control | | | |
| NORDIET n=45/43 follow-up: | healthy Nordic diet versus control diet (subjects usualWestern diet) | mildly hypercholesterolaemic subjects | Sweden |
| BARON n=NA | - | - | |
| HPT n=NA | - | - | |
| Kumanyika n=NA | - | - | |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|---|---|---|--|
| TAIM n=NA | - | - | |
| DISH n=NA | - | - | |
| Burr (DART 2) , 2003 n=1571/1543 follow-up: 36-108 months | dietary advice (to eat more oily fish) versus No such dietary advice or capsules | men being treated for angina | Parallel groups open UK |
| Burr (DART) , 1989 n=1015/1018 follow-up: 24 months | dietary advice (to eat more oily fish) versus No such dietary advice or capsulesish)ag | post-MI | Parallel groups open UK |
| Mate-Jimenez , 1991 n=19/19 follow-up: 24months | diet advice versus no advice | people with inactive Crohns disease | Parallel groups open with blind assessment Spain |
| Mediterranean diet vs control | | | |
| Lyon n=302/303 follow-up: | - | - | |
| Mediterranean diet with EOVV vs control | | | |
| PREDIMED (olive oil) , 2013 [ISRCTN35739639] n=2543/2450 follow-up: 4.8 years | Mediterranean diet supplemented with extra-virgin olive oil versus control diet (advice to reduce dietary fat) | participants who were at high cardiovascular risk, but with no cardiovascular disease | Parallel groups open Spain |
| Mediterranean diet with nuts vs control | | | |
| PREDIMED (nuts) , 2013 [ISRCTN35739639] n=2454/2450 follow-up: 4.8 years | Mediterranean diet supplemented with mixed nuts versus control diet (advice to reduce dietary fat) | participants who were at high cardiovascular risk, but with no cardiovascular disease | open Spain |
| diet vs usual diet | | | |
| Black , 1994 n=56/55 follow-up: 2.0 years | diet with 20 percent of total caloric intake as fat versus usual diet | patients with nonmelanoma skin cancer | Parallel groups open |
| DART (Burr) , 1989 n=NA follow-up: 2 years | diet advice versus usual diet | men who had recovered from MI | Factorial plan open, blind assessment |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|--|---|--|---|
| Finnish Mental Hospital (Miettinen) , 1985 n=612/610 follow-up: 6.0 years | cholesterol-lowering diet (low in saturated fats and cholesterol and relatively high in polyunsaturated fats) versus usual diet | middle-aged institutionalized women without CHD | Cluster-randomized cross-ove open, blind assessment Finland |
| Finnish Mental Hospital (Turpeinen) , 1979 n=NA follow-up: 6.0 years | cholesterol-lowering diet (low in saturated fats and cholesterol and relatively high in polyunsaturated fats) versus usual diet | middle-aged institutionalized men without CHD | Cluster-randomized cross-ove open, blind assessment Finland |
| Goteborg , 1986 n=10004/20028 follow-up: 10 years | multifactorial intervention programme versus no intervention | men, 47-55 years old at entry | Parallel groups open Sweden |
| Gteborg (Wilhelmsen) , 1986 n=10004/20028 follow-up: 10.0 years | multifactorial intervention programme versus usual care | men, 47-55 years old at entry | Parallel groups open |
| Hjermann , 1981 n=604/628 follow-up: 6.5 years | diet versus usual diet | healthy, normotensive men at high risk of coronary heart disease | Parallel groups open Sweden |
| Kallio , 1979 n=188/187 follow-up: 3.0 years | diet (multifactorial intervention programme) versus usual diet | patients below 65 years who had an acute myocardial infarction | Parallel groups open |
| Los Angeles VA (Dayton) , 1969 n=424/422 follow-up: 65279;8.0 y | diet versus usual diet | men in domiciliary care, age>55, with or without CHD | Parallel groups double blind USA |
| Minnesota coronary survey (Frantz) , 1975 n=2197/2196 follow-up: 1.1 y (max 4.5y) | cholesterol lowering diet versus control diet | 65279;Adult residents ofmental hospitals; no illness restrictions, no cholesterol concentration requirements | Parallel groups double-blind USA |
| MRC low fat , 1965 n=123/129 follow-up: 3 y | - | - | Parallel groups open |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|--|---|---|---|
| MRC Soya , 1968 n=199/194 follow-up: 3.5 y | Rgime pauvre en graisses satures + 85 g/j d'huile de soja versus usual diet | ambulatory men with recent MI | Parallel groups open, blind assessment |
| MRFIT , 1982 n=6428/6438 follow-up: 6.5 y | multifactor intervention program versus usual diet | high-risk men aged 35 to 57 years | Parallel groups open |
| Ornish , 1990 n=28/20 follow-up: 1.0 y | low-fat vegetarian diet, stopping smoking, stress management training, and moderate exercise versus usual-care | Patients with angiographically documented coronary artery disease | Parallel groups open USA |
| Oslo Diet Heart Study (Leren) , 1966 n=206/206 follow-up: 5 y (11y) | diet versus usual care | middle-aged ambulatory men with prior MI | Parallel groups open, blind assessment |
| Rose , 1965 n=28/26 follow-up: 1.2 years | Rgime restreint en graisses + 80 g/j huile de mas versus usual diet | men, <70 years | Parallel groups open |
| Singh , 1992 n=204/202 follow-up: 65279;2.0 years | strict diet versus usual diet | patients with suspected acute myocardial infarction | Parallel groups open |
| STARS (St Thomas, diet) , 1992 n=30/30 follow-up: 3.0 years | dietary advice versus usual diet | patients with angina or past myocardial infarction | open, blind assessment |
| Veterans Ad. (Dayton) , 1969 n=424/422 follow-up: 3.6 and 8 y | cholesterol lowering diet versus usual diet | men in domiciliary care, age>55, with or without CHD | Parallel groups double blind USA |
| WHI low fat , 2005 [NCT00000611] n=19541/29294 follow-up: 8.1y mean | dietary modification intervention to promote dietary change with the goals of reducing intake of total fat to 20% of energy and increasing consumption of vegetables and fruit to at least 5 servings daily and grains to at least 6 servings daily versus usual diet | postmenopausal women, aged 50 to 79 years, without prior breast cancer | Parallel groups open US |

continued...

| Trial | Treatments | Patients | Trials design and methods |
|---|---|--------------------|---|
| WHO Collaborative , 1986 n=30489/26971 follow-up: 5.5 years | multifactorial prevention versus usual diet | middle-aged men | Parallel groups open Belgium, Italy, Poland, UK |
| Woodhill , 1966 n=221/237 follow-up: <7 years | diet versus usual diet | men, 30-59 years | Parallel groups open |
| low fat diet vs mediterranean-style diet | | | |
| Tuttle , 2008 n=NA follow-up: 24 months | low-fat versus Mediterranean-style diets | First MI survivors | Parallel groups open |

More details and results :

- cholesterol lowering intervention for cardiovascular prevention in patients with LDL elevation and without CHD at <http://www.trialresultscenter.org/go-Q5>
- cholesterol lowering intervention for cardiovascular prevention in patients with prior MI or with CHD at <http://www.trialresultscenter.org/go-Q12>
- omega-3 fatty acids for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q121>
- omega-3 fatty acids for cardiovascular prevention in patients at high risk at <http://www.trialresultscenter.org/go-Q123>
- omega-3 fatty acids for cardiovascular prevention in pateints at low risk at <http://www.trialresultscenter.org/go-Q124>
- cholesterol lowering intervention for cardiovascular prevention in all chronical situations at <http://www.trialresultscenter.org/go-Q154>
- lifestyle intervention for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q282>
- diet for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q568>
- statins for cardiovascular prevention in secondary prevention at <http://www.trialresultscenter.org/go-Q689>

References

NORDIET, :

Adamsson V, Reumark A, Fredriksson IB, Hammarstrm E, Vessby B, Johansson G, Risrus U Effects of a healthy Nordic diet on cardiovascular risk factors in hypercholesterolaemic subjects: a randomized controlled trial (NORDIET). J Intern Med 2011;269:150-9 [20964740] [10.1111/j.1365-2796.2010.02290.x](https://doi.org/10.1111/j.1365-2796.2010.02290.x)

BARON, :

Baron JA, Gleason R, Crowe B, Mann JI Preliminary trial of the effect of general practice based nutritional advice. *Br J Gen Pract* 1990;40:137-41 [2115348]

HPT, :

The Hypertension Prevention Trial: three-year effects of dietary changes on blood pressure. Hypertension Prevention Trial Research Group. *Arch Intern Med* 1990;150:153-62 [2404477]

Kumanyika, :

Kumanyika SK, Hebert PR, Cutler JA, Lasser VI, Sugars CP, Steffen-Batey L, Brewer AA, Cameron M, Shepek LD, Cook NR Feasibility and efficacy of sodium reduction in the Trials of Hypertension Prevention, phase I. Trials of Hypertension Prevention Collaborative Research Group. *Hypertension* 1993;22:502-12 [8406655]

TAIM, :

Davis BR, Oberman A, Blafox MD, Wassertheil-Smoller S, Zimbaldi N, Kirchner K, Wylie-Rosett J, Langford HG Lack of effectiveness of a low-sodium/high-potassium diet in reducing antihypertensive medication requirements in overweight persons with mild hypertension. TAIM Research Group. Trial of Antihypertensive Interventions and Management. *Am J Hypertens* 1994;7:926-32 [7826557]

DISH, :

Langford HG, Blafox MD, Oberman A, Hawkins CM, Curb JD, Cutter GR, Wassertheil-Smoller S, Pressel S, Babcock C, Abernethy JD Dietary therapy slows the return of hypertension after stopping prolonged medication. *JAMA* 1985;253:657-64 [3881608]

Burr (DART 2), 2003:

Ness AR, Gallacher JE, Bennett PD, Gunnell DJ, Rogers PJ, Kessler D, Burr ML Advice to eat fish and mood: a randomised controlled trial in men with angina. *Nutr Neurosci* 2003 Feb;6:63-5 [12608739]

Burr (DART), 1989:

Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, Elwood PC, Deadman NM Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989 Sep 30;2:757-61 [2571009]

Ness AR, Whitley E, Burr ML, Elwood PC, Smith GD, Ebrahim S The long-term effect of advice to eat more fish on blood pressure in men with coronary disease: results from the diet and reinfarction trial. *J Hum Hypertens* 1999 Nov;13:729-33 [10578215]

Mate-Jimenez, 1991:**Lyon, :**

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P, Delaye J Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9 [7911176]

De Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Touboul P, Delaye J Effect of a mediterranean type of diet on the rate of cardiovascular complications in patients with coronary artery disease. Insights into the cardioprotective effect of certain nutriments. *J Am Coll Cardiol* 1996;28:1103-8 [8890801] 10.1016/S0735-1097(96)00280-X

de Lorgeril M, Renaud S, Mamelle N, Salen P, Martin JL, Monjaud I, Guidollet J, Touboul P, Delaye J Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9 [7911176]

De Lorgeril M, Salen P, Martin JL, Mamelle N, Monjaud I, Touboul P, Delaye J Effect of a mediterranean type of diet on the rate of cardiovascular complications in patients with coronary artery disease. Insights into the cardioprotective effect of certain nutriments. *J Am Coll Cardiol* 1996;28:1103-8 [8890801] 10.1016/S0735-1097(96)00280-X

PREDIMED (olive oil), 2013:

Estruch R, Ros E, Salas-Salvad J, Covas MI, D Pharm, Corella D, Ars F, Gmez-Gracia E, Ruiz-Gutierrez V, Fiol M, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pint X, Basora J, Muñoz MA, Sorlí JV, Martínez JA, Martínez-González MA Primary Prevention of Cardiovascular Disease with a Mediterranean Diet. *N Engl J Med* 2013 Feb 25;: [23432189] [10.1056/NEJMoa1200303](https://doi.org/10.1056/NEJMoa1200303)

PREDIMED (nuts), 2013:

Estruch R, Ros E, Salas-Salvad J, Covas MI, D Pharm, Corella D, Ars F, Gmez-Gracia E, Ruiz-Gutierrez V, Fiol M, Lapetra J, Lamuela-Raventos RM, Serra-Majem L, Pint X, Basora J, Muñoz MA, Sorlí JV, Martínez JA, Martínez-González MA Primary Prevention of Cardiovascular Disease with a Mediterranean Diet. *N Engl J Med* 2013 Feb 25;: [23432189] [10.1056/NEJMoa1200303](https://doi.org/10.1056/NEJMoa1200303)

Black, 1994:

Black HS, Herd JA, Goldberg LH, Wolf JE Jr, Thornby JI, Rosen T, Bruce S, Tschen JA, Foreyt JP, Scott LW Effect of a low-fat diet on the incidence of actinic keratosis. *N Engl J Med* 1994;330:1272-5 [8145782]

DART (Burr), 1989:

Burr ML, Fehily AM, Gilbert JF, Rogers S, Holliday RM, Sweetnam PM, Elwood PC, Deadman NM Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;2:757-61 [2571009]

Finnish Mental Hospital (Miettinen), 1985:

European collaborative trial of multifactorial prevention of coronary heart disease: final report on the 6-year results. World Health Organisation European Collaborative Group. *Lancet* 1986;1:869-72 [2870351]

Miettinen TA, Huttunen JK, Naukkarinen V, Strandberg T, Mattila S, Kumlin T, Sarna S Multifactorial primary prevention of cardiovascular diseases in middle-aged men. Risk factor changes, incidence, and mortality. *JAMA* 1985;254:2097-102 [4046137]

Miettinen M, Turpeinen O, Karvonen MJ, Pekkarinen M, Paavilainen E, Elosuo R Dietary prevention of coronary heart disease in women: the Finnish mental hospital study. *Int J Epidemiol* 1983;12:17-25 [6840954]

Finnish Mental Hospital (Turpeinen), 1979:

Turpeinen O, Karvonen MJ, Pekkarinen M, Miettinen M, Elosuo R, Paavilainen E Dietary prevention of coronary heart disease: the Finnish Mental Hospital Study. *Int J Epidemiol* 1979;8:99-118 [393644]

Goteborg, 1986:

Wilhelmsen L, Berglund G, Elmfeldt D, Tibblin G, Wedel H, Pennert K, Vedin A, Wilhelmsson C, Werk L The multifactor primary prevention trial in Gteborg, Sweden. *Eur Heart J* 1986;7:279-88 [3720755]

Gteborg (Wilhelmsen), 1986:

Wilhelmsen L, Berglund G, Elmfeldt D, Tibblin G, Wedel H, Pennert K, Vedin A, Wilhelmsson C, Werk L The multifactor primary prevention trial in Gteborg, Sweden. *Eur Heart J* 1986;7:279-88 [3720755]

Hjermann, 1981:

Hjermann I, Velve Byre K, Holme I, Leren P Effect of diet and smoking intervention on the incidence of coronary heart disease. Report from the Oslo Study Group of a randomised trial in healthy men. *Lancet* 1981;2:1303-10 [6118715]

Kallio, 1979:

Kallio V, Hmlinen H, Hakkila J, Luurila OJ Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. *Lancet* 1979;2:1091-4 [91836]

Los Angeles VA (Dayton), 1969:

Rogers MC Sir John Scott Burdon-Sanderson (1828-1905): a pioneer in electrophysiology. *Circulation* 1969;40:1-2 [4893441]

Dayton S, Pearce ML Diet high in unsaturated fat. A controlled clinical trial. *Minn Med* 1969;52:1237-42 [4896402]

Minnesota coronary survey (Frantz), 1975:

Frantz ID Jr, Dawson EA, Ashman PL, Gatewood LC, Bartsch GE, Kuba K, Brewer ER Test of effect of lipid lowering by diet on cardiovascular risk. The Minnesota Coronary Survey. *Arteriosclerosis* 1989;9:129-35 [2643423]

MRC low fat, 1965:

, Low-fat diet in myocardial infarction: A controlled trial. *Lancet* 1965; 2:501-4 [4158171]

MRC Soya, 1968:

, Controlled trial of soya-bean oil in myocardial infarction. *Lancet* 1968; 2:693-9 [4175085]

MRFIT, 1982:

Multiple risk factor intervention trial. Risk factor changes and mortality results. Multiple Risk Factor Intervention Trial Research Group. *JAMA* 1982;248:1465-77 [7050440]

Ornish, 1990:

Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, McLanahan SM, Kirkeeide RL, Brand RJ, Gould KL Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. *Lancet* 1990;336:129-33 [1973470]

Oslo Diet Heart Study (Leren), 1966:

Leren P, The Oslo diet-heart study. Eleven-year report. *Circulation* 1970; 42:935-42 [5477261]

Rose, 1965:

ROSE GA, THOMSON WB, WILLIAMS RT CORN OIL IN TREATMENT OF ISCHAEMIC HEART DISEASE. *Br Med J* 1965 Jun 12;1:1531-3 [14288105]

Singh, 1992:

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, Niaz MA Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. *BMJ* 1992;304:1015-9 [1586782]

STARS (St Thomas, diet), 1992:

Watts GF, Lewis B, Brunt JN, Lewis ES, Coltart DJ, Smith LD, Mann JI, Swan AV Effects on coronary artery disease of lipid-lowering diet, or diet plus cholestyramine, in the St Thomas' Atherosclerosis Regression Study (STARS) *Lancet* 1992;339:563-9 [1347091]

Veterans Ad. (Dayton), 1969:

Dayton S, Pearce ML, Hashimoto S, Dixon WJ, Tomiyasu U. A controlled clinical trial of a diet high in unsaturated fat in preventing complications of atherosclerosis. *Circulation* 1969; 40(supp 2):1-55 [0]

WHI low fat, 2005:

Howard BV, Van Horn L, Hsia J, Manson JE, Stefanick ML, Wassertheil-Smoller S, Kuller LH, LaCroix AZ, Langer RD, Lasser NL, Lewis CE, Limacher MC, Margolis KL, Mysiw WJ, Ockene JK, Parker LM, Perri MG, Phillips L, Prentice RL, Robbins J, Rossouw JE, Sarto Low-fat dietary pattern and risk of cardiovascular disease: the Women's Health Initiative Randomized Controlled Dietary Modification Trial. *JAMA* 2006 Feb 8;295:655-66 [16467234]

WHO Collaborative, 1986:

European collaborative trial of multifactorial prevention of coronary heart disease: final report on the 6-year results. World Health Organisation European Collaborative Group. *Lancet* 1986;1:869-72 [2870351]

Woodhill, 1966:

Woodhill JM, Palmer AJ, Leelarthaepin B, McGilchrist C, Blacket RB, Low fat, low cholesterol diet in secondary prevention of coronary heart disease. Adv Exp Med Biol 1978; 109:317-30 [727035]

Tuttle, 2008:

Tuttle KR, Shuler LA, Packard DP, Milton JE, Daratha KB, Bibus DM, Short RA Comparison of low-fat versus Mediterranean-style dietary intervention after first myocardial infarction (from The Heart Institute of Spokane Diet Intervention and Evaluation Trial). Am J Cardiol 2008;101:1523-30 [18489927]

3 diabetes type 2

| Trial | Treatments | Patients | Trials design and methods |
|---|---|--|--------------------------------|
| AHA 2 diet vs AHA 1 diet | | | |
| Liao , 2002 n=70 follow-up: 22 months | American Heart Association (AHA) step 2 diet (<30% of total calories as fat, <7% saturated fat, 55% carbohydrate, and <200 mg cholesterol daily) plus endurance exercise for 1 h three times a week versus AHA step 1 diet (30% of total calories as fat, 10% saturated fat, 50% carbohydrate, and <300 mg cholesterol) plus stretching exercise three times a week | Japanese American subjects with impaired glucose tolerance (WHO criteria 1998) | Parallel groups open USA |
| intensive dietary advice vs routine dietary advice | | | |
| Wein , 1999 n=200 follow-up: 4.24 y | intensive dietary advice versus routine dietary advice | women with previous gestational diabetes and currently with impaired glucose tolerance (WHO 1985 criteria) | Parallel groups open USA |

More details and results :

- prevention for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q341>
- prevention for diabetes type 2 in people with impaired glucose tolerance at <http://www.trialresultscenter.org/go-Q416>

References

Liao, 2002:

Liao D, Asberry PJ, Shofer JB, Callahan H, Matthys C, Boyko EJ, Leonetti D, Kahn SE, Austin M, Newell L, Schwartz RS, Fujimoto WY Improvement of BMI, body composition, and body fat distribution with lifestyle modification in Japanese Americans with impaired glucose tolerance. Diabetes Care 2002;25:1504-10 [12196418]

Wein, 1999:

Wein P, Beischer N, Harris C, Permezel M A trial of simple versus intensified dietary modification for prevention of progression to diabetes mellitus in women with impaired glucose tolerance. Aust N Z J Obstet Gynaecol 1999;39:162-6 [[10755770](#)]