

Clinical trials of double stenting with PES

TrialResults-center www.trialresultscenter.org

1 coronary artery disease

Trial	Treatments	Patients	Trials design and methods
double stenting with PES vs single stenting			
BBC ONE , 2008 [NCT00351260] n=250/250 follow-up: 65279;9 months	systematically stenting of both vessels with drug eluting stents(culotte or crush techniques) and with mandatory kissing balloon dilatation versus stenting of the main vessel followed by optional kissing balloon dilatation/T-stent	patients with significant coronary bifurcation lesions	Parallel groups open

More details and results :

- Drug eluting stent for coronary artery disease in bifurcation lesion at <http://www.trialresultscenter.org/go-Q214>

References

BBC ONE, 2008:

Hildick-Smith D BBC ONE British Bifurcation Coronary study Presented at TCT, Washington, 2008

Hildick-Smith D, de Belder AJ, Cooter N, Curzen NP, Clayton TC, Oldroyd KG, Bennett L, Holmberg S, Cotton JM, Glennon PE, Thomas MR, Maccarthy PA, Baumbach A, Mulvihill NT, Henderson RA, Redwood SR, Starkey IR, Stables RH Randomized trial of simple versus complex drug-eluting stenting for bifurcation lesions: the British Bifurcation Coronary Study: old, new, and evolving strategies. *Circulation* 2010;121:1235-43 [20194880] [10.1161/CIRCULATIONAHA.109.888297](https://doi.org/10.1161/CIRCULATIONAHA.109.888297)