

Clinical trials of drug-eluting stents

TrialResults-center www.trialresultscenter.org

1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
drug-eluting stents vs bare-metal stent			
DEDICATION , 2008 [NCT00192868] n=313/313 follow-up: 8 mo (15 mo, 3y)	DES currently used with or without distal protection versus BMS with or without distal protection	patients referred within 12 hours from symptom onset of an ST-elevation myocardial infarction	Factorial plan open Denmark.
PASEO , 2009 n=180/90 follow-up: 4.3 years	paclitaxel-eluting stents and sirolimus-eluting stents versus bare metal stent	patients with ST-elevation myocardial infarction within 12 hours from symptom onset	Parallel groups open

More details and results :

- myocardial revascularization for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q129>
- PCI for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q246>

References

DEDICATION, 2008:

Kelbaek H, Thuesen L, Helqvist S, Clemmensen P, Klwgaard L, Kaltoft A, Andersen B, Thuesen H, Engstrm T, Btker HE, Saunamki K, Krusell LR, Jrgensen E, Hansen HH, Christiansen EH, Ravkilde J, Kber L, Kofoed KF, Terkelsen CJ, Lassen JF Drug-eluting versus bare metal stents in patients with st-segment-elevation myocardial infarction: eight-month follow-up in the Drug Elution and Distal Protection in Acute Myocardial Infarction (DEDICATION) trial. *Circulation* 2008 Sep 9;118:1155-62 [[18725489](#)]

Kaltoft A, Kelbk H, Thuesen L, Lassen JF, Clemmensen P, Klwgaard L, Engstrm T, Btker HE, Saunamki K, Krusell LR, Jrgensen E, Tilsted HH, Christiansen EH, Ravkilde J, Kber L, Kofoed KF, Terkelsen CJ, Helqvist S Long-Term Outcome After Drug-Eluting Versus Bare-Metal Stent Implantation in Patients With ST-Segment Elevation Myocardial Infarction 3-Year Follow-Up of the Randomized DEDICATION (Drug Elution and Distal Protection in Acute Myocardial Infarction) Trial. *J Am Coll Cardiol* 2010 Jun 14;: [[20688033](#)] [10.1016/j.jacc.2010.05.009](https://doi.org/10.1016/j.jacc.2010.05.009)

PASEO, 2009:

Di Lorenzo E, De Luca G, Sauro R, Varricchio A, Capasso M, Lanzillo T, Manganelli F, Mariello C, Siano F, Pagliuca MR, Stanco G, Rosato G The PASEO (PaclitAxel or Sirolimus-Eluting Stent Versus Bare Metal Stent in Primary Angioplasty) Randomized Trial. *JACC Cardiovasc Interv* 2009 Jun;2:515-23 [[19539255](#)]

Di Lorenzo E, Sauro R, Varricchio A, Carbone G, Cortese G, Capasso M, Lanzillo T, Manganelli F, Mariello C, Siano F, Pagliuca MR, Stanco G, Rosato G, De Luca G Long-Term outcome of drug-eluting stents compared with bare metal stents in ST-segment elevation myocardial infarction: results of the paclitaxel- or sirolimus-eluting stent versus bare metal stent in Primary Angioplasty (PASEO) Randomized Trial. *Circulation* 2009;120:964-72 [19720939]

2 stable angina

Trial	Treatments	Patients	Trials design and methods
drug-eluting stents vs bare-metal stent			
ISAR-CABG <i>ongoing</i> [NCT00611910] n=NA follow-up:	DES versus BMS	Bypass Graft Lesions	open
PCI with drug-eluting stents vs CABG			
SYNTAX (diabetic) , 2010 [NCT00114972] n=NA follow-up: 1 year	paclitaxel-eluting stents versus surgical revascularization	sub group of diabetic patients with left main and/or 3-vessel disease	Parallel groups
FREEDOM , 2012 [NCT00086450] n=953/947 follow-up: 3.8 yrs (median)	percutaneous coronary stenting versus CABG	patients with diabetes and multivessel coronary artery disease	Parallel groups open international
PCI with drug-eluting stents vs CABG			
Hong , 2005 n=119/70 follow-up: 9 months	drug-eluting stents versus invasive direct coronary artery bypass (MIDCAB) surgery	proximal left anterior descending (LAD) coronary artery stenosis	Parallel groups open
VA CARDS <i>ongoing</i> [NCT00326196] n=NA follow-up:	percutaneous coronary stenting with drug eluting stents versus CABG	angiographically significant coronary artery disease in diabetes	Parallel groups open
drug-eluting stents vs CABG			
Leipzig <i>ongoing</i> [NCT00176397] n=NA follow-up:	PCI With DES versus CABG	left main coronary stenosis	

More details and results :

- myocardial revascularization for stable angina in all type of patient at <http://www.trialresultscenter.org/go-Q25>

- myocardial revascularization for stable angina in single vessel disease at <http://www.trialresultscenter.org/go-Q27>
- myocardial revascularization for stable angina in multivessels disease at <http://www.trialresultscenter.org/go-Q28>
- myocardial revascularization for stable angina in diabetic patients at <http://www.trialresultscenter.org/go-Q29>

References

ISAR-CABG, 0:

ongoing trial NCT00611910

SYNTAX (diabetic), 2010:

Banning AP, Westaby S, Morice MC, Kappetein AP, Mohr FW, Berti S, Glauber M, Kellett MA, Kramer RS, Leadley K, Dawkins KD, Serruys PW Diabetic and nondiabetic patients with left main and/or 3-vessel coronary artery disease: comparison of outcomes with cardiac surgery and paclitaxel-eluting stents. J Am Coll Cardiol 2010;55:1067-75 [[20079596](#)]

FREEDOM, 2012:

Farkouh ME, Domanski M, Sleeper LA, Siami FS, Dangas G, Mack M, Yang M, Cohen DJ, Rosenberg Y, Solomon SD, Desai AS, Gersh BJ, Magnuson EA, Lansky A, Boineau R, Weinberger J, Ramanathan K, Sousa JE, Rankin J, Bhargava B, Buse J, Hueb W, Smith CR, Muratov Strategies for Multivessel Revascularization in Patients with Diabetes. N Engl J Med 2012 Nov 4;: [[23121323](#)] [10.1056/NEJMoa1211585](#)

Hong, 2005:

Hong SJ, Lim DS, Seo HS, Kim YH, Shim WJ, Park CG, Oh DJ, Ro YM Percutaneous coronary intervention with drug-eluting stent implantation vs. minimally invasive direct coronary artery bypass (MIDCAB) in patients with left anterior descending coronary artery stenosis. Catheter Cardiovasc Interv 2005;64:75-81 [[15619278](#)]

VA CARDS, 0:

ongoing trial NCT00326196

Leipzig, 0:

ongoing trial NCT00176397

3 coronary artery disease

Trial	Treatments	Patients	Trials design and methods
drug-eluting stents vs bare-metal stent			
DEDICATION , 2008 [NCT00192868] n=313/313 follow-up: 8 mo (15 mo, 3y)	DES currently used with or without distal protection versus BMS with or without distal protection	patients referred within 12 hours from symptom onset of an ST-elevation myocardial infarction	Factorial plan open Denmark.

continued...

Trial	Treatments	Patients	Trials design and methods
PASEO , 2009 n=180/90 follow-up: 4.3 years	paclitaxel-eluting stents and sirolimus-eluting stents versus bare metal stent	patients with ST-elevation myocardial infarction within 12 hours from symptom onset	Parallel groups open
ISAR-CABG <i>ongoing</i> [NCT00611910] n=NA follow-up:	DES versus BMS	Bypass Graft Lesions	open
PCI with drug-eluting stents vs CABG			
SYNTAX (diabetic) , 2010 [NCT00114972] n=NA follow-up: 1 year	paclitaxel-eluting stents versus surgical revascularization	sub group of diabetic patients with left main and/or 3-vessel disease	Parallel groups
FREEDOM , 2012 [NCT00086450] n=953/947 follow-up: 3.8 yrs (median)	percutaneous coronary stenting versus CABG	patients with diabetes and multivessel coronary artery disease	Parallel groups open international
PCI with drug-eluting stents vs CABG			
Hong , 2005 n=119/70 follow-up: 9 months	drug-eluting stents versus invasive direct coronary artery bypass (MIDCAB) surgery	proximal left anterior descending (LAD) coronary artery stenosis	Parallel groups open
VA CARDS <i>ongoing</i> [NCT00326196] n=NA follow-up:	percutaneous coronary stenting with drug eluting stents versus CABG	angiographically significant coronary artery disease in diabetes	Parallel groups open
drug-eluting stents vs CABG			
Leipzig <i>ongoing</i> [NCT00176397] n=NA follow-up:	PCI With DES versus CABG	left main coronary stenosis	

More details and results :

- myocardial revascularization for coronary artery disease in all type of patient at <http://www.trialresultscenter.org/go-Q26>
- myocardial revascularization for coronary artery disease in diabetic patients at <http://www.trialresultscenter.org/go-Q30>
- myocardial revascularization for coronary artery disease in multivessels disease at <http://www.trialresultscenter.org/go-Q31>
- myocardial revascularization for coronary artery disease in single vessel disease at <http://www.trialresultscenter.org/go-Q32>

- Drug eluting stent for coronary artery disease in all type of patients at <http://www.trialresultscenter.org/go-Q206>
- Drug eluting stent for coronary artery disease in acute myocardial infarction at <http://www.trialresultscenter.org/go-Q208>
- Drug eluting stent for coronary artery disease in bypass graft lesion at <http://www.trialresultscenter.org/go-Q210>
- Drug eluting stent for coronary artery disease in unprotected left main artery stenosis at <http://www.trialresultscenter.org/go-Q212>

References

DEDICATION, 2008:

Kelbaek H, Thuesen L, Helqvist S, Clemmensen P, Klvgaard L, Kaltoft A, Andersen B, Thuesen H, Engstrm T, Btker HE, Saunamki K, Krusell LR, Jrgensen E, Hansen HH, Christiansen EH, Ravkilde J, Kber L, Kofoed KF, Terkelsen CJ, Lassen JF Drug-eluting versus bare metal stents in patients with st-segment-elevation myocardial infarction: eight-month follow-up in the Drug Elution and Distal Protection in Acute Myocardial Infarction (DEDICATION) trial. *Circulation* 2008 Sep 9;118:1155-62 [18725489]

Kaltoft A, Kelbk H, Thuesen L, Lassen JF, Clemmensen P, Klvgaard L, Engstrm T, Btker HE, Saunamki K, Krusell LR, Jrgensen E, Tilsted HH, Christiansen EH, Ravkilde J, Kber L, Kofoed KF, Terkelsen CJ, Helqvist S Long-Term Outcome After Drug-Eluting Versus Bare-Metal Stent Implantation in Patients With ST-Segment Elevation Myocardial Infarction 3-Year Follow-Up of the Randomized DEDICATION (Drug Elution and Distal Protection in Acute Myocardial Infarction) Trial. *J Am Coll Cardiol* 2010 Jun 14;: [20688033] 10.1016/j.jacc.2010.05.009

PASEO, 2009:

Di Lorenzo E, De Luca G, Sauro R, Varricchio A, Capasso M, Lanzillo T, Manganelli F, Mariello C, Siano F, Pagliuca MR, Stanco G, Rosato G The PASEO (PaclitAxel or Sirolimus-Eluting Stent Versus Bare Metal Stent in Primary Angioplasty) Randomized Trial. *JACC Cardiovasc Interv* 2009 Jun;2:515-23 [19539255]

Di Lorenzo E, Sauro R, Varricchio A, Carbone G, Cortese G, Capasso M, Lanzillo T, Manganelli F, Mariello C, Siano F, Pagliuca MR, Stanco G, Rosato G, De Luca G Long-Term outcome of drug-eluting stents compared with bare metal stents in ST-segment elevation myocardial infarction: results of the paclitaxel- or sirolimus-eluting stent versus bare metal stent in Primary Angioplasty (PASEO) Randomized Trial. *Circulation* 2009;120:964-72 [19720939]

ISAR-CABG, 0:

ongoing trial NCT00611910

SYNTAX (diabetic), 2010:

Banning AP, Westaby S, Morice MC, Kappetein AP, Mohr FW, Berti S, Glauber M, Kellett MA, Kramer RS, Leadley K, Dawkins KD, Serruys PW Diabetic and nondiabetic patients with left main and/or 3-vessel coronary artery disease: comparison of outcomes with cardiac surgery and paclitaxel-eluting stents. *J Am Coll Cardiol* 2010;55:1067-75 [20079596]

FREEDOM, 2012:

Farkouh ME, Domanski M, Sleeper LA, Siami FS, Dangas G, Mack M, Yang M, Cohen DJ, Rosenberg Y, Solomon SD, Desai AS, Gersh BJ, Magnuson EA, Lansky A, Boineau R, Weinberger J, Ramanathan K, Sousa JE, Rankin J, Bhargava B, Buse J, Hueb W, Smith CR, Muratov Strategies for Multivessel Revascularization in Patients with Diabetes. *N Engl J Med* 2012 Nov 4;: [23121323] 10.1056/NEJMoa1211585

Hong, 2005:

Hong SJ, Lim DS, Seo HS, Kim YH, Shim WJ, Park CG, Oh DJ, Ro YM Percutaneous coronary intervention with drug-eluting stent implantation vs. minimally invasive direct coronary artery bypass (MIDCAB) in patients with left anterior descending coronary artery stenosis. *Catheter Cardiovasc Interv* 2005;64:75-81 [15619278]

VA CARDS, 0:

ongoing trial NCT00326196

Leipzig, 0:

ongoing trial NCT00176397

Entry terms: CYPHER, TAXUS, Promus, Xience, Biomatrix, Nobori, Endeavor, Drug-Eluting Stents, Drug Eluting Stents, Drug-Eluting Stent, Drug-Coated Stents, Drug Coated Stents, Drug-Coated Stent,