

Clinical trials of early amiodarone

TrialResults-center www.trialresultscenter.org

1 post myocardial infarction

Trial	Treatments	Patients	Trials design and methods
early amiodarone vs control			
BASIS , 1990 n=98/14 follow-up: 65279;12 mo	amiodarone 1 g for 5 d; then 200 mg/d started within 4 weeks of AMI versus no amiodarone (usual care)	patients with persisting asymptomatic complex arrhythmias after myocardial infarction (Lown class 3 or 4b in >2 of 24 h)	Parallel groups open
Navarro-Lopez , 1993 n=115/123 follow-up: 24 mo	amiodarone 600 mg/d for 1 week, 400 mg/d for 1 week then 200 mg/d started 10-30 d after AMI versus no amiodarone	patients who have had MI with a left ventricular ejection fraction of 20 to 45% and >or = 3 ventricular premature complexes per hour (pairs or runs) - 3 VPOs/h, pairs or runs of VT	Parallel groups open Spain
early amiodarone vs placebo			
CAMIAT , 1991 n=NA follow-up: 24 mo	amiodarone 10 mg/kg per d for 3 weeks then 300-400 mg/d started 6-45 d after AMI versus placebo	patients with acute myocardial infarction within the previous 6-30 days and >10 VPDs/h for 18 h or a run of VT	Parallel groups double blind
Ceremuzynski , 1992 n=305/308 follow-up: 12 mo	amiodarone 800 mg/d for 1 week then 200-400 mg/d started 5-7 d after AMI versus placebo	No need for antiarrhythmic therapy	Parallel groups double blind
Hockings , 1987 n=59/70 follow-up: 642 mo	amiodarone 200 mg 3 times daily for 1 wk; then 200 mg/d started <8-10 d after AMI versus placebo	patients with AMI - Absence of VF or VT >3 beats	Parallel groups double blind

More details and results :

- antiarrhythmic drugs for post myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q251>

References

BASIS, 1990:

Burkart F, Pfisterer M, Kiowski W, Follath F, Burckhardt D Effect of antiarrhythmic therapy on mortality in survivors of myocardial infarction with asymptomatic complex ventricular arrhythmias: Basel Antiarrhythmic Study of Infarct Survival (BASIS) J Am Coll Cardiol 1990;16:1711-8 [2254558]

Navarro-Lopez, 1993:

Navarro-Lopez F, Cosin J, Marrugat J, Guindo J, Bayes de Luna A Comparison of the effects of amiodarone versus metoprolol on the frequency of ventricular arrhythmias and on mortality after acute myocardial infarction. SSSD Investigators. Spanish Study on Sudden Death. Am J Cardiol 1993;72:1243-8 [7504880]

CAMIAT , 1991:

Cairns JA, Connolly SJ, Gent M, Roberts R Post-myocardial infarction mortality in patients with ventricular premature depolarizations. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Pilot Study. Circulation 1991;84:550-7 [1860199]

Ceremuzynski, 1992:

Ceremuzynski L, Kleczar E, Krzeminska-Pakula M, Kuch J, Nartowicz E, Smielak-Korombel J, Dyduzynski A, Maciejewicz J, Zaleska T, Lazarczyk-Kedzia E Effect of amiodarone on mortality after myocardial infarction: a double-blind, placebo-controlled, pilot study. J Am Coll Cardiol 1992;20:1056-62 [1401602]

Hockings, 1987:

Hockings BE, George T, Mahrous F, Taylor RR, Hajar HA Effectiveness of amiodarone on ventricular arrhythmias during and after acute myocardial infarction. Am J Cardiol 1987;60:967-70 [3673913]

Entry terms: amiodarone, Amiobeta, Cordarone, Cordarex, Amiodarex, Kordaron, Trangorex, Amiodarona, Amiohexal, Braxan, Corbionax, Ortacrone, Rytmarone, Tachydaron, Aratac