

# Clinical trials of elastic compression stockings

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 postthrombotic syndrome

Trial	Treatments	Patients	Trials design and methods
<b>elastic compression stockings vs no compression</b>			
<a href="#">Partsch , 2004</a> n=36/17 follow-up:	compression stockings or Unna boot bandages plus walking versus no compression	patients with acute proximal DVT	Parallel groups open
<b>elastic compression stockings vs placebo</b>			
<a href="#">Ginsberg , 2001</a> n=42/40 follow-up: 57 months	below-knee stocking (20-30 mm Hg) versus placebo stocking	patients with proximal deep venous thrombosis	Parallel groups double-blind

More details and results :

- prevention for postthrombotic syndrome in all type of patients at <http://www.trialresultscenter.org/go-Q436>

## References

### Partsch, 2004:

Partsch H, Kaulich M, Mayer W Immediate mobilisation in acute vein thrombosis reduces post-thrombotic syndrome. Int Angiol 2004;23:206-12 [[15765034](#)]

### Ginsberg, 2001:

Ginsberg JS, Hirsh J, Julian J, Vander LaandeVries M, Magier D, MacKinnon B, Gent M Prevention and treatment of postphlebotic syndrome: results of a 3-part study. Arch Intern Med 2001;161:2105-9 [[11570939](#)]