

# Clinical trials of eplerenone

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 heart failure

Trial	Treatments	Patients	Trials design and methods
<b>eplerenone vs placebo</b>			
<b>EMPHASIS-HF , 2010</b> [NCT00232180] n=1364/1373 follow-up: 21 months	eplerenone versus placebo	patients with New York Heart Association class II heart failure and an ejection fraction of no more than 35%	Parallel groups double blind 29 countries
<b>EPHESUS , 2003</b> n=3319/3313 follow-up: 16 mo (mean, range 0 to 33)	eplerenone 25 mg per day initially, titrated to a maximum of 50 mg per day versus placebo	patients with acute myocardial infarction complicated by left ventricular dysfunction and heart failure	Parallel groups Double blind 37 countries
<b>REMODEL ongoing</b> [NCT00082589] n=NA follow-up:	Eplerenone versus placebo	Patients with left ventricular systolic dysfunction (EF Less Than or Equal to 35% ) and mild to moderate heart failure	Parallel groups double blind
<b>Weir ongoing</b> [NCT00132093] n=NA follow-up: 6 montsh	eplerenone versus placebo	patients with acute myocardial infarction	Parallel groups double blind

More details and results :

- diuretics for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q75>
- diuretics for heart failure in Left Ventricular Dysfunction after Myocardial Infarction at <http://www.trialresultscenter.org/go-Q76>
- aldosterone blockade for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q488>

## References

### EMPHASIS-HF, 2010:

Zannad F, McMurray JJ, Drexler H, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pitt B Rationale and design of the Eplerenone in Mild Patients Hospitalization And SurvIval Study in Heart Failure (EMPHASIS-HF). Eur J Heart Fail 2010;12:617-22 [20388647] [10.1093/eurjhf/hfq049](https://doi.org/10.1093/eurjhf/hfq049)

Zannad F, McMurray JJ, Drexler H, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pitt B Rationale and design of the Eplerenone in Mild Patients Hospitalization And SurvIval Study in Heart Failure (EMPHASIS-HF). Eur J Heart Fail 2010;12:617-22 [20388647] [10.1093/eurjhf/hfq049](https://doi.org/10.1093/eurjhf/hfq049)

Zannad F, McMurray JJ, Krum H, van Veldhuisen DJ, Swedberg K, Shi H, Vincent J, Pocock SJ, Pitt B Eplerenone in Patients with Systolic Heart Failure and Mild Symptoms. N Engl J Med 2010 Nov 14;: [21073363] 10.1056/NEJMoa1009492

**EPHESUS, 2003:**

Pitt B, White H, Nicolau J, Martinez F, Gheorghide M, Aschermann M, van Veldhuisen DJ, Zannad F, Krum H, Mukherjee R, Vincent J Eplerenone reduces mortality 30 days after randomization following acute myocardial infarction in patients with left ventricular systolic dysfunction and heart failure. J Am Coll Cardiol 2005 Aug 2;46:425-31 [16053953]

Pitt B, Remme W, Zannad F, Neaton J, Martinez F, Roniker B, Bittman R, Hurley S, Kleiman J, Gatlin M Eplerenone, a selective aldosterone blocker, in patients with left ventricular dysfunction after myocardial infarction. N Engl J Med 2003 Apr 3;348:1309-21 [12668699]

**REMODEL, :**

ongoing trial NCT00082589

**Weir, :**

ongoing trial NCT00132093

Entry terms: eplerenon, Inspra