

Clinical trials of evolocumab

TrialResults-center www.trialresultscenter.org

1 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
evolocumab vs			
Mendel 1 , 2012 [NCT01375777] n=NA follow-up:	-	-	
MENDEL 2 [NCT01763827] n=NA	-	-	
YUKAWA-1 , 2014 n=NA follow-up:	-	-	
evolocumab vs ezetimibe alone			
GAUSS 2 [NCT01763905] n=102/205 follow-up:	evolocumab 140 mg every two weeks (Q2W) or evolocumab 420 mg once monthly (QM) versus ezetimibe 10 mg	patients with statin intolerance	
evolocumab vs placebo			
GAUSS 1 , 2012 [NCT01375764] n=95/32 follow-up:	-	statin-intolerant patients	
evolocumab vs placebo (on top statins)			
DESCARTES , 2014 [NCT01516879] n=599/302 follow-up: 52 weeks	evolocumab (420 mg) every 4 weeks versus placebo	-	
FOURIER , 2017 [NCT01764633] n=NA follow-up: 2.2 years	evolocumab (either 140 mg every 2 weeks or 420 mg monthly) versus placebo	patients with atherosclerotic cardiovascular disease and LDL cholesterol levels of 70 mg per deciliter (1.8 mmol per liter) or higher who were receiving statin therapy	Parallel groups double-blind

continued...

Trial	Treatments	Patients	Trials design and methods
LAPLACE 2 , 2014 [NCT01763866] n=1117/558 follow-up:	evolucumab + statin versus placebo + statin	-	
LAPLACE-TIMI 57 [NCT01380730] n=NA follow-up:	subcutaneous injections of AMG 145 70 mg, 105 mg, or 140 mg, versus placebo	-	
RUTHERFORD-1 [NCT01375751] n=111/56 follow-up:	AMG 145 350 mg, AMG 145 420 mg versus placebo	heterozygous familial hypercholesterolemia patients	
RUTHERFORD-2 , 2015 [NCT01763918] n=220/109 follow-up:	subcutaneous evolucumab 140 mg every 2 weeks, evolucumab 420 mg monthly versus placebo	heterozygous familial hypercholesterolaemia	

More details and results :

- PCSK9 Inhibitors for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q599>

References

Mendel 1, 2012:

Koren MJ, Scott R, Kim JB, Knusel B, Liu T, Lei L, Bolognese M, Wasserman SM Efficacy, safety, and tolerability of a monoclonal antibody to proprotein convertase subtilisin/kexin type 9 as monotherapy in patients with hypercholesterolaemia (MENDEL): a randomised, double-blind, placebo-controlled, phase 2 study. *Lancet* 2012 Dec 8;380:1995-2006 [23141812] [10.1016/S0140-6736\(12\)61771-1](https://doi.org/10.1016/S0140-6736(12)61771-1)

MENDEL 2, :

YUKAWA-1, 2014:

Hirayama A, Honarpour N, Yoshida M, Yamashita S, Huang F, Wasserman SM, Teramoto T Effects of evolucumab (AMG 145), a monoclonal antibody to PCSK9, in hypercholesterolemic, statin-treated Japanese patients at high cardiovascular risk—primary results from the phase 2 YUKAWA study. *Circ J* 2014;78:1073-82 [24662398]

GAUSS 2, :

Stroes E, Colquhoun D, Sullivan D, Civeira F, Rosenson RS, Watts GF, Bruckert E, Cho L, Dent R, Knusel B, Xue A, Scott R, Wasserman SM, Rocco M Anti-PCSK9 antibody effectively lowers cholesterol in patients with statin intolerance: the GAUSS-2 randomized, placebo-controlled phase 3 clinical trial of evolucumab. *J Am Coll Cardiol* 2014 Jun 17;63:2541-8 [24694531]

GAUSS 1, 2012:

Sullivan D, Olsson AG, Scott R, Kim JB, Xue A, GebSKI V, Wasserman SM, Stein EA Effect of a monoclonal antibody to PCSK9 on low-density lipoprotein cholesterol levels in statin-intolerant patients: the GAUSS randomized trial. *JAMA* 2012 Dec 19;308:2497-506 [23128163]

DESCARTES, 2014:

Blom DJ, Hala T, Bolognese M, Lillestol MJ, Toth PD, Burgess L, Ceska R, Roth E, Koren MJ, Ballantyne CM, Monsalvo ML, Tsirtsonis K, Kim JB, Scott R, Wasserman SM, Stein EA A 52-week placebo-controlled trial of evolocumab in hyperlipidemia. *N Engl J Med* 2014 May 8;370:1809-19 [24678979]

FOURIER, 2017:

Sabatine MS, Giugliano RP, Keech AC, Honarpour N, Wiviott SD, Murphy SA, Kuder JF, Wang H, Liu T, Wasserman SM, Sever PS, Pedersen TR Evolocumab and Clinical Outcomes in Patients with Cardiovascular Disease. *N Engl J Med* 2017;376:1713-1722 [28304224]

LAPLACE 2, 2014:

Robinson JG, Nedergaard BS, Rogers WJ, Fialkow J, Neutel JM, Ramstad D, Somaratne R, Legg JC, Nelson P, Scott R, Wasserman SM, Weiss R Effect of evolocumab or ezetimibe added to moderate- or high-intensity statin therapy on LDL-C lowering in patients with hypercholesterolemia: the LAPLACE-2 randomized clinical trial. *JAMA* 2014 May 14;311:1870-82 [24825642]

LAPLACE-TIMI 57, :

Giugliano RP, Desai NR, Kohli P, Rogers WJ, Somaratne R, Huang F, Liu T, Mohanavelu S, Hoffman EB, McDonald ST, Abrahamsen TE, Wasserman SM, Scott R, Sabatine MS Efficacy, safety, and tolerability of a monoclonal antibody to proprotein convertase subtilisin/kexin type 9 in combination with a statin in patients with hypercholesterolaemia (LAPLACE-TIMI 57): a randomised, placebo-controlled, dose-ranging, phase 2 study. *Lancet* 2012 Dec 8;380:2007-17 [23141813]

RUTHERFORD-1, :

Raal F, Scott R, Somaratne R, Bridges I, Li G, Wasserman SM, Stein EA Low-density lipoprotein cholesterol-lowering effects of AMG 145, a monoclonal antibody to proprotein convertase subtilisin/kexin type 9 serine protease in patients with heterozygous familial hypercholesterolemia: the Reduction of LDL-C with PCSK9 Inhibition in Heterozygous Familial Hypercholesterolemia Disorder (RUTHERFORD) randomized trial. *Circulation* 2012 Nov 13;126:2408-17 [23129602]

RUTHERFORD-2, 2015:

Raal FJ, Stein EA, Dufour R, Turner T, Civeira F, Burgess L, Langslet G, Scott R, Olsson AG, Sullivan D, Hovingh GK, Cariou B, Gouni-Berthold I, Somaratne R, Bridges I, Scott R, Wasserman SM, Gaudet D PCSK9 inhibition with evolocumab (AMG 145) in heterozygous familial hypercholesterolaemia (RUTHERFORD-2): a randomised, double-blind, placebo-controlled trial. *Lancet* 2015 Jan 24;385:331-40 [25282519]

Entry terms: AMG 145, evolocumab, AMG-145, Repatha