

Clinical trials of felodipine or israpidine

TrialResults-center www.trialresultscenter.org

1 hypertension

Trial	Treatments	Patients	Trials design and methods
felodipine or israpidine vs diuretic or beta-blocker			
STOP-2 (CCB vs diuretic or beta-blocker) , 1999 n=2196/2213 follow-up: up to 6 years	felodipine 25 mg or israpidine 25 mg daily versus conventional antihypertensivedrugs (atenolol 50 mg, metoprolol 100 mg,pindolol 5 mg, or hydrochlorothiazide 25 mg plus amiloride2.5 mg daily	patients aged 70-84 years with hypertension (blood pressure >180 mm Hg systolic, >105 mm Hg diastolic, or both).	Parallel groups Open

More details and results :

- anti hypertensive agents for hypertension in elderly (60 years and more) at <http://www.trialresultscenter.org/go-Q9>
- anti hypertensive agents for hypertension in diabetic patients at <http://www.trialresultscenter.org/go-Q10>
- anti hypertensive agents for hypertension in all type of patient at <http://www.trialresultscenter.org/go-Q13>
- anti hypertensive agents for hypertension in uncomplicated hypertension at <http://www.trialresultscenter.org/go-Q685>

References

STOP-2 (CCB vs diuretic or beta-blocker), 1999:

Hansson L, Lindholm LH, Ekblom T, Dahlöf B, Lanke J, Schersten B, Wester PO, Hedner T, de Faire U Randomised trial of old and new antihypertensive drugs in elderly patients: cardiovascular mortality and morbidity the Swedish Trial in Old Patients with Hypertension-2 study. Lancet 1999 Nov 20;354:1751-6 [[10577635](#)]