

# Clinical trials of fenofibrate

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
<b>fenofibrate vs placebo</b>			
<b>FIELD , 2005</b> [ISRCTN64783481] n=4895/4900 follow-up: 5y	fenofibrate 200 mg daily versus placebo	aged 50-75 years, with type 2 diabetes mellitus, and not taking statin therapy at study entry	
<b>DAIS , 2001</b> n=207/211 follow-up: 3.3 years	fenofibrate 200 mg/day versus placebo	men and women with type 2 diabetes and coronary atherosclerosis	Parallel groups double-blind Canada, Finland, France, Sweden
<b>FIELD , 2005</b> [ISRCTN64783481] n=4895/4900 follow-up: 5 years	fenofibrate 200mg/d versus Placebo	participants aged 50-75 years, with type 2 diabetes mellitus, and not taking statin therapy at study entry	Parallel groups double blind Australia, New Zealand, Finland
<b>fenofibrate vs placebo (on top simvastatine)</b>			
<b>ACCORD lipid , 2010</b> [NCT00000620] n=2765/2753 follow-up: 4.7y	fenofibrate on top simvastatin versus placebo (on top simvastatine)	high-risk patients with type 2 diabetes	Factorial plan double-blind United States and Canada

More details and results :

- cholesterol lowering intervention for cardiovascular prevention in diabetic patients at <http://www.trialresultscenter.org/go-Q6>
- cholesterol lowering intervention for cardiovascular prevention in all chronic situations at <http://www.trialresultscenter.org/go-Q154>
- HDL increasing drugs for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q503>

## References

**FIELD, 2005:**

Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005;366:1849-61 [[16310551](#)]

**DAIS, 2001:**

Effect of fenofibrate on progression of coronary-artery disease in type 2 diabetes: the Diabetes Atherosclerosis Intervention Study, a randomised study. *Lancet* 2001;357:905-10 [[11289345](#)]

**FIELD, 2005:**

The need for a large-scale trial of fibrate therapy in diabetes: the rationale and design of the Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study. [*ISRCTN64783481*]. *Cardiovasc Diabetol* 2004 Dec 1;3:9 [[15571637](#)]

Scott R, Best J, Forder P, Taskinen MR, Simes J, Barter P, Keech A Fenofibrate Intervention and Event Lowering in Diabetes (FIELD) study: baseline characteristics and short-term effects of fenofibrate [*ISRCTN64783481*]. *Cardiovasc Diabetol* 2005 Aug 22;4:13 [[16111499](#)]  
*Lancet* 2005

Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005 Nov 26;366:1849-61 [[16310551](#)]

Hiukka A, Westerbacka J, Leinonen ES, Watanabe H, Wiklund O, Hulten LM, Salonen JT, Tuomainen TP, Yki-Jrvinen H, Keech AC, Taskinen MR Long-term effects of fenofibrate on carotid intima-media thickness and augmentation index in subjects with type 2 diabetes mellitus. *J Am Coll Cardiol* 2008 Dec 16;52:2190-7 [[19095138](#)] [10.1016/j.jacc.2008.09.049](#)

Hiukka A, Westerbacka J, Leinonen ES, Watanabe H, Wiklund O, Hulten LM, Salonen JT, Tuomainen TP, Yki-Jrvinen H, Keech AC, Taskinen MR Long-term effects of fenofibrate on carotid intima-media thickness and augmentation index in subjects with type 2 diabetes mellitus. *J Am Coll Cardiol* 2008 Dec 16;52:2190-7 [[19095138](#)]

Rajamani K, Colman PG, Li LP, Best JD, Voysey M, D'Emden MC, Laakso M, Baker JR, Keech AC Effect of fenofibrate on amputation events in people with type 2 diabetes mellitus (FIELD study): a prespecified analysis of a randomised controlled trial. *Lancet* 2009 May 23;373:1780-8 [[19465233](#)] [10.1016/S0140-6736\(09\)60698-X](#)

**ACCORD lipid, 2010:**

Effects of Combination Lipid Therapy in Type 2 Diabetes Mellitus. *N Engl J Med* 2010 Mar 14;: [[20228404](#)] [10.1056/NEJMoa1001282](#)

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. *N Engl J Med* 2010 Jun 29;: [[20587587](#)] [10.1056/NEJMoa1001288](#)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010 Jun 29;: [[20594588](#)] [10.1016/S0140-6736\(10\)60576-4](#)

## 2 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
fenofibrate vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
<b>FIELD , 2005</b> [ISRCTN64783481] n=4895/4900 follow-up: 5y	fenofibrate 200 mg daily versus placebo	aged 50-75 years, with type 2 diabetes mellitus, and not taking statin therapy at study entry	
<b>fenofibrate vs placebo (on top simvastatine)</b>			
<b>ACCORD lipid , 2010</b> [NCT00000620] n=2765/2753 follow-up: 4.7y	fenofibrate on top simvastatin versus placebo (on top simvastatine)	high-risk patients with type 2 diabetes	Factorial plan double-blind United States and Canada

More details and results :

- cholesterol lowering intervention for diabetes type 2 in diabetic patients with or without hypercholesterolemia at <http://www.trialresultscenter.org/go-Q85>

## References

### FIELD, 2005:

Keetch A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forder P, Pillai A, Davis T, Glasziou P, Drury P, Kesniemi YA, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): randomised controlled trial. *Lancet* 2005;366:1849-61 [[16310551](#)]

### ACCORD lipid, 2010:

Effects of Combination Lipid Therapy in Type 2 Diabetes Mellitus. *N Engl J Med* 2010 Mar 14;: [[20228404](#)] [10.1056/NEJMoa1001282](https://doi.org/10.1056/NEJMoa1001282)

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. *N Engl J Med* 2010 Jun 29;: [[20587587](#)] [10.1056/NEJMoa1001288](https://doi.org/10.1056/NEJMoa1001288)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010 Jun 29;: [[20594588](#)] [10.1016/S0140-6736\(10\)60576-4](https://doi.org/10.1016/S0140-6736(10)60576-4)

Entry terms: Procetofen