

# Clinical trials of flecainide

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 atrial fibrillation

Trial	Treatments	Patients	Trials design and methods
<b>flecainide vs no treatment</b>			
<a href="#">Van Gelder , 1989</a> n=36/37 follow-up: 12 months	Flecainide 200-300 mg/d versus no treatment	Any persistent AF or AFl	Parallel groups open
<b>flecainide vs placebo</b>			
<a href="#">Carunchio (flecainide vs placebo) , 1995</a> n=20/26 follow-up: 12 months	Flecainide 200 mg/d versus placebo	patients with recurrent AF with >3 episodes in previous 1 year	Parallel groups open
<b>flecainide vs digoxin</b>			
<a href="#">Steinbeck (flecainide vs digoxin) , 1988</a> n=15/15 follow-up: 12 months	-	Paroxysmal symptomatic AF of any duration	Parallel groups open
<b>flecainide vs propafenone</b>			
<a href="#">Aliot , 1996</a> n=48/49 follow-up: 12 months	Flecainide 100-200mg/d versus Propafenone 600 mg/d	patients with paroxysmal AF documented any time before	Parallel groups open
<b>flecainide vs quinidine</b>			
<a href="#">Naccarelli , 1996</a> n=117/122 follow-up: 12 months	Flecainide 200-300 mg/d versus Quinidine 1-1,5 g/d	Any documented symptomatic AF	Parallel groups double blind

More details and results :

- antiarrhythmic drugs for atrial fibrillation in maintaining sinus rhythm after cardioversion at <http://www.trialresultscenter.org/go-Q113>

## References

Van Gelder, 1989:

Van Gelder IC, Crijns HJ, Van Gilst WH, Van Wijk LM, Hamer HP, Lie KI Efficacy and safety of flecainide acetate in the maintenance of sinus rhythm after electrical cardioversion of chronic atrial fibrillation or atrial flutter. *Am J Cardiol* 1989;64:1317-21 [2511744]

Abstracts from the 61st scientific sessions. American Heart Association. Washington, DC, November 14-17, 1988. *Circulation* 1988;78:IIA-IIIG, III-II775 [3168202]

**Carunchio (flecainide vs placebo), 1995:**

Carunchio A, Fera MS, Mazza A, Burattini M, Greco G, Galati A, Ceci V [A comparison between flecainide and sotalol in the prevention of recurrences of paroxysmal atrial fibrillation] *G Ital Cardiol* 1995;25:51-68 [7642012]

**Steinbeck (flecainide vs digoxin), 1988:**

**Aliot, 1996:**

Aliot E, Denjoy I Comparison of the safety and efficacy of flecainide versus propafenone in hospital out-patients with symptomatic paroxysmal atrial fibrillation/flutter. The Flecainide AF French Study Group. *Am J Cardiol* 1996;77:66A-71A [8607394]

**Naccarelli, 1996:**

Naccarelli GV, Dorian P, Hohnloser SH, Coumel P Prospective comparison of flecainide versus quinidine for the treatment of paroxysmal atrial fibrillation/flutter. The Flecainide Multicenter Atrial Fibrillation Study Group. *Am J Cardiol* 1996;77:53A-59A [8607392]

Entry terms: Flecainide, Flecatap, Flecainide Acetate, Flecainide Monoacetate, R818, Tambocor, Apocard, Flcane, Flecadura, Flecainid-Isis, Flecainid Isis,