

# Clinical trials of insulin glulisine + glargine

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 diabetes type 2

| Trial  | Treatments   | Patients  | Trials design and methods                       |
|--|--|---|---|
| <b>insulin glulisine + glargine vs glargine once daily</b> |  |   |   |
| <b>Owens , 2011</b><br>n=49/57<br>follow-up: 3 months      | basal+bolus (single dose of insulin glulisine immediately prior to the main meal)<br>versus<br>basal insulin (glargin) | patients withT2DM using any basal insulin and HbA1c >7.0% after 3-month of insulin glargine titrated to optimize fasting bloodglucose control | Parallel groups<br>open-label<br>US, UK, Russia |

More details and results :

- insulin therapy for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q548>

## References

### Owens, 2011:

Owens DR, Luzio SD, Sert-Langeron C, Riddle MC Effects of initiation and titration of a single pre-prandial dose of insulin glulisine while continuing titrated insulin glargine in type 2 diabetes: a 6-month 'proof-of-concept' study. Diabetes Obes Metab 2011;13:1020-7 [21679291] [10.1111/j.1463-1326.2011.01459.x](https://doi.org/10.1111/j.1463-1326.2011.01459.x)

Entry terms: glargine