

Clinical trials of intensive

TrialResults-center www.trialresultscenter.org

1 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
intensive lipid-lowering therapy vs diet			
FATS , 1990 [NCT00000512] n=94/52 follow-up: 2.5 years	intensive lipid-lowering therapy with various drugs versus placebo	men no more than 62 years of age who had apolipoprotein B levels greater than or equal to 125 mg per deciliter, documented coronary artery disease, and a family history of vascular disease	Parallel groups open Japan

More details and results :

- cholesterol lowering intervention for cardiovascular prevention in all chronic situations at <http://www.trialresultscenter.org/go-Q154>

References

FATS, 1990:

Brown G, Albers JJ, Fisher LD, Schaefer SM, Lin JT, Kaplan C, Zhao XQ, Bisson BD, Fitzpatrick VF, Dodge HT Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. N Engl J Med 1990 Nov 8;323:1289-98 [2215615]

2 hypertension

Trial	Treatments	Patients	Trials design and methods
intensive vs usual			
ACCORD (blood pressure) , 2010 [NCT00000620] n=2363/2371 follow-up: 4.7 y	intensive blood-pressure control, targeting a systolic pressure of less than 120 mm Hg versus standard blood-pressure control	high-risk patients with type 2 diabetes, high HbA1c concentrations (>7.5%), and cardiovascular disease (or ≥ 2 cardiovascular risk factors)	Factorial plan open United States, Canada
more intensive blood pressure lowering strategie vs less intensive blood pressure lowering strategie			

continued...

Trial	Treatments	Patients	Trials design and methods
PAST-BP , 2015 n=NA	-	-	
Wei , 2013 n=NA follow-up: 4 years (mean)	BP <=140/90 mm Hg versus BP <=150/90 mm Hg	Chinese hypertensive patients older than 70 years	Parallel groups China
SPS3 , 2013 [NCT00059306.] n=NA follow-up:	less than 130 mm Hg versus 130-149 mm Hg	patients lived in North America, Latin America, and Spain and had recent, MRI-defined symptomatic lacunar infarctions	Parallel groups open-label
HOMED-BP , 2012 n=NA follow-up: 5.3 years (median)	tight control (<125/<80 mm Hg (TC)) of HBP versus usual control (125-134/80-84 mm Hg (UC))	with an untreated systolic/diastolic HBP of 135-179/85-119 mm Hg	Parallel groups
VANLISH , 2010 n=NA follow-up: 3.07 years (median)	strict blood pressure control (<140 mm Hg) versus moderate blood pressure control (>or =140 mm Hg to <150 mm Hg)	patients aged 70 to 84 years with isolated systolic hypertension (sitting blood pressure 160 to 199 mm Hg)	Parallel groups open-label
JATOS , 2008 n=2212/2206 follow-up:	strict treatment to maintain systolic blood pressure below 140 mmHg versus mild treatment to maintain systolic blood pressure below 160 but at or above 140 mmHg	elderly hypertensive patients with essential hypertension (65-85 years old, with a pretreatment systolic blood pressure of above 160 mmHg)	Parallel groups open-label
UKPDS-HDS , 1998 n=758/390 follow-up: 8.4 years	blood pressure of <150/85 mm Hg (with the use of an angiotensin converting enzyme inhibitor captopril or a beta blocker atenolol as main treatment) versus less tight control aiming at a blood pressure of <180/105 mm Hg	patients with type 2 diabetes	Parallel groups open-label UK
SPRINT , 2015 [NCT01206062] n=4678/4683 follow-up:	target of 120 mm Hg versus target of 140 mm Hg	high-risk hypertensive adults 50 years of age and older with one additional cardiovascular risk factor or preexisting kidney disease	Parallel groups open
Cardio-Sis , 2009 [NCT00421863] n=558/553 follow-up: 2 years	tighter control of systolic BP with a goal of <130 mm Hg versus usual control, with a goal of <140 mm Hg	nondiabetic patients with hypertension and with SBP of 150 mm Hg or higher confirmed at two different times	Parallel groups open Italy

continued...

Trial	Treatments	Patients	Trials design and methods
AASK , 2002 n=540/554 follow-up: (range 3-6.4y)	arterial pressure goal of 92 mm Hg or lower versus usual mean arterial pressure goal of 102 to 107 mm Hg/pj	African-Americans,with diastolic blood pressure higher than 94mmHg and a glomerular filtration rate between 20 and 65 ml/min per 1.73 m2	Parallel groups open USA
ABCD target (H) , 2000 n=237/233 follow-up: 5 year	intensive treatment with a diastolic blood pressure goal of 75 mmHg versus moderate treatment with a diastolic blood pressure goal of 80-89 mmHg	diabetes patients with DBP >=90 mmHg	Parallel groups open
ABCD target (N) , 2002 n=237/243 follow-up:	intensive treatment (diastolic blood pressure decrease of 10 mmHg below baseline DBP) versus moderate treatment (diastolic blood pressure goal of 80-89 mmHg)	diabetes patients with diastolic blood pressure between 80 and 89mmHg	Parallel groups open
HOT , 1994 n=12526/6264 follow-up: 3.8 y	less or equal than 85 mmHg, or less or equal than 80 mmHg versus less or equal than 90 mmHg	patients with diastolic blood pressure between 100 mmHg and 115 mmHg	Factorial plan open 26 countries
REIN-2 , 2005 n=169/169 follow-up: 36 months	intensified (systolic/diastolic <130/80 mm Hg) blood-pressure control versus conventional (diastolic <90 mm Hg) blood-pressure control	patients with non-diabetic proteinuric nephropathies receiving background treatment with the ACE inhibitor ramipril	open
MDRD , 1994 n=840 follow-up: 2.2 y	low target blood pressure (mean arterial pressure <92 mm Hg) versus usual target blood pressure (mean arterial pressure <107 mm Hg)	patients with predominantly nondiabetic kidney disease and a glomerular filtration rate of 13 to 55 mL/min per 1.73 m2	open
Toto , 1995 n=42/35 follow-up:	strict blood pressure control (DBP 65 to 80 mm Hg) versus usual blood pressure control (DBP 85 to 95 mm Hg)	non-diabetic patients (age 25 to 73) with long-standing hypertension (DBP >or = 95 mm Hg), chronic renal insufficiency (GFR <or = 70 m/min/1.73 m2) and a normal urine sediment	open
ACCORD blood pressure , 2008 [NCT00000620] n=2362/2371 follow-up: 4.7y	intensive therapy, targeting a systolic pressure of less than 120 mm Hg versus standard therapy, targeting a systolic pressure of less than 140 mm Hg	patients with a median glyated hemoglobin level of 8.1% at high risk for cardiovascular events	Factorial plan open USA, Canada

continued...

Trial	Treatments	Patients	Trials design and methods
ESH-CHL-SHOT [NCT01563731] n=NA	<i>ongoing</i> -	-	

More details and results :

- anti hypertensive agents for hypertension in diabetic patients at <http://www.trialresultscenter.org/go-Q10>
- intensive blood pressure control for hypertension in all type of patients at <http://www.trialresultscenter.org/go-Q336>
- intensive blood pressure control for hypertension in diabetic patients at <http://www.trialresultscenter.org/go-Q343>
- intensive blood pressure control for hypertension in non diabetic patients at <http://www.trialresultscenter.org/go-Q344>
- intensive blood pressure control for hypertension in patients with chronic kidney disease at <http://www.trialresultscenter.org/go-Q495>

References

ACCORD (blood pressure), 2010:

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. N Engl J Med 2010 Jun 29;: [20587587] [10.1056/NEJMoa1001288](https://doi.org/10.1056/NEJMoa1001288)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. Lancet 2010 Jun 29;: [20594588] [10.1016/S0140-6736\(10\)60576-4](https://doi.org/10.1016/S0140-6736(10)60576-4)

Chew EY, Ambrosius WT, Howard LT, Greven CM, Johnson S, Danis RP, Davis MD, Genuth S, Domanski M Rationale, design, and methods of the Action to Control Cardiovascular Risk in Diabetes Eye Study (ACCORD-EYE). Am J Cardiol 2007;99:103i-111i [17599420] [10.1016/j.amjcard.2007.03.028](https://doi.org/10.1016/j.amjcard.2007.03.028)

Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT Effects of intensive glucose lowering in type 2 diabetes. N Engl J Med 2008;358:2545-59 [18539917] [10.1056/NEJMoa0802743](https://doi.org/10.1056/NEJMoa0802743)

Cushman WC, Evans GW, Byington RP, Goff DC Jr, Grimm RH Jr, Cutler JA, Simons-Morton DG, Basile JN, Corson MA, Probstfield JL, Katz L, Peterson KA, Friedewald WT, Buse JB, Bigger JT, Gerstein HC, Ismail-Beigi F Effects of intensive blood-pressure control in type 2 diabetes mellitus. N Engl J Med 2010;362:1575-85 [20228401] [10.1056/NEJMoa1001286](https://doi.org/10.1056/NEJMoa1001286)

PAST-BP, 2015:

Wei, 2013:

Wei Y, Jin Z, Shen G, Zhao X, Yang W, Zhong Y, Wang J Effects of intensive antihypertensive treatment on Chinese hypertensive patients older than 70 years. J Clin Hypertens (Greenwich) 2013;15:420-7 [23730991] [10.1111/jch.12094](https://doi.org/10.1111/jch.12094)

SPS3, 2013:

Benavente OR, Coffey CS, Conwit R, Hart RG, McClure LA, Pearce LA, Pergola PE, Szychowski JM Blood-pressure targets in patients with recent lacunar stroke: the SPS3 randomised trial. Lancet 2013;382:507-15 [23726159] [10.1016/S0140-6736\(13\)60852-1](https://doi.org/10.1016/S0140-6736(13)60852-1)

HOMED-BP, 2012:

Asayama K, Ohkubo T, Metoki H, Obara T, Inoue R, Kikuya M, Thijs L, Staessen JA, Imai Y Cardiovascular outcomes in the first trial of antihypertensive therapy guided by self-measured home blood pressure. *Hypertens Res* 2012;35:1102-10 [22895063] [10.1038/hr.2012.125](#)

VANLISH, 2010:

Ogihara T, Saruta T, Rakugi H, Matsuoka H, Shimamoto K, Shimada K, Imai Y, Kikuchi K, Ito S, Eto T, Kimura G, Imaizumi T, Takishita S, Ueshima H Target blood pressure for treatment of isolated systolic hypertension in the elderly: valsartan in elderly isolated systolic hypertension study. *Hypertension* 2010;56:196-202 [20530299] [10.1161/HYPERTENSIONAHA.109.146035](#)

JATOS, 2008:

Principal results of the Japanese trial to assess optimal systolic blood pressure in elderly hypertensive patients (JATOS). *Hypertens Res* 2008;31:2115-27 [19139601] [10.1291/hypres.31.2115](#)

UKPDS-HDS, 1998:

Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13 [9732337]

SPRINT, 2015:

A Randomized Trial of Intensive versus Standard Blood-Pressure Control. *N Engl J Med* 2015 Nov 9; [26551272] [10.1056/NEJMoa1511939](#)

Cardio-Sis, 2009:

Verdecchia P, Staessen JA, Angeli F, de Simone G, Achilli A, Ganau A, Mureddu G, Pede S, Maggioni AP, Lucci D, Reboldi G Usual versus tight control of systolic blood pressure in non-diabetic patients with hypertension (Cardio-Sis): an open-label randomised trial. *Lancet* 2009;374:525-33 [19683638]

AASK, 2002:

Wright JT Jr, Bakris G, Greene T, Agodoa LY, Appel LJ, Charleston J, Cheek D, Douglas-Baltimore JG, Gassman J, Glassock R, Hebert L, Jamerson K, Lewis J, Phillips RA, Toto RD, Middleton JP, Rostand SG Effect of blood pressure lowering and antihypertensive drug class on progression of hypertensive kidney disease: results from the AASK trial. *JAMA* 2002;288:2421-31 [12435255]

ABCD target (H) , 2000:

Estacio RO, Jeffers BW, Gifford N, Schrier RW Effect of blood pressure control on diabetic microvascular complications in patients with hypertension and type 2 diabetes. *Diabetes Care* 2000;23 Suppl 2:B54-64 [10860192]

ABCD target (N) , 2002:

Schrier RW, Estacio RO, Esler A, Mehler P Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. *Kidney Int* 2002;61:1086-97 [11849464] [10.1046/j.1523-1755.2002.00213.x](#)

HOT, 1994:

Hansson L, Zanchetti A, Carruthers SG, Dahlf B, Elmfeldt D, Julius S, Mnard J, Rahm KH, Wedel H, Westerling S Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *Lancet* 1998;351:1755-62 [9635947]

Hansson L, Zanchetti A The Hypertension Optimal Treatment (HOT) Study: 24-month data on blood pressure and tolerability. *Blood Press* 1997;6:313-7 [9360003]

Hansson L, Zanchetti A The Hypertension Optimal Treatment (HOT) Study-patient characteristics: randomization, risk profiles, and early blood pressure results. *Blood Press* 1994;3:322-7 [7866597]

Zanchetti A, Hansson L, Dahlf B, Elmfeldt D, Kjeldsen S, Kolloch R, Larochelle P, McInnes GT, Mallion JM, Ruilope L, Wedel H Effects of individual risk factors on

the incidence of cardiovascular events in the treated hypertensive patients of the Hypertension Optimal Treatment Study. HOT Study Group. *J Hypertens* 2001;19:1149-59 [11403365]

REIN-2, 2005:

Ruggenti P, Perna A, Loriga G, Ganeva M, Ene-Iordache B, Turturro M, Lesti M, Perticucci E, Chakarski IN, Leonardis D, Garini G, Sessa A, Basile C, Alpa M, Scanziani R, Sorba G, Zoccali C, Remuzzi G Blood-pressure control for renoprotection in patients with non-diabetic chronic renal disease (REIN-2): multicentre, randomised controlled trial. *Lancet* 2005;365:939-46 [15766995] 10.1016/S0140-6736(05)71082-5

Tight blood pressure control and risk of macrovascular and microvascular complications in type 2 diabetes: UKPDS 38. UK Prospective Diabetes Study Group. *BMJ* 1998;317:703-13 [9732337]

MDRD, 1994:

Klahr S, Levey AS, Beck GJ, Caggiula AW, Hunsicker L, Kusek JW, Striker G The effects of dietary protein restriction and blood-pressure control on the progression of chronic renal disease. Modification of Diet in Renal Disease Study Group. *N Engl J Med* 1994;330:877-84 [8114857]

Lazarus JM, Bourgoignie JJ, Buckalew VM, Greene T, Levey AS, Milas NC, Paranandi L, Peterson JC, Porush JG, Rauch S, Soucie JM, Stollar C Achievement and safety of a low blood pressure goal in chronic renal disease. The Modification of Diet in Renal Disease Study Group. *Hypertension* 1997;29:641-50 [9040451]

Peterson JC, Adler S, Burkart JM, Greene T, Hebert LA, Hunsicker LG, King AJ, Klahr S, Massry SG, Seifter JL Blood pressure control, proteinuria, and the progression of renal disease. The Modification of Diet in Renal Disease Study. *Ann Intern Med* 1995;123:754-62 [7574193]

Sarnak MJ, Greene T, Wang X, Beck G, Kusek JW, Collins AJ, Levey AS The effect of a lower target blood pressure on the progression of kidney disease: long-term follow-up of the modification of diet in renal disease study. *Ann Intern Med* 2005;142:342-51 [15738453]

Toto, 1995:

Toto RD, Mitchell HC, Smith RD, Lee HC, McIntire D, Pettinger WA "Strict" blood pressure control and progression of renal disease in hypertensive nephrosclerosis. *Kidney Int* 1995;48:851-9 [7474675]

ACCORD blood pressure, 2008:

Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59 [18539917] 10.1056/NEJMoa0802743

Cushman WC, Grimm RH Jr, Cutler JA, Evans GW, Capes S, Corson MA, Sadler LS, Alderman MH, Peterson K, Bertoni A, Basile JN Rationale and design for the blood pressure intervention of the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Am J Cardiol* 2007;99:44i-55i [17599425] 10.1016/j.amjcard.2007.03.005

Effects of Intensive Blood-Pressure Control in Type 2 Diabetes Mellitus. *N Engl J Med* 2010 Mar 14;: [20228401] 10.1056/NEJMoa1001286

ESH-CHL-SHOT, :

ongoing trial NCT01563731

3 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
intensive dietary advice vs routine dietary advice			

continued...

Trial	Treatments	Patients	Trials design and methods
Wein , 1999 n=200 follow-up: 4.24 y	intensive dietary advice versus routine dietary advice	women with previous gestational diabetes and currently with impaired glucose tolerance (WHO 1985 criteria)	Parallel groups open USA
intensive vs usual			
ABCD normotensives , 1993 n=237/243 follow-up: 5.3 y	intensive (10 mm Hg below the baseline DBP) DBP control versus moderate (80 to 89 mm Hg) DBP control	normotensive type 2 diabetic patients	Parallel groups open
ACCORD (blood pressure) , 2010 [NCT00000620] n=2363/2371 follow-up: 4.7 y	intensive blood-pressure control, targeting a systolic pressure of less than 120 mm Hg versus standard blood-pressure control	high-risk patients with type 2 diabetes, high HbA1c concentrations (>7.5%), and cardiovascular disease (or >=2 cardiovascular risk factors)	Factorial plan open United States, Canada
HOT <=80 (diabetic subgroup) , 1998 n=499 follow-up: 3.8y	target diastolic blood pressure <=80 mmHg versus target diastolic blood pressure <=90 mmHg	patients aged 50-80 years with hypertension and diastolic blood pressure between 100 mm Hg and 115 mm Hg; diabetics subgroup	Parallel groups open 26 countries
intensive glycemic control vs conventional treatment			
ACCORD , 2008 [NCT00000620] n=5128/5123 follow-up: 3.5y (5y)	very intensive glycemic control through currently available means (targetinga glycosylated hemoglobin <6%) during a mean of 3.7 years versus standard glycemic control (targeting a glycosylated hemoglobin 7.0-7.9%)	patients with type 2 diabetes mellitus at high risk of death and stroke (pre-existing heart disease or two or more additional risk factors for heart disease)	Factorial plan open USA, Canada
ADDITION , 2010 [NCT00237549] n=1678/1379 follow-up: 5 year	intensive multifactorial treatment versus routine care	patients with newly diagnosed type 2 diabetes	Parallel groups open Denmark, United Kingdom, the Netherlands
ADVANCE , 2008 [NCT00145925] n=5571/5569 follow-up: median 5 y	intensive glucose-lowering treatments HbA1C <=6.5% using gliclazide(modified release) plus other drugs versus standard glucose-lowering treatments (targetglycated hemoglobin levels defined on the basisof local guidelines)	patients with type 2 diabetes	Parallel groups open 20 countries

continued...

Trial	Treatments	Patients	Trials design and methods
Kumamoto (primary prev) , 1995 n=28/27 follow-up: 8.0y	intensive glycemic control with multiple insulin injection treatment versus conventional insulin injection treatment (1-2 daily injections)	patients with non-insulin-dependent diabetes mellitus and with no retinopathy and urinary albumin excretions <30 mg/24 h	Parallel groups open Japan
Kumamoto (secondary prev) , 1995 n=27/28 follow-up: 8.0y	multiple insulin injection treatment versus conventional insulin injection treatment (1-2 daily injections)	patients with non-insulin-dependent diabetes mellitus and simple retinopathy	Parallel groups open Japan
Steno 2 , 2003 n=80/80 follow-up: 7.8 y	targeted, intensified, multifactorial intervention versus conventional treatment on modifiable risk factors for cardiovascular disease	patients with type 2 diabetes and microalbuminuria	Parallel groups open Denmark
UKPDS 33 , 1998 n=2729/1138 follow-up: 10.3 y	intensive policy with a sulphonylurea (chlorpropamide, glibenclamide, or glipizide) or with insulin; fasting plasma glucose <6.0 mmol/L versus conventional policy with diet	newly diagnosed patients with type 2 diabetes who after 3 months diet treatment had a mean of two fasting plasma glucose concentrations of 6.1150 mmol/L	Parallel groups open UK
VA CSDM , 1997 n=75/78 follow-up: 2.3y	intensive glycemic control (stepped plan from 1 evening injection of insulin, alone or with glipizide, to multiple daily injections, target to attain near-normal glycemia levels) versus standard treatment (1 insulin injection every morning)	non-insulin-dependent diabetes mellitus patients	Parallel groups open USA
VADT , 2008 [NCT00032487] n=892/899 follow-up: 5.6y	intensive glucose control versus standard glucose control	military veterans who had a suboptimal response to therapy for type 2 diabetes	Parallel groups open US
more intensive blood pressure lowering strategie vs less intensive blood pressure lowering strategie			
ABCD target (H) , 2000 n=237/233 follow-up: 5 year	intensive treatment with a diastolic blood pressure goal of 75 mmHg versus moderate treatment with a diastolic blood pressure goal of 80-89 mmHg	diabetes patients with DBP >=90 mmHg	Parallel groups open

continued...

Trial	Treatments	Patients	Trials design and methods
ABCD target (N) , 2002 n=237/243 follow-up:	intensive treatment (diastolic blood pressure decrease of 10 mmHg below baseline DBP) versus moderate treatment (diastolic blood pressure goal of 80-89 mmHg)	diabetes patients with diastolic blood pressure between 80 and 89mmHg	Parallel groups open

More details and results :

- anti hypertensive agents for diabetes type 2 in patients with hypertension at <http://www.trialresultscenter.org/go-Q83>
- intensive glycemc control for diabetes type 2 in Type 1 and 2 diabetes at <http://www.trialresultscenter.org/go-Q240>
- intensive glycemc control for diabetes type 2 in type 2 diabetes (NIDD) at <http://www.trialresultscenter.org/go-Q267>
- prevention for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q341>
- anti hypertensive agents for diabetes type 2 in patients with or without hypertension at <http://www.trialresultscenter.org/go-Q414>
- prevention for diabetes type 2 in people with impaired glucose tolerance at <http://www.trialresultscenter.org/go-Q416>
- intensive therapy for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q459>

References

Wein, 1999:

Wein P, Beischer N, Harris C, Permezel M A trial of simple versus intensified dietary modification for prevention of progression to diabetes mellitus in women with impaired glucose tolerance. Aust N Z J Obstet Gynaecol 1999;39:162-6 [[10755770](#)]

ABCD normotensives, 1993:

Savage S, Johnson Nagel N, Estacio RO, Feig PU, MacCarthy EP, Lukken NJ, Ziegler R, Schrier RW The ABCD (Appropriate Blood Pressure Control in Diabetes) trial. Rationale and design of a trial of hypertension control (moderate or intensive) in type II diabetes. Online J Curr Clin Trials 1993;Doc No 104:[6250 words; 128 paragraphs] [[8305994](#)]

Schrier RW, Estacio RO, Esler A, Mehler P Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. Kidney Int 2002;61:1086-97 [[11849464](#)] [10.1046/j.1523-1755.2002.00213.x](https://doi.org/10.1046/j.1523-1755.2002.00213.x)

ACCORD (blood pressure), 2010:

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. N Engl J Med 2010 Jun 29;: [[20587587](#)] [10.1056/NEJMoa1001288](https://doi.org/10.1056/NEJMoa1001288)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. Lancet 2010 Jun 29;: [[20594588](#)] [10.1016/S0140-6736\(10\)60576-4](https://doi.org/10.1016/S0140-6736(10)60576-4)

Chew EY, Ambrosius WT, Howard LT, Greven CM, Johnson S, Danis RP, Davis MD, Genuth S, Domanski M Rationale, design, and methods of the Action to Control Cardiovascular Risk in Diabetes Eye Study (ACCORD-EYE). *Am J Cardiol* 2007;99:103i-111i [17599420] [10.1016/j.amjcard.2007.03.028](https://doi.org/10.1016/j.amjcard.2007.03.028)

Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59 [18539917] [10.1056/NEJMoa0802743](https://doi.org/10.1056/NEJMoa0802743)

Cushman WC, Evans GW, Byington RP, Goff DC Jr, Grimm RH Jr, Cutler JA, Simons-Morton DG, Basile JN, Corson MA, Probstfield JL, Katz L, Peterson KA, Friedewald WT, Buse JB, Bigger JT, Gerstein HC, Ismail-Beigi F Effects of intensive blood-pressure control in type 2 diabetes mellitus. *N Engl J Med* 2010;362:1575-85 [20228401] [10.1056/NEJMoa1001286](https://doi.org/10.1056/NEJMoa1001286)

HOT <=80 (diabetic subgroup), 1998:

Hansson L, Zanchetti A, Carruthers SG, Dahlf B, Elmfeldt D, Julius S, Mnard J, Rahn KH, Wedel H, Westerling S Effects of intensive blood-pressure lowering and low-dose aspirin in patients with hypertension: principal results of the Hypertension Optimal Treatment (HOT) randomised trial. *Lancet* 1998;351:1755-62 [9635947]

Wesseling H [Lowering of diastolic blood pressure ; or = 90 mmHg should not be attempted, except in type 2 diabetics; the 'Hypertension optimal treatment' (HOT) trial] *Ned Tijdschr Geneeskd* 1999 Jun 5;143:1188-91 [10389531]

Hansson L The Hypertension Optimal Treatment study and the importance of lowering blood pressure. *J Hypertens Suppl* 1999 Feb;17:S9-13 [10340838]

Lopes AA, Andrade J, Noblat AC, Silveira MA Reduction in diastolic blood pressure and cardiovascular mortality in nondiabetic hypertensive patients. A reanalysis of the HOT study. *Arq Bras Cardiol* 2001;77:132-7 [11514824]

Zanchetti A; Hansson L Intensive blood pressure lowering: Contrast between benefits in diabetics and other groups of hypertensive patients and increased risk in smokers: Further analysis of the hypertension optimal treatment (HOT) study Abstract *JOURNAL OF HYPERTENSION* 2000;18(Supp: 2):S92-S92

Hansson L, Zanchetti A The Hypertension Optimal Treatment (HOT) Study—patient characteristics: randomization, risk profiles, and early blood pressure results. *Blood Press* 1994;3:322-7 [7866597]

Hansson L, Zanchetti A The Hypertension Optimal Treatment (HOT) Study: 24-month data on blood pressure and tolerability. *Blood Press* 1997;6:313-7 [9360003]

ACCORD, 2008:

Gerstein HC, Miller ME, Byington RP, Goff DC Jr, Bigger JT, Buse JB, Cushman WC, Genuth S, Ismail-Beigi F, Grimm RH Jr, Probstfield JL, Simons-Morton DG, Friedewald WT Effects of intensive glucose lowering in type 2 diabetes. *N Engl J Med* 2008;358:2545-59 [18539917]

Goff DC Jr, Gerstein HC, Ginsberg HN, Cushman WC, Margolis KL, Byington RP, Buse JB, Genuth S, Probstfield JL, Simons-Morton DG Prevention of cardiovascular disease in persons with type 2 diabetes mellitus: current knowledge and rationale for the Action to Control Cardiovascular Risk in Diabetes (ACCORD) trial. *Am J Cardiol* 2007;99:4i-20i [17599424]

Effects of Medical Therapies on Retinopathy Progression in Type 2 Diabetes. *N Engl J Med* 2010 Jun 29;: [20587587] [10.1056/NEJMoa1001288](https://doi.org/10.1056/NEJMoa1001288)

Ismail-Beigi F, Craven T, Banerji MA, Basile J, Calles J, Cohen RM, Cuddihy R, Cushman WC, Genuth S, Grimm RH Jr, Hamilton BP, Hoogwerf B, Karl D, Katz L, Krikorian A, O'Connor P, Pop-Busui R, Schubart U, Simmons D, Taylor H, Thomas A, Weiss D, Hramiak I Effect of intensive treatment of hyperglycaemia on microvascular outcomes in type 2 diabetes: an analysis of the ACCORD randomised trial. *Lancet* 2010 Jun 29;: [20594588] [10.1016/S0140-6736\(10\)60576-4](https://doi.org/10.1016/S0140-6736(10)60576-4)

Klein BE Reduction in Risk of Progression of Diabetic Retinopathy. *N Engl J Med* 2010 Jun 29;: [20587586] [10.1056/NEJMe1005667](https://doi.org/10.1056/NEJMe1005667)

Gerstein HC, Miller ME, Genuth S, Ismail-Beigi F, Buse JB, Goff DC Jr, Probstfield JL, Cushman WC, Ginsberg HN, Bigger JT, Grimm RH Jr, Byington RP, Rosenberg YD, Friedewald WT Long-term effects of intensive glucose lowering on cardiovascular outcomes. *N Engl J Med* 2011 Mar 3;364:818-28 [21366473] [10.1056/NEJMoa1006524](https://doi.org/10.1056/NEJMoa1006524)

ADDITION, 2010:

Sandbaek A, Griffin SJ, Rutten G, Davies M, Stolk R, Khunti K, Borch-Johnsen K, Wareham NJ, Lauritzen T Stepwise screening for diabetes identifies people with high but modifiable coronary heart disease risk. The ADDITION study. *Diabetologia* 2008;51:1127-34 [[18443762](#)] [10.1007/s00125-008-1013-0](#)

Griffin SJ, Borch-Johnsen K, Davies MJ, Khunti K, Rutten GE, Sandbk A, Sharp SJ, Simmons RK, van den Donk M, Wareham NJ, Lauritzen T Effect of early intensive multifactorial therapy on 5-year cardiovascular outcomes in individuals with type 2 diabetes detected by screening (ADDITION-Europe): a cluster-randomised trial. *Lancet* 2011 Jun 24;: [[21705063](#)] [10.1016/S0140-6736\(11\)60698-3](#)

ADVANCE, 2008:

Patel A, MacMahon S, Chalmers J, Neal B, Billot L, Woodward M, Marre M, Cooper M, Glasziou P, Grobbee D, Hamet P, Harrap S, Heller S, Liu L, Mancia G, Mogensen CE, Pan C, Poulter N, Rodgers A, Williams B, Bompont S, de Galan BE, Joshi R, Travert F Intensive blood glucose control and vascular outcomes in patients with type 2 diabetes. *N Engl J Med* 2008;358:2560-72 [[18539916](#)]

Kumamoto (primary prev), 1995:

Ohkubo Y, Kishikawa H, Araki E, Miyata T, Isami S, Motoyoshi S, Kojima Y, Furuyoshi N, Shichiri M Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995;28:103-17 [[7587918](#)]

Kumamoto (secondary prev), 1995:

Shichiri M, Kishikawa H, Ohkubo Y, Wake N Long-term results of the Kumamoto Study on optimal diabetes control in type 2 diabetic patients. *Diabetes Care* 2000;23 Suppl 2:B21-9 [[10860187](#)]

Ohkubo Y, Kishikawa H, Araki E, Miyata T, Isami S, Motoyoshi S, Kojima Y, Furuyoshi N, Shichiri M Intensive insulin therapy prevents the progression of diabetic microvascular complications in Japanese patients with non-insulin-dependent diabetes mellitus: a randomized prospective 6-year study. *Diabetes Res Clin Pract* 1995;28:103-17 [[7587918](#)]

Steno 2, 2003:

Gaede P, Vedel P, Larsen N, Jensen GV, Parving HH, Pedersen O Multifactorial intervention and cardiovascular disease in patients with type 2 diabetes. *N Engl J Med* 2003;348:383-93 [[12556541](#)]

UKPDS 33, 1998:

Intensive blood-glucose control with sulphonylureas or insulin compared with conventional treatment and risk of complications in patients with type 2 diabetes (UKPDS 33). UK Prospective Diabetes Study (UKPDS) Group. *Lancet* 1998;352:837-53 [[9742976](#)]

VA CSDM, 1997:

Abraira C, Colwell J, Nuttall F, Sawin CT, Henderson W, Comstock JP, Emanuele NV, Levin SR, Pacold I, Lee HS Cardiovascular events and correlates in the Veterans Affairs Diabetes Feasibility Trial. Veterans Affairs Cooperative Study on Glycemic Control and Complications in Type II Diabetes. *Arch Intern Med* 1997;157:181-8 [[9009975](#)]

VADT, 2008:

Duckworth W, Abraira C, Moritz T, Reda D, Emanuele N, Reaven PD, Zieve FJ, Marks J, Davis SN, Hayward R, Warren SR, Goldman S, McCarren M, Vitek ME, Henderson WG, Huang GD Glucose Control and Vascular Complications in Veterans with Type 2 Diabetes. *N Engl J Med* 2008 Dec 17;: [[19092145](#)]

ABCD target (H) , 2000:

Estacio RO, Jeffers BW, Gifford N, Schrier RW Effect of blood pressure control on diabetic microvascular complications in patients with hypertension and type 2 diabetes. *Diabetes Care* 2000;23 Suppl 2:B54-64 [[10860192](#)]

ABCD target (N) , 2002:

Schrier RW, Estacio RO, Esler A, Mehler P Effects of aggressive blood pressure control in normotensive type 2 diabetic patients on albuminuria, retinopathy and strokes. *Kidney Int* 2002;61:1086-97 [[11849464](#)] [10.1046/j.1523-1755.2002.00213.x](#)

Entry terms: intensive blood pressure lowering strategies, intensive treatment, tighter control of blood pressure, low target blood pressure, strict blood pressure control, intensified blood-pressure control