

Clinical trials of kissing balloon

TrialResults-center www.trialresultscenter.org

1 coronary artery disease

Trial	Treatments	Patients	Trials design and methods
kissing balloon vs no kissing balloon			
Nordic-Baltic Bifurcation Study III , 2009 [NCT00914199] n=238/239 follow-up: 6 mo	Kissing balloon dilatation post-stenting of the main artery (one-stent technique) versus no kissing balloon dilatation	patients with bifurcation lesions	Parallel groups open Denmark, Finland, Latvia, Sweden, Norway

More details and results :

- Drug eluting stent for coronary artery disease in bifurcation lesion at <http://www.trialresultscenter.org/go-Q214>

References

Nordic-Baltic Bifurcation Study III, 2009:

Niemel M, Kervinen K, Erglis A, Holm NR, Maeng M, Christiansen EH, Kumsars I, Jegere S, Dombrovskis A, Gunnes P, Stavnes S, Steigen TK, Trovik T, Eskola M, Vikman S, Romppanen H, Mkikallio T, Hansen KN, Thayssen P, Aberg L, Jensen LO, Hervold A, Airaks Randomized Comparison of Final Kissing Balloon Dilatation Versus No Final Kissing Balloon Dilatation in Patients With Coronary Bifurcation Lesions Treated With Main Vessel Stenting: The Nordic-Baltic Bifurcation Study III. *Circulation* 2011;123:79-86 [[21173348](https://pubmed.ncbi.nlm.nih.gov/21173348/)] [10.1161/CIRCULATIONAHA.110.966879](https://doi.org/10.1161/CIRCULATIONAHA.110.966879)