

Clinical trials of naftidrofuryl IV

TrialResults-center www.trialresultscenter.org

1 peripheral vascular diseases

Trial	Treatments	Patients	Trials design and methods
naftidrofuryl IV vs control			
Meehan , 1982 n=24/16 follow-up: 7 days	Naftidrofuryl (Nafronyl): par voie IV (200 mg*2/ jour pendant 2 heures dans 500 ml de srum sal isotonique) et par voie orale (200mg*3/ jour)+ repos au lit + rchauffement pendant 7 jours versus Repos au lit + rchauffement	AOMI stade III	Parallel groups Simple aveugle
naftidrofuryl IV vs placebo			
Testart , 1994 n=20/17 follow-up: 8 jours	2 daily infusions of 400 mg of naftidrofuryl for 8 days versus placebo	AOMI stade III ou IV	Parallel groups Double aveugle
D'Hooge D , 2001 n=116/119 follow-up: 6 mois	- effet de naftidrofuryl sur la qualitt de vie de patients souffrant de claudication intermittente stable : 3x200mg pdt 6 mois versus - placebo pdt 6 mois	- 235 patients - ge moyen 66,5 ans - 66,4% d'hommes	Parallel groups Double aveugle
naftidrofuryl IV vs prostacyclin			
Negus , 1987 n=15/14 follow-up: 24 h	Naftidrofuryl 0.02 mg/kg/min, intra-arterial, 3 days versus intra-arterial prostacyclin 8 ng/kg/min, 3 days	Stade de la maladie: III: 55.2% ; IV: 44.8% Dure de la douleur (semaines): groupe contrle: 30 +/-54.2 groupe ttt: 19.7 +/- 44.6	Parallel groups Double aveugle

More details and results :

- vasodilators therapy for peripheral vascular diseases in all type of patients at <http://www.trialresultscenter.org/go-Q49>

References

Meehan, 1982:

The usefulness of naftidrofuryl in severe peripheral ischaemia—a symptomatic assessment using linear analogue scales. Meehan SE, Preece PE, Walker WF *Angiology* 1982 Oct;33:625-34 [[6751162](#)]

Testart, 1994:

[Evaluation of the analgesic effect of naftidrofuryl in permanent ischemia in patients with arterial diseases. Double-blind versus placebo study] Testart J, Guidicelli H, Glanddier G, Mosnier M *Ann Cardiol Angeiol (Paris)* 1994 Nov;43:542-7 [[7864561](#)]

D'Hooge D, 2001:

D. D'Hooge; P.Lehert, D.L.Clement Naftidrofuryl in quality of life (NIQOL) A Belgian study. *Int Angiol* 2001;20:288-94

Negus, 1987:

Intra-arterial prostacyclin compared to Praxilene in the management of severe lower limb ischaemia: a double blind trial. Negus D, Irving JD, Friedgood A *J Cardiovasc Surg (Torino)* 1987 Mar-Apr;28:196-9 [[2435738](#)]

Entry terms: naftidrofuryl