

# Clinical trials of niacin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 post myocardial infarction

Trial	Treatments	Patients	Trials design and methods
<b>clofibrate+niacin vs placebo</b>			
<a href="#">Carlson (Stockholm) , 1977</a> n=279/276 follow-up: 5 years	clofibrate, 1 g twice daily, and nicotinic acid 1 g three times daily versus control	survivors of a myocardial infarction below 70 years of age	Parallel groups open Sweden

More details and results :

- cholesterol lowering intervention for post myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q45>

## References

### Carlson (Stockholm), 1977:

Carlson LA, Danielson M, Ekberg I, Klintemar B, Rosenhamer G, Reduction of myocardial reinfarction by the combined treatment with clofibrate and nicotinic acid. *Atherosclerosis* 1977; 28:81-6 [[911371](#)]

Carlson LA, Rosenhamer G, Reduction of mortality in the Stockholm Ischaemic Heart Disease Secondary Prevention Study by combined treatment with clofibrate and nicotinic acid. *Acta Med Scand* 1988; 223:405-18 [[3287837](#)]

## 2 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
<b>niacin vs control</b>			
<a href="#">VA drugs , 1968</a> n=77/143 follow-up: 3.2 years	-	-	Parallel groups double blind
<b>niacin+colestipol vs control</b>			
<a href="#">UCSF SCOR , 1990</a> n=72 follow-up: 26 months	Niacin 07.5 g colestipol 1520 g versus Conventional therapy	patients with heterozygous familial hypercholesterolemia	

continued...

Trial	Treatments	Patients	Trials design and methods
<b>clofibrate+niacin vs placebo</b>			
<b>Carlson (Stockholm) , 1977</b> n=279/276 follow-up: 5 years	clofibrate, 1 g twice daily, and nicotinic acid 1 g three times daily versus control	survivors of a myocardial infarction below 70 years of age	Parallel groups open Sweden
<b>colestipol-niacin vs placebo</b>			
<b>CLAS , 1987</b> n=94/94 follow-up: 2 ans	Colestipol + Niacin 30 g / j 3-12 g / j (titr sur chaque patient sur la base de la baisse de cholestrol sanguin) versus placebo: methyl cellulose	Patients coronariens avec antcedent de revascularisation chirurgicale coronarienne.	Parallel groups Non dterminable
<b>CLAS , 1987</b> n=NA follow-up: 65279;2 years	colestipol + niacin versus placebo	nonsmoking men aged 40 to 59 years with previous coronary bypass surgery	Parallel groups double blind
<b>niacin vs placebo</b>			
<b>CDP niacin , 1975</b> n=1119/2789 follow-up: 6.2 years	niacin 3 mg/d versus placebo	Hommes, de 30 64 ans	Parallel groups double blind
<b>niacin+colestipol vs placebo</b>			
<b>FATS , 1990</b> n=48/54 follow-up: 2.5 years	niacin (1 g four times a day) and colestipol (10 g three times a day) versus placebo (or colestipol if the low-density lipoprotein [LDL] cholesterol level was elevated)	men no more than 62 years of age with apolipoprotein B levels greater than or equal to 125 mg per deciliter, documented coronary artery disease, and a family history of vascular disease	Parallel groups double-blind
<b>niacin vs placebo (on top statin)</b>			
<b>AIM-HIGH , 2011</b> [NCT00120289] n=1718/1691 follow-up: 32 months	high-dose, extended-release niacin in gradually increasing doses up to 2000 mg daily (+ simvastatin) versus placebo	patients with a history of cardiovascular disease, high triglycerides, and low levels of HDL cholesterol	Parallel groups double blind US, Canada
<b>HPS 2-Thrive</b> [NCT00461630] n=12838/12835 follow-up: 3.9y (median)	2 g of extended-release niacin and 40 mg of laropiprant versus placebo	patients with vascular disease	Parallel groups double blind UK, Scandinavia, China

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Oxford Niaspan Study , 2009</b> [NCT00232531] n=35/36 follow-up: 1 year	niacin 2g daily (added to statin therapy) versus placebo (statins alone)	patients with low HDL-C (<40 mg/dl) and either a type 2 diabetes with coronary heart disease or a carotid/peripheral atherosclerosis	Parallel groups double blind USA
<b>ARBITER 2 , 2009</b> n=87/80 follow-up: 1 y	long-acting niacin target dose of 1 g/day (added to statin therapy) versus placebo	patients with known coronary artery disease and well controlled on statin therapy	Parallel groups double blind USA
<b>HATS , 2001</b> n=73/73 follow-up: 3 y	simvastatin plus niacin versus placebo	patients with coronary disease, low HDL cholesterol levels and normal LDL cholesterol levels	Factorial plan double blind USA, Canada
<b>niacin vs ezetimibe</b>			
<b>ARBITER 6-HALTS (niacin vs ezetimibe) , 2009</b> [NCT00397657] n=97/111 follow-up: 14 months	extended-release niacin 1 g/d, titrated to max tolerable dose up to 2 g/d (HDL-focused strategy) versus ezetimibe 10 mg/d (LDL-focused strategy)	patients with known coronary or vascular disease or coronary risk equivalents	Parallel groups open US
<b>niacin+ezetimibe vs simvastatin+ezetimibe</b>			
<b>Guyton , 2008</b> n=NA follow-up: 24 weeks	Niacin 2 g ezetimibe 10 mg simvastatin 20 mg versus Ezetimibe 10 mg simvastatin 20 mg	patients with type IIa or IIb hyperlipidemia	Parallel groups double-blind

More details and results :

- cholesterol lowering intervention for cardiovascular prevention in patients with prior MI or with CHD at <http://www.trialresultscenter.org/go-Q12>
- cholesterol lowering intervention for cardiovascular prevention in all chronic situations at <http://www.trialresultscenter.org/go-Q154>
- niacin for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q326>
- HDL increasing drugs for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q503>

## References

VA drugs, 1968:

Schoch HK. The US Veterans Administration Cardiology drug lipid study: an interim report *Adv Exp Med Biol.* 1968;4:405-420

#### **UCSF SCOR, 1990:**

Kane JP, Malloy MJ, Ports TA, Phillips NR, Diehl JC, Havel RJ Regression of coronary atherosclerosis during treatment of familial hypercholesterolemia with combined drug regimens. *JAMA* 1990;264:3007-12 [[2243428](#)]

#### **Carlson (Stockholm), 1977:**

Carlson LA, Danielson M, Ekberg I, Klintemar B, Rosenhamer G, Reduction of myocardial reinfarction by the combined treatment with clofibrate and nicotinic acid. *Atherosclerosis* 1977; 28:81-6 [[911371](#)]

Carlson LA, Rosenhamer G, Reduction of mortality in the Stockholm Ischaemic Heart Disease Secondary Prevention Study by combined treatment with clofibrate and nicotinic acid. *Acta Med Scand* 1988; 223:405-18 [[3287837](#)]

#### **CLAS, 1987:**

Blankenhorn DH, Brooks SH. Angiographic trials of lipid-lowering therapy. *Arteriosclerosis* 1981; 1: 242-249.

The Cholesterol Lowering Atherosclerosis Study (CLAS): design, methods, and baseline results. Blankenhorn DH, Johnson RL, Nessim SA, Azen SP, Sanmarco ME, Selzer RH *Control Clin Trials* 1987 Dec;8:356-87 [[3327654](#)]

Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. Blankenhorn DH, Nessim SA, Johnson RL, Sanmarco ME, Azen SP, Cashin-Hemphill L *JAMA* 1987 Jun 19;257:3233-40 [[3295315](#)]

Comparison of computer- and human-derived coronary angiographic end-point measures for controlled therapy trials. Mack WJ, Selzer RH, Pogoda JM, Lee PL, Shircore AM, Azen SP, Blankenhorn DH *Arterioscler Thromb* 1992 Mar;12:348-56 [[1547194](#)]

#### **CLAS, 1987:**

Blankenhorn DH, Johnson RL, Nessim SA, Azen SP, Sanmarco ME, Selzer RH The Cholesterol Lowering Atherosclerosis Study (CLAS): design, methods, and baseline results. *Control Clin Trials* 1987;8:356-87 [[3327654](#)]

Cashin-Hemphill L, Mack WJ, Pogoda JM, Sanmarco ME, Azen SP, Blankenhorn DH Beneficial effects of colestipol-niacin on coronary atherosclerosis. A 4-year follow-up. *JAMA* 1990;264:3013-7 [[2243429](#)]

Blankenhorn DH, Nessim SA, Johnson RL, Sanmarco ME, Azen SP, Cashin-Hemphill L Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. *JAMA* 1987;257:3233-40 [[3295315](#)]

Blankenhorn DH, Azen SP, Crawford DW, Nessim SA, Sanmarco ME, Selzer RH, Shircore AM, Wickham EC Effects of colestipol-niacin therapy on human femoral atherosclerosis. *Circulation* 1991;83:438-47 [[1991366](#)]

#### **CDP niacin, 1975:**

, Clofibrate and niacin in coronary heart disease. *JAMA* 1975; 231:360-81 [[1088963](#)]

#### **FATS, 1990:**

Brown G, Albers JJ, Fisher LD, Schaefer SM, Lin JT, Kaplan C, Zhao XQ, Bisson BD, Fitzpatrick VF, Dodge HT Regression of coronary artery disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein B. *N Engl J Med* 1990;323:1289-98 [[2215615](#)]

#### **AIM-HIGH, 2011:**

Boden WE, Probstfield JL, Anderson T, Chaitman BR, Desvignes-Nickens P, Koprowicz K, McBride R, Teo K, Weintraub W Niacin in patients with low HDL cholesterol levels receiving intensive statin therapy. *N Engl J Med* 2011;365:2255-67 [[22085343](#)] [10.1056/NEJMoa1107579](#)

#### **HPS 2-Thrive, :**

Landray MJ, Haynes R, Hopewell JC, Parish S, Aung T, Tomson J, Wallendszus K, Craig M, Jiang L, Collins R, Armitage J Effects of extended-release niacin with laropiprant in high-risk patients. N Engl J Med 2014 Jul 17;371:203-12 [25014686] 10.1056/NEJMoa1300955

#### **Oxford Niaspan Study, 2009:**

Lee JM, Robson MD, Yu LM, Shirodaria CC, Cunnington C, Kylintireas I, Digby JE, Bannister T, Handa A, Wiesmann F, Durrington PN, Channon KM, Neubauer S, Choudhury RP Effects of high-dose modified-release nicotinic Acid on atherosclerosis and vascular function a randomized, placebo-controlled, magnetic resonance imaging study. J Am Coll Cardiol 2009;54:1787-94 [19874992]

#### **ARBITER 2, 2009:**

Taylor AJ, Sullenberger LE, Lee HJ, Lee JK, Grace KA Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol (ARBITER) 2: a double-blind, placebo-controlled study of extended-release niacin on atherosclerosis progression in secondary prevention patients treated with statins. Circulation 2004;110:3512-7 [15537681]

#### **HATS, 2001:**

Brown BG, Zhao XQ, Chait A, Fisher LD, Cheung MC, Morse JS, Dowdy AA, Marino EK, Bolson EL, Alaupovic P, Frohlich J, Albers JJ Simvastatin and niacin, antioxidant vitamins, or the combination for the prevention of coronary disease. N Engl J Med 2001 Nov 29;345:1583-92 [11757504]

#### **ARBITER 6-HALTS (niacin vs ezetimibe), 2009:**

Taylor AJ, Villines TC, Stanek EJ, Devine PJ, Griffen L, Miller M, Weissman NJ, Turco M Extended-release niacin or ezetimibe and carotid intima-media thickness. N Engl J Med 2009 Nov 26;361:2113-22 [19915217]

Villines TC, Stanek EJ, Devine PJ, Turco M, Miller M, Weissman NJ, Griffen L, Taylor AJ The ARBITER 6-HALTS Trial (Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol 6-HDL and LDL Treatment Strategies in Atherosclerosis) Final Results and the Impact of Medication Adherence, Dose, and Treatment Duration. J Am Coll Cardiol 2010 Apr 8;: [20399059] 10.1016/j.jacc.2010.03.017

#### **Guyton, 2008:**

Guyton JR, Brown BG, Fazio S, Polis A, Tomassini JE, Tershakovec AM Lipid-altering efficacy and safety of ezetimibe/simvastatin coadministered with extended-release niacin in patients with type IIa or type IIb hyperlipidemia. J Am Coll Cardiol 2008;51:1564-72 [18420099] 10.1016/j.jacc.2008.03.003

### 3 peripheral vascular diseases

Trial	Treatments	Patients	Trials design and methods
<b>colestipol-niacin vs placebo</b>			
CLAS , 1987 n=94/94 follow-up: 2 ans	Colestipol + Niacin 30 g / j 3-12 g / j (titr sur chaque patient sur la base de la baisse de cholestrol sanguin) versus placebo: methyl cellulose	Patients coronariens avec antcdent de revascularisation chirurgicale coronarienne.	Parallel groups Non dterminable

More details and results :

- cholesterol lowering intervention for peripheral vascular diseases in all type of patients at <http://www.trialresultscenter.org/go-Q52>

## References

### CLAS, 1987:

Blankenhorn DH, Brooks SH. Angiographic trials of lipid-lowering therapy. *Arteriosclerosis* 1981; 1: 242-249.

The Cholesterol Lowering Atherosclerosis Study (CLAS): design, methods, and baseline results. Blankenhorn DH, Johnson RL, Nessim SA, Azen SP, Sanmarco ME, Selzer RH *Control Clin Trials* 1987 Dec;8:356-87 [[3327654](#)]

Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. Blankenhorn DH, Nessim SA, Johnson RL, Sanmarco ME, Azen SP, Cashin-Hemphill L *JAMA* 1987 Jun 19;257:3233-40 [[3295315](#)]

Comparison of computer- and human-derived coronary angiographic end-point measures for controlled therapy trials. Mack WJ, Selzer RH, Pogoda JM, Lee PL, Shircore AM, Azen SP, Blankenhorn DH *Arterioscler Thromb* 1992 Mar;12:348-56 [[1547194](#)]

Entry terms: Niacin, Nicotinic Acid, 3-Pyridinecarboxylic Acid, 3 Pyridinecarboxylic Acid, Induracin, Nicamin, Nico-400, Nico 400, Nico400, Nicobid, Nicocap, Nicolar, Nicotinate, Wampocap, Enduracin, Lithium Nicotinate