

# Clinical trials of once daily dalteparin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 venous thrombosis

Trial	Treatments	Patients	Trials design and methods
<b>once daily dalteparin vs twice daily dalteparin</b>			
<b>Holmstrm , 1992</b> n=50/51 follow-up:	once daily dalteparin 200 U (anti-FXa)/kg for at least 5 days versus twice daily dalteparin 100 U (anti-FXa)/kg for at least 5 days	Patients with a first occurrence of DVT in the lower limb, confirmed with phlebography	Parallel groups open Sweden
<b>Partsch , 1996</b> n=76/64 follow-up:	Fragmin administered 200 IU/kg once daily for at least 7 days versus Fragmin 100 IU/kg twice daily for at least 7 days	patients presented with DVT extending into the iliofemoral segment diagnosed by duplex ultrasonography	Parallel groups NA Austria

More details and results :

- antithrombotics for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q101>
- LMWH for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q203>
- heparin (UFH or LMWH) for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q204>

## References

### Holmstrm, 1992:

Holmstrm M, Berglund MC, Granquist S, Bratt G, Trnebohm E, Lockner D Fragmin once or twice daily subcutaneously in the treatment of deep venous thrombosis of the leg. Thromb Res 1992;67:49-55 [[1332213](#)]

### Partsch, 1996:

Partsch H, Kechavarz B, Mostbeck A, Khn H, Lipp C Frequency of pulmonary embolism in patients who have iliofemoral deep vein thrombosis and are treated with once- or twice-daily low-molecular-weight heparin. J Vasc Surg 1996;24:774-82 [[8918323](#)]

Entry terms: dalteparin