

# Clinical trials of once daily logiparin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 venous thrombosis

Trial	Treatments	Patients	Trials design and methods
<b>once daily logiparin vs twice daily logiparin</b>			
<a href="#">Siegbahn , 1989</a> n=10/10 follow-up:	Once daily logiparin 150 XaI U/kgp, imag versus twice daily logiparin 75 XaI U/kg	patients with a venographically confirmed episode of DVT	Parallel groups single blind Sweden and Denmark

More details and results :

- antithrombotics for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q101>
- LMWH for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q203>
- heparin (UFH or LMWH) for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q204>

## References

### Siegbahn, 1989:

Siegbahn A, Y-Hassan S, Boberg J, Bylund H, Neerstrand HS, Ostergaard P, Hedner U Subcutaneous treatment of deep venous thrombosis with low molecular weight heparin. A dose finding study with LMWH-Novo. Thromb Res 1989;55:767-78 [[2551070](#)]

Entry terms: logiparin, Logiparin, LHN-1,