

Clinical trials of once daily logiparin

TrialResults-center www.trialresultscenter.org

1 venous thrombosis

| Trial | Treatments | Patients | Trials design and methods |
|--|---|---|---|
| once daily logiparin vs twice daily logiparin | | | |
| Siegbahn , 1989 n=10/10 follow-up: | Once daily logiparin 150 XaI U/kgp, imag versus twice daily logiparin 75 XaI U/kg | patients with a venographically confirmed episode of DVT | Parallel groups single blind Sweden and Denmark |

More details and results :

- antithrombotics for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q101>
- LMWH for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q203>
- heparin (UFH or LMWH) for venous thrombosis in all type of patients at <http://www.trialresultscenter.org/go-Q204>

References

Siegbahn, 1989:

Siegbahn A, Y-Hassan S, Boberg J, Bylund H, Neerstrand HS, Ostergaard P, Hedner U Subcutaneous treatment of deep venous thrombosis with low molecular weight heparin. A dose finding study with LMWH-Novo. Thromb Res 1989;55:767-78 [[2551070](#)]

Entry terms: logiparin, Logiparin, LHN-1,