

Clinical trials of out of hospital Dalteparin

TrialResults-center www.trialresultscenter.org

1 thrombosis prevention

Trial	Treatments	Patients	Trials design and methods
out of hospital Dalteparin vs standard prophylaxis			
Dahl , 1997 n=134/131	65279;in hospital thromboprophylaxis followed by out of hospital Dalteparin 5000 IU once a day for a total duration of 35 days versus Dalteparin 5000 IU once a day for 7 days (dextran day 0 and day 1)	THR	
Lassen , 1998 n=140/141	65279;in hospital thromboprophylaxis followed by out of hospital Dalteparin 5000 IU once a day for a total duration of 35 days versus Dalteparin 5000 IU once a day for 7 days	THR	
Hull , 2000 n=389/180	65279;in hospital thromboprophylaxis followed by out of hospital Dalteparin 5000 IU once a day for a total duration of 35 days versus Dalteparin 5000 IU once a day or warfarin for 6 days	THR	

More details and results :

- antithrombotics for thrombosis prevention in orthopedic surgery at <http://www.trialresultscenter.org/go-Q37>

References

Dahl, 1997:

Dahl OE, Andreassen G, Aspelin T, Mller C, Mathiesen P, Nyhus S, Abdelnoor M, Solhaug JH, Arnesen H Prolonged thromboprophylaxis following hip replacement surgery—results of a double-blind, prospective, randomised, placebo-controlled study with dalteparin (Fragmin) Thromb Haemost 1997;77:26-31 [9031444]

Lassen, 1998:

Lassen MR, Borris LC, Anderson BS, Jensen HP, Skej Bro HP, Andersen G, Petersen AO, Siem P, Hrlyck E, Jensen BV, Thomsen PB, Hansen BR, Erin-Madsen J, Mller JC, Rotwitt L, Christensen F, Nielsen JB, Jrgensen PS, Paaske B, Trholm C, Hvidt P, Jensen NK Efficacy and safety of prolonged thromboprophylaxis with a low molecular weight heparin (dalteparin) after total hip arthroplasty—the Danish Prolonged Prophylaxis (DaPP) Study. *Thromb Res* 1998;89:281-7 [[9669750](#)]

Hull, 2000:

Hull RD, Pineo GF, Francis C, Bergqvist D, Fellenius C, Soderberg K, Holmqvist A, Mant M, Dear R, Baylis B, Mah A, Brant R Low-molecular-weight heparin prophylaxis using dalteparin extended out-of-hospital vs in-hospital warfarin/out-of-hospital placebo in hip arthroplasty patients: a double-blind, randomized comparison. North American Fragmin Trial Investigators. *Arch Intern Med* 2000;160:2208-15 [[10904465](#)]

Entry terms: dalteparin