

Clinical trials of polymer free sirolimus stent

TrialResults-center www.trialresultscenter.org

1 coronary artery disease

Trial	Treatments	Patients	Trials design and methods
polymer free sirolimus stent vs sirolimus eluting stent			
ISAR TEST 3 (PF) , 2009 n=201/202 follow-up: 12 months	polymer free 2% rapamycin (479 mg rapamycin/cm2) stent versus permanent-polymer rapamycin-eluting stent (Cypher) (140 mg rapamycin/cm2)	Patients with de novo coronary lesions in native vessels	open Germany

More details and results :

- myocardial revascularization for coronary artery disease in all type of patient at <http://www.trialresultscenter.org/go-Q26>
- Drug eluting stent for coronary artery disease in all type of patients at <http://www.trialresultscenter.org/go-Q206>

References

ISAR TEST 3 (PF), 2009:

Mehilli J, Byrne RA, Wiecek A, Iijima R, Schulz S, Bruskin O, Pache J, Wessely R, Schmig A, Kastrati A Eur Heart J 2008 Aug;29:1975-82 [[18550554](https://doi.org/10.1093/eurheartj/ehn253)]
[10.1093/eurheartj/ehn253](https://doi.org/10.1093/eurheartj/ehn253)