

# Clinical trials of rivaroxaban (long duration)

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 DVT prophylaxis

Trial	Treatments	Patients	Trials design and methods
<b>rivaroxaban (long duration) vs enoxaparin (short duration)</b>			
<b>RECORD 2 , 2008</b> [NCT00332020] n=1252/1257 follow-up: 30-42 days	extended thromboprophylaxis with rivaroxaban 10mg once daily for 31-39 days versus enoxaparin 40mg subcutaneous once daily for 10-14 days	patients undergoing elective total hip replacement	Parallel groups double blind 21 countries worldwide

More details and results :

- antithrombotics for DVT prophylaxis in orthopedic surgery at <http://www.trialresultscenter.org/go-Q37>
- antithrombotics for DVT prophylaxis in elective hip replacement at <http://www.trialresultscenter.org/go-Q39>
- anticoagulant for DVT prophylaxis in orthopedic surgery at <http://www.trialresultscenter.org/go-Q184>
- direct factor Xa inhibitors for DVT prophylaxis in all type of patients at <http://www.trialresultscenter.org/go-Q371>
- direct oral anticoagulant (DAO) for DVT prophylaxis in all type of patients at <http://www.trialresultscenter.org/go-Q393>
- direct oral anticoagulant (DAO) for DVT prophylaxis in elective hip replacement at <http://www.trialresultscenter.org/go-Q395>
- direct oral anticoagulant (DAO) for DVT prophylaxis in orthopaedic surgery at <http://www.trialresultscenter.org/go-Q475>

## References

### RECORD 2, 2008:

Kakkar AK, Brenner B, Dahl OE, Eriksson BI, Mouret P, Muntz J, Soglian AG, Pap AF, Misselwitz F, Haas S Extended duration rivaroxaban versus short-term enoxaparin for the prevention of venous thromboembolism after total hip arthroplasty: a double-blind, randomised controlled trial. Lancet 2008 Jun 24; [18582928]

Entry terms: rivaroxaban, Xarelto, BAY 59-7939,