Clinical trials of rosiglitazone and metformin

TrialResults-center www.trialresultscenter.org

1 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
rosiglitazone and metformin vs placebo			
CANOE , 2010 [NCT00116932] n=103/104 follow-up: 3.9y (median)	rosiglitazone (2 mg) and metformin (500 mg) twice-daily versus placebo	patients with impaired glucose tolerance	Parallel groups double-blind

More details and results :

- prevention for diabetes type 2 in all type of patients at http://www.trialresultscenter.org/go-Q341

- prevention for diabetes type 2 in people with impaired glucose tolerance at http://www.trialresultscenter.org/go-Q416

References

CANOE, 2010:

Zinman B, Harris SB, Neuman J, Gerstein HC, Retnakaran RR, Raboud J, Qi Y, Hanley AJ Low-dose combination therapy with rosiglitazone and metformin to prevent type 2 diabetes mellitus (CANOE trial): a double-blind randomised controlled study. Lancet 2010 Jul 10;376:103-11 [20605202] 10.1016/S0140-6736(10)60746-5

2 impaired fasting glucose

Trial	Treatments	Patients	Trials design and methods
rosiglitazone and metformin vs placebo			
CANOE , 2010 [NCT00116932] n=103/104 follow-up: 3.9y (median)	rosiglitazone (2 mg) and metformin (500 mg) twice-daily versus placebo	patients with impaired glucose tolerance	Parallel groups double-blind

More details and results :

- prevention for impaired fasting glucose in all type of patients at http://www.trialresultscenter.org/go-Q342

References

CANOE, 2010:

Zinman B, Harris SB, Neuman J, Gerstein HC, Retnakaran RR, Raboud J, Qi Y, Hanley AJ Low-dose combination therapy with rosiglitazone and metformin to prevent type 2 diabetes mellitus (CANOE trial): a double-blind randomised controlled study. Lancet 2010 Jul 10;376:103-11 [20605202] 10.1016/S0140-6736(10)60746-5

Entry terms: metformin, Metformin, Dimethylguanylguanidine, Dimethylbiguanidine, Glucophage,