

# Clinical trials of spironolactone

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 heart failure

Trial	Treatments	Patients	Trials design and methods
<b>spironolactone vs control</b>			
<a href="#">Cicoira , 2002</a> n=54/52 follow-up: 12 months	spironolactone 12.5 to 50 mg/day versus control	patients with chronic heart failure	Parallel groups open
<a href="#">Cicoira , 2004</a> n=47/46 follow-up: 12 months	spironolactone versus control	chronic heart failure patients	open
<a href="#">Ramires , 2000</a> n=19/16 follow-up: 20 weeks	spironolactone versus standard medical treatment	patients with systolic dysfunction and NYHA class III CHF secondary to dilated or ischemic cardiomyopathy	Parallel groups open
<b>spironolactone vs placebo</b>			
<a href="#">Agostoni , 2005</a> n=14/15 follow-up: 6 months	spironolactone 25mg/d versus placebo	stable chronic heart failure patients with reduced influences lung diffusion (DLCO)	Parallel groups open Italy
<a href="#">Barr , 1995</a> n=28/14 follow-up: 8 weeks	spironolactone 50 to 100 mg/day, titrated to blood pressure and plasma potassium (added to an angiotensin-converting enzyme inhibitor) versus placebo	patients with New York Heart Association II to III congestive heart failure	Parallel groups double blind
<a href="#">Farquharson , 2000</a> n=10/10 follow-up: 4 weeks	spironolactone 50 mg/d versus placebo	patients with NYHA class II to III chronic heart failure on standard diuretic/ACE inhibitor therapy	double blind
<a href="#">Macdonald , 2004</a> n=43/43 follow-up: 3 months	spironolactone 12.5-50 mg/d versus placebo	patients with New York Heart Association class I-II congestive heart failure taking optimal treatment (including beta blockers)	Cross over double blind
<a href="#">MacFadyen , 1997</a> n=21/16 follow-up: 8 weeks	spironolactone (50-100 mg/day) versus placebo	patients with stable chronic heart failure	Parallel groups double blind

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Mottram , 2004</b> n=30 follow-up: 6 months	spironolactone 25 mg/d versus placebo	hypertensive patients with diastolic heart failure	double blind
<b>RALES , 1998</b> n=822/841 follow-up: 24 mo	spironolactone (25 to 50 mg daily) versus placebo	patients with severe heart failure	Parallel groups Open World
<b>Tsutamoto , 2001</b> n=20/17 follow-up: 12 weeks	spironolactone 25 mg daily versus placebo	patients with mild-to-moderate nonischemic congestive heart failure	Parallel groups double blind Japan
<b>Yee , 2001</b> n=28/28 follow-up: 4 weeks	spironolactone 50mg/d versus placebo	patients with New York Heart Association class II to IV congestive heart failure	double blind
<b>TOPCAT , 2014</b> [NCT00094302] n=3445 follow-up: 3.3 years	spironolactone (15 to 45 mg daily) versus placebo	patients with heart failure and a preserved left ventricular ejection fraction of 45% or more	Parallel groups double-blind
<b>PIE II</b> <i>ongoing</i> [NCT00123955] n=NA follow-up: 9 months	Spironolactone 25mg tablet daily for 9 months versus placebo	elderly patients with isolated diastolic heart failure	Parallel groups double blind
<b>spironolactone+captopril vs captopril</b>			
<b>Han , 1994</b> n=19/16 follow-up: 4 weeks	captopril plus spironolactone versus captopril alone	patients with refractory CHF and New York Heart Association functional class IV without renal dysfunction, hypotension and hyperkalemia	open China
<b>furosemide + spironolactone vs prenalterol</b>			
<b>Dalhstrom , 1986</b> n=10/10 follow-up: 12 weeks	intensified treatment with diuretics (furosemide- spironolactone) versus prenalterol 100-200 mg daily in addition to their basal treatment	patients with severe chronic congestive heart failure (CHF) due to ischaemic heart disease treated with digitalis and diuretics	Cross over double blind
<b>spironolactone+furosemide vs spironolactone+butizide</b>			
<b>Mauersberger , 1985</b> n=22 follow-up:	spironolactone 50mg + furosemide 20 mg versus spironolactone 50mg + butizide 5mg	patients with congestive heart failure	open
<b>spironolactone vs spironolactone</b>			

continued...

Trial	Treatments	Patients	Trials design and methods
Nouvel essai <i>ongoing</i> [NCT00125437] n=NA follow-up:	spironolactone larger dose versus spironolactone standard dose	severe congestive heart failure in patients with nonischemic cardiomyopathy	Parallel groups single blind

More details and results :

- diuretics for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q75>
- diuretics for heart failure in patients with preserved-LVEF heart failure at <http://www.trialresultscenter.org/go-Q236>
- diuretics for heart failure in elderly at <http://www.trialresultscenter.org/go-Q314>
- aldosterone blockade for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q488>
- mineralocorticoid receptor antagonists for heart failure in all type of patients at <http://www.trialresultscenter.org/go-Q665>
- mineralocorticoid receptor antagonists for heart failure in HF pEF at <http://www.trialresultscenter.org/go-Q666>

## References

### Cicoira, 2002:

Cicoira M, Zanolla L, Rossi A, Golia G, Franceschini L, Brighetti G, Marino P, Zardini P, Long-term, dose-dependent effects of spironolactone on left ventricular function and exercise tolerance in patients with chronic heart failure. *J Am Coll Cardiol* 2002;40:304-10. [[12106936](#)]

### Cicoira, 2004:

Cicoira M, Rossi A, Bonapace S, Zanolla L, Perrot A, Francis DP, Golia G, Franceschini L, Osterziel KJ, Zardini P, Effects of ACE gene insertion/deletion polymorphism on response to spironolactone in patients with chronic heart failure. *Am J Med* 2004;116:657-61. [[15121491](#)] [10.1016/j.amjmed.2003.12.033](https://doi.org/10.1016/j.amjmed.2003.12.033)

### Ramires, 2000:

Ramires FJ, Mansur A, Coelho O, Maranhão M, Gruppi CJ, Mady C, Ramires JA, Effect of spironolactone on ventricular arrhythmias in congestive heart failure secondary to idiopathic dilated or to ischemic cardiomyopathy. *Am J Cardiol* 2000;85:1207-11. [[10802002](#)]

### Agostoni, 2005:

Agostoni P, Magini A, Andreini D, Contini M, Apostolo A, Bussotti M, Cattadori G, Palermo P, Spironolactone improves lung diffusion in chronic heart failure. *Eur Heart J* 2005;26:159-64. [[15618072](#)] [10.1093/eurheartj/ehi023](https://doi.org/10.1093/eurheartj/ehi023)

### Barr, 1995:

Barr CS, Lang CC, Hanson J, Arnott M, Kennedy N, Struthers AD, Effects of adding spironolactone to an angiotensin-converting enzyme inhibitor in chronic congestive heart failure secondary to coronary artery disease. *Am J Cardiol* 1995;76:1259-65. [[7503007](#)]

### Farquharson, 2000:

Farquharson CA, Struthers AD, Spironolactone increases nitric oxide bioactivity, improves endothelial vasodilator dysfunction, and suppresses vascular angiotensin I/angiotensin II conversion in patients with chronic heart failure. *Circulation* 2000;101:594-7. [[10673249](#)]

**Macdonald, 2004:**

Macdonald JE, Kennedy N, Struthers AD, Effects of spironolactone on endothelial function, vascular angiotensin converting enzyme activity, and other prognostic markers in patients with mild heart failure already taking optimal treatment. *Heart* 2004;90:765-70. [[15201246](#)] [10.1136/hrt.2003.017368](#)

**MacFadyen, 1997:**

MacFadyen RJ, Barr CS, Struthers AD, Aldosterone blockade reduces vascular collagen turnover, improves heart rate variability and reduces early morning rise in heart rate in heart failure patients. *Cardiovasc Res* 1997;35:30-4. [[9302344](#)]

**Mottram, 2004:**

Mottram PM, Haluska B, Leano R, Cowley D, Stowasser M, Marwick TH, Effect of aldosterone antagonism on myocardial dysfunction in hypertensive patients with diastolic heart failure. *Circulation* 2004;110:558-65. [[15277317](#)] [10.1161/01.CIR.0000138680.89536.A9](#)

**RALES, 1998:**

Effectiveness of spironolactone added to an angiotensin-converting enzyme inhibitor and a loop diuretic for severe chronic congestive heart failure (the Randomized Aldactone Evaluation Study [RALES]). *Am J Cardiol* 1996;78:902-7 [[8888663](#)]

Pitt B, Zannad F, Remme WJ, Cody R, Castaigne A, Perez A, Palensky J, Wittes J The effect of spironolactone on morbidity and mortality in patients with severe heart failure. Randomized Aldactone Evaluation Study Investigators. *N Engl J Med* 1999;341:709-17 [[10471456](#)] [10.1056/NEJM199909023411001](#)

Pitt D, ACE inhibitor co-therapy in patients with heart failure: rationale for the Randomized Aldactone Evaluation Study (RALES). *Eur Heart J* 1995;16 Suppl N:107-10. [[8682055](#)]

**Tsutamoto, 2001:**

Tsutamoto T, Wada A, Maeda K, Mabuchi N, Hayashi M, Tsutsui T, Ohnishi M, Sawaki M, Fujii M, Matsumoto T, Matsui T, Kinoshita M, Effect of spironolactone on plasma brain natriuretic peptide and left ventricular remodeling in patients with congestive heart failure. *J Am Coll Cardiol* 2001;37:1228-33. [[11300427](#)]

**Yee, 2001:**

Yee KM, Pringle SD, Struthers AD, Circadian variation in the effects of aldosterone blockade on heart rate variability and QT dispersion in congestive heart failure. *J Am Coll Cardiol* 2001;37:1800-7. [[11401114](#)]

**TOPCAT, 2014:**

Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Harty B, Heitner JF, Kenwood CT, Lewis EF, O'Meara E, Probstfield JL, Shaburishvili T, Shah SJ, Solomon SD, Sweitzer NK, Yang S, McKinlay SM Spironolactone for heart failure with preserved ejection fraction. *N Engl J Med* 2014;370:1383-92 [[24716680](#)] [10.1056/NEJMoa1313731](#)

**PIE II, :**

ongoing trial NCT00123955

**Han, 1994:**

Han YL, Tong M, Jing QM, Hu XL, Liu JQ, Combined therapy of captopril and spironolactone for refractory congestive heart failure. *Chin Med J (Engl)* 1994;107:688-92. [[7805462](#)]

**Dalhstrom, 1986:**

Jorde R, Pettersen JE, Burhol PG Lack of effect of exogenous or endogenous gastric inhibitory polypeptide on the elimination rate of Intralipid in man. *Acta Med Scand* 1984;216:19-23 [[6485877](#)]

Dahlstrm U, Areskog M, Wranne B, Karlsson E Prenalterol as long-term therapy for chronic congestive heart failure. A randomized cross-over trial. *Acta Med Scand* 1984;216:199-207 [[6149671](#)]

### Mauersberger, 1985:

Mauersberger H, Rangoonwala B, Ehrlich E, [Comparative study of 2 diuretic-containing combination preparations in patients with edematous heart failure] Wien Med Wochenschr 1985;135:205-13. [4013351]

### Nouvel essai, :

ongoing trial NCT00125437

## 2 heart failure with preserved LVEF

Trial	Treatments	Patients	Trials design and methods
<b>spironolactone vs placebo</b>			
TOPCAT, 2014 [NCT00094302] n=3445 follow-up: 3.3 years	spironolactone (15 to 45 mg daily) versus placebo	patients with heart failure and a preserved left ventricular ejection fraction of 45% or more	Parallel groups double-blind

More details and results :

- All mechanism for heart failure with preserved LVEF in all type of patients at <http://www.trialresultscenter.org/go-Q237>

## References

### TOPCAT, 2014:

Pitt B, Pfeffer MA, Assmann SF, Boineau R, Anand IS, Claggett B, Clausell N, Desai AS, Diaz R, Fleg JL, Gordeev I, Harty B, Heitner JF, Kenwood CT, Lewis EF, O'Meara E, Probstfield JL, Shaburishvili T, Shah SJ, Solomon SD, Sweitzer NK, Yang S, McKinlay SM Spironolactone for heart failure with preserved ejection fraction. N Engl J Med 2014;370:1383-92 [24716680] 10.1056/NEJMoa1313731

Entry terms: Veroshpiron, Verospirone, Spiractin, Spirobeta, Spirogamma, Spirolang, Spiro-Isis, Spiro-Isis, Spironone, Spirospare, Verospiron, Aldactone, Aldactone A, Aquareduct, duraspiron, Espironolactona Alter, Espironolactona Mundogen, Flumach, Frumikal, Jenaspiron, Novo-Spiroton, Novo Spiroton, NovoSpiroton, Practon, Spiro L.U.T., spiro von ct,