

Clinical trials of sympathectomie

TrialResults-center www.trialresultscenter.org

1 peripheral vascular diseases

| Trial | Treatments | Patients | Trials design and methods |
|---|---|------------------------------------|---------------------------|
| sympathectomie vs contrle | | | |
| Barnes RW , 1977 n=25/26 follow-up: 3 ans | sympathectomie bilatrale L2-L4 versus absence de sympathectomie | AOMI stade II (80%) et III (20%) | Parallel groups Ouvert |
| Satiani B , 1982 n=29/38 follow-up: 11 mois | sympathectomie bilatrale L2-L4 versus absence de sympathectomie | AOMI stade non precis | Parallel groups Ouvert |

More details and results :

- vasodilators therapy for peripheral vascular diseases in all type of patients at <http://www.trialresultscenter.org/go-Q49>

References

Barnes RW, 1977:

Value of concomitant sympathectomy in aortoiliac reconstruction. Results of a prospective, randomized study. Barnes RW, Baker WH, Shanik G, Maixner W, Hayes AC, Lin R, Clarke W Arch Surg 1977 Nov;112:1325-30 [[921531](#)]

Satiani B, 1982:

Prospective randomized study of concomitant lumbar sympathectomy with aortoiliac reconstruction. Satiani B, Liapis CD, Hayes JP, Kimmins S, Evans WE Am J Surg 1982 Jun;143:755-60 [[7091512](#)]