

Clinical trials of t-PA + streptokinase

TrialResults-center www.trialresultscenter.org

1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
t-PA + streptokinase vs streptokinase			
GUSTO tPA-SK Hiv , 1993 n=10374/20251 follow-up: 30 d	tPA en IV 1 mg/kg, sans dpasser 90 mg, dont 10 % en bolus + streptokinase 1 MU en 60 min + hparine en IV (5000 U en bolus, 1000 U/h (de prfrence 1200 U/h si >80 kg), poursuivi au moins 48 h) versus Streptokinase 1.5 MU en 60 min + hparine SC (12500 U 2 fois/j commence 4h aprs thrombolytique) combin streptokinase (1.5 MU en 60 min) + hparine en IV (5000 U en bolus, puis 1000 U/h (1200 U/h si >80 kg) poursuivi au moins 48 h)	Hommes et femmes	Parallel groups International 15 countries

More details and results :

- myocardial revascularization for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q129>
- fibrinolysis for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q260>

References

GUSTO tPA-SK Hiv, 1993:

An international randomized trial comparing four thrombolytic strategies for acute myocardial infarction. The GUSTO investigators. N Engl J Med 1993 Sep 2;329:673-82 [[8204123](#)]

Entry terms: streptokinase