

# Clinical trials of taspoglutide 10mg once weekly

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 diabetes type 2

Trial	Treatments	Patients	Trials design and methods
<b>taspoglutide 10mg once weekly vs placebo</b>			
<a href="#">Nauck 10 once weekly vs PBO , 2009</a> [NCT00423501] n=257/49 follow-up: 12 weeks	taspoglutide, either 5, 10, or 20 mg once weekly or 10 or 20 mg once every 2 weeks for 8 weeks versus placebo	patients with type 2 diabetes inadequately controlled with metformin	Parallel groups double-blind

More details and results :

- insulin secretagogues peptides (incretins) for diabetes type 2 in all type of patients at <http://www.trialresultscenter.org/go-Q381>
- antidiabetic drugs for diabetes type 2 in patients inadequately controlled on metformin at <http://www.trialresultscenter.org/go-Q509>
- antidiabetic drugs for diabetes type 2 in patients inadequately controlled on monotherapy at <http://www.trialresultscenter.org/go-Q512>

## References

### Nauck 10 once weekly vs PBO, 2009:

Nauck MA, Ratner RE, Kapitza C, Berria R, Boldrin M, Balena R, Treatment with the human once-weekly glucagon-like peptide-1 analog taspoglutide in combination with metformin improves glycemic control and lowers body weight in patients with type 2 diabetes inadequately controlled with metformin alone: a double-blind placebo-controlled study. Diabetes Care 2009;32:1237-43. [[19366970](#)] [10.2337/dc08-1961](#)

Entry terms: taspoglutide, taspoglutide,