

Clinical trials of transfer for primary angioplasty

TrialResults-center www.trialresultscenter.org

1 acute myocardial infarction

Trial	Treatments	Patients	Trials design and methods
transfer for primary angioplasty vs immediate thrombolysis			
AIR-PAMI , 2002 n=71/66 follow-up:	Transfer for Primary Angioplasty versus immediate thrombolysis (various thrombolytic)	Patients with high-risk AMI (age >70 years, anterior MI, Killip class II/III, heart rate >100 beats/min or systolic BP <100 mm Hg), eligible for thrombolytic therapy	Parallel groups open
DANAMI-2 , 2003 n=567/562 follow-up: 30 days	Transfer for Primary Angioplasty versus immediate thrombolysis with tPA (accelerated infusion)	patients with myocardial infarction with ST-segment elevation	Parallel groups open
PRAGUE-2 , 2003 n=429/421 follow-up: 30 days	immediate transport for primary percutaneous coronary intervention versus immediate thrombolysis with streptokinase	patients with acute ST elevation myocardial infarction presenting within <12 h to the nearest community hospital without a catheter laboratory	Parallel groups open

More details and results :

- myocardial revascularization for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q129>
- PCI for acute myocardial infarction in all type of patients at <http://www.trialresultscenter.org/go-Q246>

References

AIR-PAMI , 2002:

Grines CL, Westerhausen DR Jr, Grines LL, Hanlon JT, Logemann TL, Niemela M, Weaver WD, Graham M, Boura J, O'Neill WW, Balestrini C A randomized trial of transfer for primary angioplasty versus on-site thrombolysis in patients with high-risk myocardial infarction: the Air Primary Angioplasty in Myocardial Infarction study. J Am Coll Cardiol 2002;39:1713-9 [[12039480](#)]

DANAMI-2 , 2003:

Fosbl EL, Thune JJ, Kelbaek H, Andersen HR, Saunamki K, Nielsen TT, Mortensen LS, Kber L Long-term outcome of primary angioplasty compared with fibrinolysis across age groups: a Danish Multicenter Randomized Study on Fibrinolytic Therapy Versus Acute Coronary Angioplasty in Acute Myocardial Infarction (DANAMI-2) substudy. Am Heart J 2008 Aug;156:391-6 [[18657676](#)]

Madsen JK, Grande P, Saunamki K, Thayssen P, Kassis E, Eriksen U, Rasmussen K, Hauns S, Nielsen TT, Haghfelt T, Fritz-Hansen P, Hjelms E, Paulsen PK, Alstrup P, Arendrup H, Niebuhr-Jrgensen U, Andersen LI Danish multicenter randomized study of invasive versus conservative treatment in patients with inducible ischemia after thrombolysis in acute myocardial infarction (DANAMI). DANish trial in Acute Myocardial Infarction. *Circulation* 1997 Aug 5;96:748-55 [[9264478](#)]

Madsen JK, Nielsen TT, Grande P, Eriksen UH, Saunamki K, Thayssen P, Kassis E, Rasmussen K, Hauns S, Haghfelt T, Fritz-Hansen P, Hjelms E, Paulsen PK, Alstrup P, Arendrup H, Niebuhr-Jrgensen U, Andersen LI Revascularization compared to medical treatment in patients with silent vs. symptomatic residual ischemia after thrombolysed myocardial infarction—the DANAMI study. *Cardiology* 2007;108:243-51 [[17114878](#)]

Andersen HR, Nielsen TT, Rasmussen K, Thuesen L, Kelbaek H, Thayssen P, Abildgaard U, Pedersen F, Madsen JK, Grande P, Villadsen AB, Krusell LR, Haghfelt T, Lomholt P, Husted SE, Vigholt E, Kjaergard HK, Mortensen LS A comparison of coronary angioplasty with fibrinolytic therapy in acute myocardial infarction. *N Engl J Med* 2003 Aug 21;349:733-42 [[12930925](#)]

PRAGUE-2 , 2003:

Widimsk P, Budesnsk T, Vorc D, Groch L, Zelzko M, Aschermann M, Branny M, St'sek J, Formnek P Long distance transport for primary angioplasty vs immediate thrombolysis in acute myocardial infarction. Final results of the randomized national multicentre trial—PRAGUE-2. *Eur Heart J* 2003;24:94-104 [[12559941](#)]

Widimsk P, Budesnsk T, Vorc D, Groch L, Zelzko M, Aschermann M, Branny M, St'sek J, Formnek P Long distance transport for primary angioplasty vs immediate thrombolysis in acute myocardial infarction. Final results of the randomized national multicentre trial—PRAGUE-2. *Eur Heart J* 2003;24:94-104 [[12559941](#)]