

# Clinical trials of warfarin + aspirin

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 cardiovascular prevention

Trial	Treatments	Patients	Trials design and methods
<b>warfarin + aspirin vs placebo</b>			
<b>Thrombosis Prevention trial (W plus A) , 1998</b> [NCT00000614] n=1277/1272 follow-up: median 6.8 y	warfarin adjusted dose for INR of 1.5 + aspirin 75 mg daily versus placebo	men aged between 45 years and 69 years at high risk of IHD	NA double blind UK

More details and results :

- anticoagulant for cardiovascular prevention in all type of patients at <http://www.trialresultscenter.org/go-Q709>
- anticoagulant for cardiovascular prevention in primary prevention at <http://www.trialresultscenter.org/go-Q710>

## References

### Thrombosis Prevention trial (W plus A), 1998:

Thrombosis prevention trial: randomised trial of low-intensity oral anticoagulation with warfarin and low-dose aspirin in the primary prevention of ischaemic heart disease in men at increased risk. The Medical Research Council's General Practice Research Framework. Lancet 1998;351:233-41 [[9457092](#)]

Meade TW, Wilkes HC, Stirling Y, Brennan PJ, Kelleher C, Browne W Randomized controlled trial of low dose warfarin in the primary prevention of ischaemic heart disease in men at high risk: design and pilot study. Eur Heart J 1988;9:836-43 [[3053176](#)]

## 2 atrial fibrillation

Trial	Treatments	Patients	Trials design and methods
<b>warfarin + aspirin vs warfarin standard dose</b>			
<b>AFASAK II (warfarin low dose+aspirin vs warfarin standard dose) , 1998</b> n=171/170 follow-up: 3.5 years	warfarin fixed low dose(1.25mg/d) + aspirin(300mg/d) versus warfarin standard dose(target INR 2.0-3.0)	chronic non valvular atrial fibrillation	Parallel groups Open Denmark

continued...

Trial	Treatments	Patients	Trials design and methods
SPAF III , 1996 n=521/523 follow-up: 1.1 years	warfarin low dose(target INR 1.2-1.5)+ aspirin 325 mg/d versus warfarin standard dose(target INR 2.0-3.0)	non rheumatic atrial fibrillation,patients with at least one additional thromboembolic risk factor(high risk patients)	Parallel groups Open USA,Canada

More details and results :

- antithrombotics for atrial fibrillation in primary prevention of thromboembolic events at <http://www.trialresultscenter.org/go-Q57>

## References

### AFASAK II (warfarin low dose+aspirin vs warfarin standard dose), 1998:

Gullov AL, Koefoed BG, Petersen P, Pedersen TS, Andersen ED, Godtfredsen J, Boysen G Fixed minidose warfarin and aspirin alone and in combination vs adjusted-dose warfarin for stroke prevention in atrial fibrillation: Second Copenhagen Atrial Fibrillation, Aspirin, and Anticoagulation Study. Arch Intern Med 1998 Jul 27;158:1513-21 [[9679792](#)]

### SPAF III, 1996:

Adjusted-dose warfarin versus low-intensity, fixed-dose warfarin plus aspirin for high-risk patients with atrial fibrillation: Stroke Prevention in Atrial Fibrillation III randomised clinical trial. Lancet. 1996 Sep 7;348(9028):633-8. [[8782752](#)]

Entry terms: warfarin