

Clinical trials of warfarin low dose + aspirin

TrialResults-center www.trialresultscenter.org

1 atrial fibrillation

Trial	Treatments	Patients	Trials design and methods
warfarin low dose + aspirin vs control			
SAFT(warfarin low dose + aspirin vs no treatment) , 2003 n=334/334 follow-up: 33 months	warfarin low dose (1.25 mg/d) + aspirin 75 mg/d versus no treatment	Low-medium risk patients with non valvular atrial fibrillation.	Parallel groups Open Sweden

More details and results :

- antithrombotics for atrial fibrillation in primary prevention of thromboembolic events at <http://www.trialresultscenter.org/go-Q57>

References

SAFT(warfarin low dose + aspirin vs no treatment), 2003:

Edvardsson N, Juul-Moller S, Omblus R, Pehrsson K Effects of low-dose warfarin and aspirin versus no treatment on stroke in a medium-risk patient population with atrial fibrillation. J Intern Med 2003 Jul;254:95-101 [[12823646](#)]

Edvardsson N, Juul-Moller S, Omblus R, Pehrsson K Effects of low-dose warfarin and aspirin versus no treatment on stroke in a medium-risk patient population with atrial fibrillation. J Intern Med 2003 Jul;254:95-101 [[12823646](#)]

Entry terms: warfarin