

Clinical trials of antithrombotics for venous thrombosis in all type of patients

TrialResults-center www.trialresultscenter.org

1 anticoagulant

Trial	Treatments	Patients	Trials design and methods
warfarin vs discontinuation			
PREVENT (Ridker) , 2003 n=255/253 follow-up: 2.1 years	extension with low-intensity warfarin (target INR, 1.5 to 2.0) versus placebo	Patients with idiopathic venous thromboembolism who had received full-dose anticoagulation therapy for a median of 6.5 months	Parallel groups
Agnelli , 2003 n=NA follow-up: 33 months	continuation for 3 or 9 additional months of warfarin or other oral anticoagulant was adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 3 months)	patients who had had 3 months of oral anticoagulant therapy without experiencing recurrence or bleeding after a first episode of pulmonary embolism	Parallel groups open Italy
Agnelli , 2001 n=NA follow-up: 33 months	continuation for 9 additional months; warfarin or acenocoumarol adjusted to achieve a target INR between 2.0 and 3.0 versus discontinuation (after 3 months months)	Patients with a first episode of idiopathic proximal deep venous thrombosis who had completed three months of oral anticoagulant therapy	Parallel groups open Italy
LAFIT (Kearon) , 1999 n=NA follow-up:	Continuation of the oral anticoagulant therapy up to 24 months, warfarin was adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 3 months)	patients who had completed 3 months of anticoagulant therapy for a first episode of idiopathic venous thromboembolism	
ELAET (Kearon) , 2004 n=NA follow-up: 11 months (after randomizatio)	continuation for 2 additional months of warfarin adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 1 months)	-	Parallel groups double blind Canada, US
Levine , 1995 n=NA follow-up: 11 months after randomization.	continuation for 2 months of warfarin adjusted INR value of 2.0 to 3.0 versus Discontinue oral anticoagulant therapy (after 1 months)	Patients with venographically confirmed acute proximal DVT who had received four weeks of warfarin after initial heparin and whose four week IPG was normal	Parallel groups double blind Canada, Italy
DURAC (Schulman) , 1997 n=NA follow-up: Four years after randomization	indefinite warfarin or dicoumarol adjusted for a target INR between 2.0 and 2.85 versus 6 months warfarin or dicoumarol adjusted for a target INR between 2.0 and 2.85	-	Parallel groups open Sweden
heparin+warfarin vs placebo			

continued...

Trial	Treatments	Patients	Trials design and methods
Ott import , 1998 n=11/12	anticoagulants (s.c. heparin followed by oral warfarin) (duration NA) versus s.c. saline followed by oral placebo tablets	-	double blind Denmark
heparin+phenprocoumon vs phenylbutazone			
Nielsen import , 1994 n=48/42	heparin and phenprocoumon for 3 months versus phenylbutazone	-	open Denmark

References

PREVENT (Ridker), 2003:

Ridker PM, Goldhaber SZ, Danielson E, Rosenberg Y, Eby CS, Deitcher SR, Cushman M, Moll S, Kessler CM, Elliott CG, Paulson R, Wong T, Bauer KA, Schwartz BA, Miletich JP, Bounameaux H, Glynn RJ Long-term, low-intensity warfarin therapy for the prevention of recurrent venous thromboembolism. *N Engl J Med* 2003;348:1425-34 [[12601075](#)]

Agnelli, 2003:

Agnelli G, Prandoni P, Becattini C, Silingardi M, Taliani MR, Miccio M, Imberti D, Poggio R, Ageno W, Pogliani E, Porro F, Zonzin P Extended oral anticoagulant therapy after a first episode of pulmonary embolism. *Ann Intern Med* 2003;139:19-25 [[12834314](#)]

Agnelli, 2001:

Agnelli G, Prandoni P, Santamaria MG, Bagatella P, Iorio A, Bazzan M, Moia M, Guazzaloca G, Bertoldi A, Tomasi C, Scannapieco G, Ageno W Three months versus one year of oral anticoagulant therapy for idiopathic deep venous thrombosis. Warfarin Optimal Duration Italian Trial Investigators. *N Engl J Med* 2001;345:165-9 [[11463010](#)]

LAFIT (Kearon), 1999:

Kearon C, Gent M, Hirsh J, Weitz J, Kovacs MJ, Anderson DR, Turpie AG, Green D, Ginsberg JS, Wells P, MacKinnon B, Julian JA A comparison of three months of anticoagulation with extended anticoagulation for a first episode of idiopathic venous thromboembolism. *N Engl J Med* 1999;340:901-7 [[10089183](#)]

Kearon C, Gent M, Hirsh J, Weitz J, Kovacs MJ, Anderson DR, Turpie AG, Green D, Ginsberg JS, Wells P, MacKinnon B, Julian JA A comparison of three months of anticoagulation with extended anticoagulation for a first episode of idiopathic venous thromboembolism. *N Engl J Med* 1999;340:901-7 [[10089183](#)]

ELAET (Kearon), 2004:

Kearon C, Ginsberg JS, Anderson DR, Kovacs MJ, Wells P, Julian JA, Mackinnon B, Demers C, Douketis J, Turpie AG, Van Nguyen P, Green D, Kassis J, Kahn SR, Solymoss S, Desjardins L, Geerts W, Johnston M, Weitz JI, Hirsh J, Gent M Comparison of 1 month with 3 months of anticoagulation for a first episode of venous thromboembolism associated with a transient risk factor. *J Thromb Haemost* 2004;2:743-9 [[15099280](#)]

Levine, 1995:

Levine MN, Hirsh J, Gent M, Turpie AG, Weitz J, Ginsberg J, Geerts W, LeClerc J, Neemeh J, Powers P Optimal duration of oral anticoagulant therapy: a randomized trial comparing four weeks with three months of warfarin in patients with proximal deep vein thrombosis. *Thromb Haemost* 1995;74:606-11 [[8584992](#)]

DURAC (Schulman), 1997:

Schulman S, Granqvist S, Holmström M, Carlsson A, Lindmarker P, Nicol P, Eklund SG, Nordlander S, Lrfars G, Leijd B, Linder O, Loogna E The duration of oral anticoagulant therapy after a second episode of venous thromboembolism. The Duration of Anticoagulation Trial Study Group. *N Engl J Med* 1997;336:393-8 [[9010144](#)]

Ott import, 1998:

Ott P, Eldrup E, Oxholm P [Value of anticoagulant therapy in deep venous thrombosis in the lower limb in elderly, mobilized patients. A double-blind placebo controlled study with open therapeutic guidance] *Ugeskr Laeger* 1988;150:218-21 [[3287734](#)]

Nielsen import, 1994:

Nielsen HK, Husted SE, Krusell LR, Fasting H, Charles P, Hansen HH Silent pulmonary embolism in patients with deep venous thrombosis. Incidence and fate in a randomized, controlled trial of anticoagulation versus no anticoagulation. *J Intern Med* 1994;235:457-61 [[8182402](#)]

Nielsen HK, Husted SE, Krusell LR, Fasting H, Charles P, Hansen HH, Nielsen BO, Petersen JB, Bechgaard P Anticoagulant therapy in deep venous thrombosis. A randomized controlled study. *Thromb Res* 1994;73:215-26 [8191414]

2 direct anti-Xa

Trial	Treatments	Patients	Trials design and methods
apixaban 2.5mg vs discontinuation			
AMPLIFY-EXT 2.5mg , 2012 [NCT00633893] n=842/829 follow-up: 12 mo	Extended Treatment with apixaban 2.5 mg twice daily 12 months versus placebo	patients who have completed their intended treatment for deep vein thrombosis or pulmonary embolism	Parallel groups double blind
apixaban 5mg vs discontinuation			
AMPLIFY-EXT 5mg , 2012 [NCT00633893] n=815/829 follow-up: 12 mo	Extended Treatment with apixaban 5 mg twice daily 12 months versus placebo	patients who have completed their intended treatment for deep vein thrombosis or pulmonary embolism	double blind
rivaroxaban vs discontinuation			
EINSTEIN-extension , 2009 [NCT00439725] n=602/595 follow-up:	rivaroxaban 20 mg once-daily for an additional 6 or 12 months versus placebo	patients who had completed six to 12 months of anticoagulant treatment for an acute episode of VTE	Parallel groups double blind 28 countries
apixaban (without LMWH) vs LMWH/VKA			
AMPLIFY , 2013 [NCT00643201] n=2691/2704 follow-up: 6 mo	apixaban 10 mg twice daily for 7 days then 5 mg, twice daily, 6 months versus conventional therapy: enoxaparin 1mg/kg twice daily until INR>=2 then warfarin for an INR between 2-4, once daikly, 6 months	patients with deep vein thrombosis or pulmonary embolism	Parallel groups double blind
Botticelli DVT , 2008 [NCT00252005] n=358/118 follow-up:	apixaban 5 mg twice-daily, 10 mg twice-daily, or 20 mg once-daily for 84-91 days versus low molecular weight heparin followed by vitamin K antagonists	patients with symptomatic deep vein thrombosis	Parallel groups open
rivaroxaban (without LMWH) vs LMWH/VKA			
Einstein-DVT Dose-Ranging Study , 2008 n=NA follow-up:	rivaroxaban 20, 30, or 40 mg once daily versus low-molecular-weight heparin followed by vitamin K antagonists	patients with deep vein thrombosis	open
Einstein-DVT Evaluation , 2010 [NCT00440193] n=1731/1718 follow-up:	rivaroxaban 15 mg twice daily for 3 weeks, then 20 mg daily versus enoxaparin 1 mg/kg twice daily >=5 days, then warfarin with target INR between 2-3	Patients with Confirmed Acute Symptomatic Deep-Vein Thrombosis without Pulmonary Embolism	Parallel groups open (assessor-blind)

continued...

Trial	Treatments	Patients	Trials design and methods
Einstein-PE Evaluation , 2012 [NCT00439777] n=2419/2413 follow-up: 9.8 months	rivaroxaban (15 mg twice daily for 3 weeks, followed by 20 mg once daily) for 3, 6, or 12 months versus standard therapy with enoxaparin followed by an adjusted-dose vitamin K antagonist	patients who had acute symptomatic pulmonary embolism with or without deep-vein thrombosis	Parallel groups open 38 countries
ximelagatran vs LMWH/VKA			
Fiessinger , 2005 n=NA follow-up:	ximelagatran 36 mg twice daily versus subcutaneous enoxaparin, 1 mg/kg twice daily, for 5 to 20 days followed by warfarin adjusted to maintain an international normalized ratio of 2.0 to 3.0.	patients with acute deep vein thrombosis	double blind
ximelagatran (without LMWH) vs LMWH/VKA			
THRIVE I , 2003 n=NA follow-up:	oral ximelagatran (24, 36, 48 or 60 mg twice daily) for 2 weeks versus dalteparin and warfarin for 2 weeks	Patients with acute DVT	

References

AMPLIFY-EXT 2.5mg, 2012:

Agnelli G, Buller HR, Cohen A, Curto M, Gallus AS, Johnson M, Porcari A, Raskob GE, Weitz JI Apixaban for Extended Treatment of Venous Thromboembolism. N Engl J Med 2012 Dec 8;; [23216615] [10.1056/NEJMoa1207541](https://doi.org/10.1056/NEJMoa1207541)

AMPLIFY-EXT 5mg, 2012:

Agnelli G, Buller HR, Cohen A, Curto M, Gallus AS, Johnson M, Porcari A, Raskob GE, Weitz JI Apixaban for Extended Treatment of Venous Thromboembolism. N Engl J Med 2012 Dec 8;; [23216615] [10.1056/NEJMoa1207541](https://doi.org/10.1056/NEJMoa1207541)

EINSTEIN-extension, 2009:

Buller H R. Once daily oral rivaroxaban versus placebo in the long term treatment of recurrent symptomatic venous thromboembolism. The Einstein-extension study. ASH, 8 dcembre 2009

Oral Rivaroxaban for Symptomatic Venous Thromboembolism. N Engl J Med 2010 Dec 3;; [21128814] [10.1056/NEJMoa1007903](https://doi.org/10.1056/NEJMoa1007903)

AMPLIFY, 2013:

Agnelli G, Buller HR, Cohen A, Curto M, Gallus AS, Johnson M, Masiukiewicz U, Pak R, Thompson J, Raskob GE, Weitz JI Oral apixaban for the treatment of acute venous thromboembolism. N Engl J Med 2013 Aug 29;369:799-808 [23808982] [10.1056/NEJMoa1302507](https://doi.org/10.1056/NEJMoa1302507)

Botticelli DVT, 2008:

Buller H, Deitchman D, Prins M, Segers A Efficacy and safety of the oral direct factor Xa inhibitor apixaban for symptomatic deep vein thrombosis. The Botticelli DVT dose-ranging study. J Thromb Haemost 2008 Aug;6:1313-8 [18541000]

Einstein-DVT Dose-Ranging Study, 2008:

Buller HR, Lensing AW, Prins MH, Agnelli G, Cohen A, Gallus AS, Misselwitz F, Raskob G, Schellong S, Segers A A dose-ranging study evaluating once-daily oral administration of the factor Xa inhibitor rivaroxaban in the treatment of patients with acute symptomatic deep vein thrombosis: the Einstein-DVT Dose-Ranging Study. Blood 2008;112:2242-7 [18621928]

Einstein-DVT Evaluation, 2010:

Oral Rivaroxaban for Symptomatic Venous Thromboembolism. N Engl J Med 2010 Dec 3;; [21128814] [10.1056/NEJMoa1007903](https://doi.org/10.1056/NEJMoa1007903)

Einstein-PE Evaluation, 2012:

Oral Rivaroxaban for the Treatment of Symptomatic Pulmonary Embolism. N Engl J Med 2012 Mar 26; [22449293] 10.1056/NEJMoa1113572

Fiessinger , 2005:

Fiessinger JN, Huisman MV, Davidson BL, Bounameaux H, Francis CW, Eriksson H, Lundstrm T, Berkowitz SD, Nystrm P, Thorsn M, Ginsberg JS Ximelagatran vs low-molecular-weight heparin and warfarin for the treatment of deep vein thrombosis: a randomized trial. JAMA 2005 Feb 9;293:681-9 [15701909]

THRIVE I, 2003:

Eriksson H, Whlander K, Gustafsson D, Welin LT, Frison L, Schulman S A randomized, controlled, dose-guiding study of the oral direct thrombin inhibitor ximelagatran compared with standard therapy for the treatment of acute deep vein thrombosis: THRIVE I. J Thromb Haemost 2003 Jan;1:41-7 [12871538]

3 extended anticoagulant

Trial	Treatments	Patients	Trials design and methods
warfarin vs discontinuation			
PREVENT (Ridker) , 2003 n=255/253 follow-up: 2.1 years	extension with low-intensity warfarin (target INR, 1.5 to 2.0) versus placebo	Patients with idiopathic venous thromboembolism who had received full-dose anticoagulation therapy for a median of 6.5 months	Parallel groups
Agnelli , 2003 n=NA follow-up: 33 months	continuation for 3 or 9 additional months of warfarin or other oral anticoagulant was adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 3 months)	patients who had had 3 months of oral anticoagulant therapy without experiencing recurrence or bleeding after a first episode of pulmonary embolism	Parallel groups open Italy
Agnelli , 2001 n=NA follow-up: 33 months	continuation for 9 additional months; warfarin or acenocoumarol adjusted to achieve a target INR between 2.0 and 3.0 versus discontinuation (after 3 months months)	Patients with a first episode of idiopathic proximal deep venous thrombosis who had completed three months of oral anticoagulant therapy	Parallel groups open Italy
LAFIT (Kearon) , 1999 n=NA follow-up:	Continuation of the oral anticoagulant therapy up to 24 months, warfarin was adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 3 months)	patients who had completed 3 months of anticoagulant therapy for a first episode of idiopathic venous thromboembolism	
ELAET (Kearon) , 2004 n=NA follow-up: 11 months (after randomizatio)	continuation for 2 additional months of warfarin adjusted to achieve a target INR between 2.0 and 3.0. versus discontinuation (after 1 months)	-	Parallel groups double blind Canada, US
Levine , 1995 n=NA follow-up: 11 months after randomization.	continuation for 2 months of warfarin adjusted INR value of 2.0 to 3.0 versus Discontinue oral anticoagulant therapy (after 1 months)	Patients with venographically confirmed acute proximal DVT who had received four weeks of warfarin after initial heparin and whose four week IPG was normal	Parallel groups double blind Canada, Italy

continued...

Trial	Treatments	Patients	Trials design and methods
DURAC (Schulman) , 1997 n=NA follow-up: Four years after randomization	indefinite warfarin or dicoumarol adjusted for a target INR between 2.0 and 2.85 versus 6 months warfarin or dicoumarol adjusted for a target INR between 2.0 and 2.85	-	Parallel groups open Sweden

References

PREVENT (Ridker), 2003:

Ridker PM, Goldhaber SZ, Danielson E, Rosenberg Y, Eby CS, Deitcher SR, Cushman M, Moll S, Kessler CM, Elliott CG, Paulson R, Wong T, Bauer KA, Schwartz BA, Miletich JP, Bounameaux H, Glynn RJ Long-term, low-intensity warfarin therapy for the prevention of recurrent venous thromboembolism. *N Engl J Med* 2003;348:1425-34 [[12601075](#)]

Agnelli, 2003:

Agnelli G, Prandoni P, Becattini C, Silingardi M, Taliani MR, Miccio M, Imberti D, Poggio R, Ageno W, Pogliani E, Porro F, Zonzin P Extended oral anticoagulant therapy after a first episode of pulmonary embolism. *Ann Intern Med* 2003;139:19-25 [[12834314](#)]

Agnelli, 2001:

Agnelli G, Prandoni P, Santamaria MG, Bagatella P, Iorio A, Bazzan M, Moia M, Guazzaloca G, Bertoldi A, Tomasi C, Scannapieco G, Ageno W Three months versus one year of oral anticoagulant therapy for idiopathic deep venous thrombosis. Warfarin Optimal Duration Italian Trial Investigators. *N Engl J Med* 2001;345:165-9 [[11463010](#)]

LAFIT (Kearon), 1999:

Kearon C, Gent M, Hirsh J, Weitz J, Kovacs MJ, Anderson DR, Turpie AG, Green D, Ginsberg JS, Wells P, MacKinnon B, Julian JA A comparison of three months of anticoagulation with extended anticoagulation for a first episode of idiopathic venous thromboembolism. *N Engl J Med* 1999;340:901-7 [[10089183](#)]

Kearon C, Gent M, Hirsh J, Weitz J, Kovacs MJ, Anderson DR, Turpie AG, Green D, Ginsberg JS, Wells P, MacKinnon B, Julian JA A comparison of three months of anticoagulation with extended anticoagulation for a first episode of idiopathic venous thromboembolism. *N Engl J Med* 1999;340:901-7 [[10089183](#)]

ELAET (Kearon), 2004:

Kearon C, Ginsberg JS, Anderson DR, Kovacs MJ, Wells P, Julian JA, Mackinnon B, Demers C, Douketis J, Turpie AG, Van Nguyen P, Green D, Kassis J, Kahn SR, Solymoss S, Desjardins L, Geerts W, Johnston M, Weitz JI, Hirsh J, Gent M Comparison of 1 month with 3 months of anticoagulation for a first episode of venous thromboembolism associated with a transient risk factor. *J Thromb Haemost* 2004;2:743-9 [[15099280](#)]

Levine, 1995:

Levine MN, Hirsh J, Gent M, Turpie AG, Weitz J, Ginsberg J, Geerts W, LeClerc J, Neemeh J, Powers P Optimal duration of oral anticoagulant therapy: a randomized trial comparing four weeks with three months of warfarin in patients with proximal deep vein thrombosis. *Thromb Haemost* 1995;74:606-11 [[8584992](#)]

DURAC (Schulman), 1997:

Schulman S, Granqvist S, Holmström M, Carlsson A, Lindmarker P, Nicol P, Eklund SG, Nordlander S, Lrfars G, Leijd B, Linder O, Loogna E The duration of oral anticoagulant therapy after a second episode of venous thromboembolism. The Duration of Anticoagulation Trial Study Group. *N Engl J Med* 1997;336:393-8 [[9010144](#)]

4 extended LMWH

Trial	Treatments	Patients	Trials design and methods
Enoxaparin vs acenocoumarol			

continued...

Trial	Treatments	Patients	Trials design and methods
Veiga , 2000 n=50/50 follow-up: 6-9 mo	UFH, APTT 1.52.0d followed by Enoxaparin 4,000 IU qd versus UFH, APTT 1.52.0d followed by Acenocoumarol target INR 2-3	patients with objective diagnosis of DVT by Venography	open
Nadroparin vs acenocoumarol			
Lopez-Beret , 2001 n=81/77 follow-up: 6-9 mo	LMWH, 1,025 IU/10 kg bid followed by Nadroparin 1,025 IU/10 kg bid versus LMWH, 1,025 IU/10 kg bid followed by Acenocoumarol target INR 2-3	patients with objective diagnosis of DVT by compression ultrasonography	open
Lopaciuk , 1999 n=101/101 follow-up: 9 mo	LMWH, 85 UI/kg bid followed by Nadroparin 85 IU/kg qd versus LMWH, 85 UI/kg bid followed by Acenocoumarol target INR 2-3	patients with objective diagnosis of DVT by Venography	open
Tinzaparin vs acenocoumarol			
Romera , 2009 n=119/122 follow-up: 12 months	tinzaparin SC 175 IU anti-Xa per kg once daily for 6 months versus acenocoumarol for target INR 2-3 for 6 months after initial LMWH (until INR 2-3)	patients with symptomatic proximal DVT of the lowerlimbs confirmed by compression duplex ultrasound scan	Parallel groups open Spain
Enoxaparin vs coumarin			
Gonzalez-Fajardo , 2008 n=85/80 follow-up: 1y, 5y	long-term anticoagulant treatment with enoxaparin during at least 3 months versus long-term anticoagulant treatment with coumarin during at least 3 months	patients with symptomatic, unilateral, first-episode DVT	Parallel groups open, blind assessment Spain
Bemiparin vs warfarin			
Kakkar , 2003 n=221/103 follow-up: 3 mo	LMWH, 115 IU/kg qd followed by Bemiparin 3,500 IU qd versus A: UFH, 30/40,000IU qd; B: LMWH, 115 IU/kg qd followed by Warfarin target INR 2-3	patients with objective diagnosis of DVT by Venography/compression ultrasonography	open
Dalteparin vs warfarin			
Lee , 2003 n=336/336 follow-up: 6 mo	LMWH, 200 IU/kg qd followed by Dalteparin 150 IU/kg qd versus LMWH, 200 IU/kg qd followed by Warfarin target INR 2-3	patients with cancer and objective diagnosis of DVT by Venography/compression ultrasonography	open
Das , 1996 n=50/55 follow-up: 3 mo	UFH followed by Dalteparin 5,000 IU qd versus UFH followed by Warfarin target INR 2-3	patients with objective diagnosis of DVT by Venography	open
Enoxaparin vs warfarin			

continued...

Trial	Treatments	Patients	Trials design and methods
Deitcher , 2003 n=51/30 follow-up: 6 mo	LMWH: 1a, 1 mg/kg q12h; 1b, 1 mg/kg qd12h followed by Enoxaparin 1a: 1 mg/kg qd; 1b: 1.5 mg/kg qd versus LMWH, 1 mg/kg q12h followed by Warfarin target INR 2-3	patients with objective diagnosis of DVT	open
Meyer , 2002 n=71/58 follow-up: 3 mo	LMWH, 1.5 mg/kg qd followed by Enoxaparin 1.5 mg/Kg qd versus LMWH, 1.5 mg/kg qd followed by Warfarin target INR 2-3	patients with cancer and objective diagnosis of DVT by Venography/compression ultrasonography	open
Gonzalez-Fajardo , 1999 n=93/92 follow-up: 9 mo	LMWH, 4,000 IU bid followed by Enoxaparin 4,000 IU qd versus UFH followed by Warfarin target INR 2-3	patients with objective diagnosis of DVT by Venography	Parallel groups open
Pini , 1994 n=93/94 follow-up: 9 mo	UFH, APTT 1.31.9 followed by Enoxaparin 4,000 IU qd versus UFH, APTT 1.31.9 followed by Warfarin target INR 2-3.5	patients with objective diagnosis of DVT by Venography (diagnosed by strain-gauge plethysmography plus D-dimer latex assay and confirmed by venography)	open
Tinzaparin vs warfarin			
Hull , 2002 n=369/368 follow-up: 9 mo	LMWH, 175 IU/kg qd followed by Tinzaparin 175 IU/kg qd versus UFH 5 d, followed by UFH therapeutic APTT followed by Warfarin target INR 2-3	patients with objective diagnosis of DVT by Venography/compression ultrasonography	open

References

Veiga, 2000:

Veiga F, Escriv A, Maluenda MP, Lpez Rubio M, Margalet I, Lezana A, Gallego J, Ribera JM Low molecular weight heparin (enoxaparin) versus oral anticoagulant therapy (acenocoumarol) in the long-term treatment of deep venous thrombosis in the elderly: a randomized trial. *Thromb Haemost* 2000;84:559-64 [[11057850](#)]

Lopez-Beret, 2001:

Lpez-Beret P, Orgaz A, Fontcuberta J, Doblaz M, Martinez A, Lozano G, Romero A Low molecular weight heparin versus oral anticoagulants in the long-term treatment of deep venous thrombosis. *J Vasc Surg* 2001;33:77-90 [[11137927](#)]

Lopaciuk, 1999:

Lopaciuk S, Bielska-Falda H, Noszczyk W, Bielawiec M, Witkiewicz W, Filipecki S, Michalak J, Ciesielski L, Mackiewicz Z, Czestochowska E, Zawilska K, Cencora A Low molecular weight heparin versus acenocoumarol in the secondary prophylaxis of deep vein thrombosis. *Thromb Haemost* 1999;81:26-31 [[9974369](#)]

Romera, 2009:

Romera A, Cairols MA, Vila-Coll R, Mart X, Colom E, Bonell A, Lapiedra O A randomised open-label trial comparing long-term sub-cutaneous low-molecular-weight heparin compared with oral-anticoagulant therapy in the treatment of deep venous thrombosis. *Eur J Vasc Endovasc Surg* 2009;37:349-56 [[19121589](#)]

Gonzlez-Fajardo, 2008:

Gonzlez-Fajardo JA, Martin-Pedrosa M, Castrodeza J, Tamames S, Vaquero-Puerta C Effect of the anticoagulant therapy in the incidence of post-thrombotic syndrome and recurrent thromboembolism: Comparative study of enoxaparin versus coumarin. *J Vasc Surg* 2008;48:953-9 [[18639417](#)]

Kakkar, 2003:

Kakkar VV, Gebaska M, Kadziola Z, Saba N, Carrasco P Low-molecular-weight heparin in the acute and long-term treatment of deep vein thrombosis. *Thromb Haemost* 2003;89:674-80 [12669122]

Lee, 2003:

Lee AY, Levine MN, Baker RI, Bowden C, Kakkar AK, Prins M, Rickles FR, Julian JA, Haley S, Kovacs MJ, Gent M Low-molecular-weight heparin versus a coumarin for the prevention of recurrent venous thromboembolism in patients with cancer. *N Engl J Med* 2003;349:146-53 [12853587]

Lee AY, Rickles FR, Julian JA, Gent M, Baker RI, Bowden C, Kakkar AK, Prins M, Levine MN Randomized comparison of low molecular weight heparin and coumarin derivatives on the survival of patients with cancer and venous thromboembolism. *J Clin Oncol* 2005;23:2123-9 [15699480]

Das, 1996:

Das SK, Cohen AT, Edmondson RA, Melissari E, Kakkar VV Low-molecular-weight heparin versus warfarin for prevention of recurrent venous thromboembolism: a randomized trial. *World J Surg* 1996;20:521-6; discussion 526-7 [8661630]

Deitcher, 2003:

Deitcher SR, Kessler MG, Lyons RM., imag Treatment of venous thromboembolic events (VTE) in patients with active malignancy: a randomized study of enoxaparin alone versus initial enoxaparin followed by warfarin for a 180-day period [abstract].*s-e Blood* 2003; 102:322a

Meyer, 2002:

Meyer G, Marjanovic Z, Valcke J, Lorcerie B, Gruel Y, Solal-Celigny P, Le Maignan C, Extra JM, Cottu P, Farge D Comparison of low-molecular-weight heparin and warfarin for the secondary prevention of venous thromboembolism in patients with cancer: a randomized controlled study. *Arch Intern Med* 2002;162:1729-35 [12153376]

Gonzalez-Fajardo, 1999:

Gonzalez-Fajardo JA, Arreba E, Castrodeza J, Perez JL, Fernandez L, Agundez I, Mateo AM, Carrera S, Gutierrez V, Vaquero C Venographic comparison of subcutaneous low-molecular weight heparin with oral anticoagulant therapy in the long-term treatment of deep venous thrombosis. *J Vasc Surg* 1999;30:283-92 [10436448]

Gonzalez-Fajardo JA, Martin-Pedrosa M, Castrodeza J, Tamames S, Vaquero-Puerta C Effect of the anticoagulant therapy in the incidence of post-thrombotic syndrome and recurrent thromboembolism: Comparative study of enoxaparin versus coumarin. *J Vasc Surg* 2008 Oct;48:953-9 [18639417]

Pini, 1994:

Pini M, Aiello S, Manotti C, Pattacini C, Quintavalla R, Poli T, Tagliaferri A, Dettori AG Low molecular weight heparin versus warfarin in the prevention of recurrences after deep vein thrombosis. *Thromb Haemost* 1994;72:191-7 [7831650]

Hull, 2002:

Hull RD, Pineo GF, Mah AF, et al, for the LITE Study A randomized trial evaluating long-term lowmolecular- weight heparin therapy for three months versus intravenous heparin followed by warfarin sodium [abstract].*582258 Blood* 2002; 100:148a

5 home treatment

Trial	Treatments	Patients	Trials design and methods
LMWH at home vs UFH in hospital			
Koopman , 1996 n=202/198 follow-up: 12 weeks	home treatment with twice daily injections of nadroparin at a dose adjusted for patients weight; versus UH (APTT adjusted dose, continuous intravenous infusion of 1250 IU per hour after initial intravenous bolus of 5000 IU) in hospital.	patients with acute symptomatic proximal DVT proven by venography or duplex scan	Parallel groups open The Netherlands, France, Italy, New Zealand Australia

continued...

Trial	Treatments	Patients	Trials design and methods
Boccalon , 2000 n=99/101 follow-up: 6 months	home treatment with sub-cutaneous injection of LMWH (dalteparin sodium, enoxaparin sodium or nadroparin calcium as chosen by the attending physician) at the recommended dose followed by anticoagulant for 6months versus Sub-cutaneous injection of LMWH(dalteparin sodium, enoxaparin sodium or nadroparin calcium as chosen by attending physician) at the recommended dose followed by anticoagulant for 6 months initially in hospital for 10 +/- 2 days then at home	patientst with confirmed diagnosis (by ultrasonography or venography) of proximal DVT not more than 30 days before enrolment	Parallel groups NA France
Levine , 1996 n=247/253 follow-up: 90 days	home treatment by Sub-cutaneous enoxaparin 1 mg per kg body weight twice a day for at least 5 days versus UH (APTT adjusted dose, continuous intravenous infusion of 20,000 IU after initial intravenous bolus of 5000 IU) in hospital for at least 5 days	patients with acute proximal DVT proven on venography or duplex scan	Parallel groups open Canada
Ramacciotti , 2004 n=104/97 follow-up:	home treatment by once daily Subcutaneous injection of enoxaparin at a dose of 1.5 mg/kg for 5-10 days versus in hospital intravenous bolus injection of 5000 IU of UFH followed by intravenous 500 IU/kg/day adjusted to maintain an aPTT of 1.5-2.5 times the normal value for 5-10 days.	patientst with DVT symptoms for greater than or equal to 10 days and proximal lower limb DVT confirmed by duplex ultrasound or venography	Parallel groups open Brazil
Daskalopoulos , 2005 n=55/53 follow-up:	home treatment with single sub- cutaneous injection of LMWH (tinzaparin sodium) in a weight adjusted dose (175 anti Xa IU/Kg) daily for 6 months versus Intravenous bolus of 5000IU UFH followed by intravenous infusion of UFH for 5-7 days. APTT was measured after 4 hours of the initiation of heparin administration and was repeated 6 hours thereafter to reach the therapeutic range (ratio: 1.5-2.5). Oral an	patients with acute proximal DVT confirmed by colour duplex UScan not more than 1 week onset	Parallel groups open Greece

continued...

Trial	Treatments	Patients	Trials design and methods
Chong , 2005 n=150/148 follow-up: 24 months	once daily sub-cutaneous injection of enoxaparin 1.5mg/kg for a minimum of 5 days plus 10mg of warfarin for 3 months adjusted to achieve INR above 2 and within range accepted by the investigator versus 5000 IU bolus of unfractionated heparin (UFH) for a minimum of 5 days plus 10mg warfarin started on day 1 of the treatment for 3 months	patients with diagnosis of symptomatic lower extremity DVT (proximal or distal) confirmed by either contrast venography and/or ultrasonography, be suitable for treatment in an outpatient setting	Parallel groups open Australia, New Zealand, Poland, South Africa

References

Koopman, 1996:

Koopman MM, Prandoni P, Piovella F, Ockelford PA, Brandjes DP, van der Meer J, Gallus AS, Simonneau G, Chesterman CH, Prins MH Treatment of venous thrombosis with intravenous unfractionated heparin administered in the hospital as compared with subcutaneous low-molecular-weight heparin administered at home. The Tasman Study Group. N Engl J Med 1996;334:682-7 [8594426]

van den Belt AG, Bossuyt PM, Prins MH, Gallus AS, Bller HR Replacing inpatient care by outpatient care in the treatment of deep venous thrombosis—an economic evaluation. TASMAN Study Group. Thromb Haemost 1998;79:259-63 [9493572]

Boccalon, 2000:

Boccalon H, Elias A, Chal JJ, Cadne A, Gabriel S Clinical outcome and cost of hospital vs home treatment of proximal deep vein thrombosis with a low-molecular-weight heparin: the Vascular Midi-Pyrenees study. Arch Intern Med 2000;160:1769-73 [10871969]

Levine, 1996:

Levine M, Gent M, Hirsh J, Leclerc J, Anderson D, Weitz J, Ginsberg J, Turpie AG, Demers C, Kovacs M A comparison of low-molecular-weight heparin administered primarily at home with unfractionated heparin administered in the hospital for proximal deep-vein thrombosis. N Engl J Med 1996;334:677-81 [8594425]

O'Brien B, Levine M, Willan A, Goeree R, Haley S, Blackhouse G, Gent M Economic evaluation of outpatient treatment with low-molecular-weight heparin for proximal vein thrombosis. Arch Intern Med 1999;159:2298-304 [10547169]

Ramacciotti, 2004:

Ramacciotti E, Arajo GR, Lastoria S, Maffei FH, Karaoglan de Moura L, Michaelis W, Sandri JL, Dietrich-Neto F An open-label, comparative study of the efficacy and safety of once-daily dose of enoxaparin versus unfractionated heparin in the treatment of proximal lower limb deep-vein thrombosis. Thromb Res 2004;114:149-53 [15342210]

Daskalopoulos, 2005:

Daskalopoulos ME, Daskalopoulou SS, Tzortzis E, Sfridis P, Nikolaou A, Dimitroulis D, Kakissis I, Liapis CD Long-term treatment of deep venous thrombosis with a low molecular weight heparin (tinzaparin): a prospective randomized trial. Eur J Vasc Endovasc Surg 2005;29:638-50 [15878544]

Chong, 2005:

Chong BH, Brighton TA, Baker RI, Thurlow P, Lee CH Once-daily enoxaparin in the outpatient setting versus unfractionated heparin in hospital for the treatment of symptomatic deep-vein thrombosis. J Thromb Thrombolysis 2005;19:173-81 [16082604]

6 indirect anti-XA

Trial	Treatments	Patients	Trials design and methods
fondaparinux vs enoxaparin			

continued...

Trial	Treatments	Patients	Trials design and methods
MATISSE , 2004 n=1098/1107 follow-up: 3 months	fondaparinux 7.5 mg subcutaneously once daily for at least 5 days and until vitamin K antagonists induced an INR greater than 2.0. versus enoxaparin, 1 mg/kg of body weight, subcutaneously twice daily for at least 5 days and until vitamin K antagonists induced an INR greater than 2.0.	patients with acute symptomatic deep venous thrombosis	Parallel groups double blind international

References

MATISSE, 2004:

Biller HR, Davidson BL, Decousus H, Gallus A, Gent M, Piovella F, Prins MH, Raskob G, Segers AE, Cariou R, Leeuwenkamp O, Lensing AW Fondaparinux or enoxaparin for the initial treatment of symptomatic deep venous thrombosis: a randomized trial. *Ann Intern Med* 2004;140:867-73 [[15172900](#)]

7 Low molecular weight heparin

Trial	Treatments	Patients	Trials design and methods
Dalteparin vs unfractionated heparin			
Holm et al , 1986 n=29/27 follow-up: Hospital Stay	Dalteparin Subcutaneous twice daily adjusted for 7 Days, 57-107 U/kg BID versus unfractionated heparin subcutaneous twice daily 16000-30000 U	-	
Bratt et al , 1985 n=25/29 follow-up: 23 Months (mean)	Dalteparin Intravenous (adjusted) for >=5 Days, 120 U/kg BID versus unfractionated heparin intravenous APPTx1.7-3.5	-	
Bratt et al , 1990 n=60/60 follow-up: 65279;Hospital stay	Dalteparin Subcutaneous twice daily adjusted for >= 5 Days, 120 U/kg BID versus unfractionated heparin intravenous APPTx2-4	-	
Lindmarker et al , 1993 n=101/103 follow-up: 6 Months	Dalteparin Subcutaneous once daily for >= 5 Days, 200 U/kg BID versus unfractionated heparin intravenous APPTx1.5-3	-	
Enoxaparin vs unfractionated heparin			

continued...

Trial	Treatments	Patients	Trials design and methods
Simonneau et al , 1993 n=67/67 follow-up: 3 Months	Enoxaparin Subcutaneous twice daily for 0 Days, 100 U/kg BID versus unfractionated heparin intravenous APPTx1.5-2.5	-	
Minocloparine vs unfractionated heparin			
Faivre et al , 1988 n=33/37 follow-up: 10 Days	Minocloparine (CY222) Subcutaneous twice daily for 10 Days,155 U/kg BID versus unfractionated heparin subcutaneous twice daily APPTx2-3	-	
Nadroparin vs unfractionated heparin			
Collaborative European Multicentre , 1991 n=70/66 follow-up: 12 Weeks	Nadroparin Subcutaneous twice daily for 10 Days, 90 U/kg BID versus unfractionated heparin intravenous APPTx1.5-2	-	
Prandoni et al , 1992 n=85/85 follow-up: 6 Months	Nadroparin Subcutaneous twice daily for >=0 Days, 90 U/kg BID versus unfractionated heparin intravenous APPTx1.5-2	-	
Lopaciuk et al , 1992 n=74/75 follow-up: 3 Months	Nadroparin Subcutaneous twice daily for 10 Days, 92 U/kg BID versus unfractionated heparin subcutaneous twice daily APPTx1.5-2.5	-	
Tinzaparin vs unfractionated heparin			
Hull et al , 1992 n=213/219 follow-up: 3 Months	Tinzaparin Subcutaneous once daily for >= Days, 175 U/kg BID versus unfractionated heparin intravenous APPTx2-3	-	

References

Holm et al , 1986:

Bratt et al , 1985:

Bratt G, Tornebohm E, Granqvist S, Aberg W, Lockner D A comparison between low molecular weight heparin (KABI 2165) and standard heparin in the intravenous treatment of deep venous thrombosis. *Thromb Haemost* 1985 Dec 17;54:813-7 [[3911482](#)]

Bratt et al, 1990:

Bratt G, Aberg W, Johansson M, Tornebohm E, Granqvist S, Lockner D Two daily subcutaneous injections of fragmin as compared with intravenous standard heparin in the treatment of deep venous thrombosis (DVT). *Thromb Haemost* 1990 Dec 28;64:506-10 [[1964751](#)]

Lindmarker et al , 1993:

Ruan C, Gu J, Wang X, Chu X, Pan J Application of GPIIIa gene Taq I polymorphism to determination of carrier status in Glanzmann's thrombasthenia families of Chinese origin. *Thromb Haemost* 1993 Jan 11;69:64-9 [[8095357](#)]

Simonneau et al , 1993:

Simonneau G, Charbonnier B, Decousus H, Planchon B, Ninet J, Sie P, Silsiguen M, Combe S Subcutaneous low-molecular-weight heparin compared with continuous intravenous unfractionated heparin in the treatment of proximal deep vein thrombosis. Arch Intern Med 1993 Jul 12;153:1541-6 [8391792]

Faivre et al , 1988:

Faivre R, Neuhart Y, Kieffer Y, Apfel F, Magnin D, Didier D, Toulemonde F, Bassand JP, Maurat JP [A new treatment of deep venous thrombosis: low molecular weight heparin fractions. Randomized study] Presse Med 1988 Feb 13;17:197-200 [2965375]

Collaborative European Multicentre, 1991:

A randomised trial of subcutaneous low molecular weight heparin (CY 216) compared with intravenous unfractionated heparin in the treatment of deep vein thrombosis. A collaborative European multicentre study. Thromb Haemost 1991 Mar 4;65:251-6 [1646490]

Prandoni et al , 1992:

Prandoni P, Lensing AW, Buller HR, Carta M, Cogo A, Vigo M, Casara D, Ruol A, ten Cate JW Comparison of subcutaneous low-molecular-weight heparin with intravenous standard heparin in proximal deep-vein thrombosis. Lancet 1992 Feb 22;339:441-5 [1346817]

Lopaciuk et al , 1992:

Lopaciuk S, Meissner AJ, Filipecki S, Zawilska K, Sowier J, Ciesielski L, Bielawiec M, Glowinski S, Czestochowska E Subcutaneous low molecular weight heparin versus subcutaneous unfractionated heparin in the treatment of deep vein thrombosis: a Polish multicenter trial. Thromb Haemost 1992 Jul 6;68:14-8 [1325076]

Hull et al , 1992:

Hull RD, Raskob GE, Pineo GF, Green D, Trowbridge AA, Elliott CG, Lerner RG, Hall J, Sparling T, Brettell HR Subcutaneous low-molecular-weight heparin compared with continuous intravenous heparin in the treatment of proximal-vein thrombosis. N Engl J Med 1992 Apr 9;326:975-82 [1545850]

8 once daily LMWH

Trial	Treatments	Patients	Trials design and methods
once daily dalteparin vs twice daily dalteparin			
Holmstrm , 1992 n=50/51 follow-up:	once daily dalteparin 200 U (anti-FXa)/kg for at least 5 days versus twice daily dalteparin 100 U (anti-FXa)/kg for at least 5 days	Patients with a first occurrence of DVT in the lower limb, confirmed with phlebography	Parallel groups open Sweden
Partsch , 1996 n=76/64 follow-up:	Fragmin administered 200 IU/kg once daily for at least 7 days versus Fragmin 100 IU/kg twice daily for at least 7 days	patients presented with DVT extending into the iliofemoral segment diagnosed by duplex ultrasonography	Parallel groups NA Austria
once daily enoxaparin vs twice daily enoxaparin			
Merli , 2001 n=298/312 follow-up:	enoxaparin 1.5 mg/kg body weight once daily versus S.c. enoxaparin at fixed dosages of 1.0 mg/kg of body weight twice daily	patients with a symptomatic lower-extremity DVT confirmed by venography or ultrasonography (including patients with confirmed PE)	Parallel groups double blind Europe, United States of America and Australia, image/pj
once daily logiparin vs twice daily logiparin			

continued...

Trial	Treatments	Patients	Trials design and methods
Siegbahn , 1989 n=10/10 follow-up:	Once daily logiparin 150 XaI U/kgp, imag versus twice daily logiparin 75 XaI U/kg	patients with a venographically confirmed episode of DVT	Parallel groups single blind Sweden and Denmark
once daily nadroparin vs twice daily nadroparin			
Charbonnier , 1998 n=316/335 follow-up:	Once daily nadroparin 20,500 (AXa IU/ml)continued for at least 5 days versus twice daily nadroparin 10,250 (AXa IU/ml)continued for at least 5 days	patients with acute symptomatic proximal DVT in popliteal vein or above documented by venography	Parallel groups double blind Europe
once daily enoxaparin vs UFH			
Merli (once daily vs UFH) , 2001 n=298/290 follow-up: 3 months	Initial therapy with enoxaparin 1.5 mg/kg body weight once daily versus Initial therapy with dose-adjusted intravenous unfractionated heparin	patients with a symptomatic lower-extremity DVT confirmed by venography or ultrasonography (including patients with confirmed PE)	Parallel groups partialy blinded Europe, United States of America and Australia, image/pj

References

Holmstrm, 1992:

Holmostrm M, Berglund MC, Granquist S, Bratt G, Trnebohm E, Lockner D Fragmin once or twice daily subcutaneously in the treatment of deep venous thrombosis of the leg. *Thromb Res* 1992;67:49-55 [[1332213](#)]

Partsch, 1996:

Partsch H, Kechavarz B, Mostbeck A, Khn H, Lipp C Frequency of pulmonary embolism in patients who have iliofemoral deep vein thrombosis and are treated with once- or twice-daily low-molecular-weight heparin. *J Vasc Surg* 1996;24:774-82 [[8918323](#)]

Merli, 2001:

Merli G, Spiro TE, Olsson CG, Abildgaard U, Davidson BL, Eldor A, Elias D, Grigg A, Musset D, Rodgers GM, Trowbridge AA, Yusen RD, Zawilska K Subcutaneous enoxaparin once or twice daily compared with intravenous unfractionated heparin for treatment of venous thromboembolic disease. *Ann Intern Med* 2001;134:191-202 [[11177331](#)]

Siegbahn, 1989:

Siegbahn A, Y-Hassan S, Boberg J, Bylund H, Neerstrand HS, Ostergaard P, Hedner U Subcutaneous treatment of deep venous thrombosis with low molecular weight heparin. A dose finding study with LMWH-Novo. *Thromb Res* 1989;55:767-78 [[2551070](#)]

Charbonnier, 1998:

Charbonnier BA, Fiessinger JN, Banga JD, Wenzel E, d'Azemar P, Sagnard L Comparison of a once daily with a twice daily subcutaneous low molecular weight heparin regimen in the treatment of deep vein thrombosis. FRAXODI group. *Thromb Haemost* 1998;79:897-901 [[9609216](#)]

Merli (once daily vs UFH), 2001:

Merli G, Spiro TE, Olsson CG, Abildgaard U, Davidson BL, Eldor A, Elias D, Grigg A, Musset D, Rodgers GM, Trowbridge AA, Yusen RD, Zawilska K Subcutaneous enoxaparin once or twice daily compared with intravenous unfractionated heparin for treatment of venous thromboembolic disease. *Ann Intern Med* 2001 Feb 6;134:191-202 [[11177331](#)]

9 oral direct thrombin inhibitor

Trial	Treatments	Patients	Trials design and methods
ximelagatran vs discontinuation			
THRIVE III , 2003 n=612/611 follow-up: 18 months	ximelagatran 24 mg twice daily for 18 months versus placebo for 18 months	patients with venous thromboembolism who had undergone six months of anticoagulant therapy	Parallel groups double blind 18 countries
heparin/dabigatran vs heparin/VKA			
RE-COVER , 2009 [NCT00291330] n=1274/1265 follow-up: 6 months	dabigatran 150 mg twice daily in a fixed-dose versus warfarin dose-adjusted to an INR between 2.0 and 3.0	patients with acute venous thromboembolism , treated with low molecular weight or unfractionated heparin for 5 to 11 days	Parallel groups double blind

References

THRIVE III, 2003:

Schulman S, Whlander K, Lundstrm T, Clason SB, Eriksson H Secondary prevention of venous thromboembolism with the oral direct thrombin inhibitor ximelagatran. N Engl J Med 2003 Oct 30;349:1713-21 [[14585939](#)]

Schulman S, Lundstrm T, Wlander K, Billing Clason S, Eriksson H Ximelagatran for the secondary prevention of venous thromboembolism: a complementary follow-up analysis of the THRIVE III study. Thromb Haemost 2005 Oct;94:820-4 [[16270637](#)]

RE-COVER, 2009:

Schulman S, Eriksson H, Goldhaber SZ, et al. Dabigatran etexilate versus warfarin in the treatment of venous thromboembolism. Oral presentation, ASH Annual Meeting 2009, Abstract number 19394, 6 December 2009

Schulman S, Kearon C, Kakkar AK, Mismetti P, Schellong S, Eriksson H, Baanstra D, Schnee J, Goldhaber SZ Dabigatran versus warfarin in the treatment of acute venous thromboembolism. N Engl J Med 2009 Dec 10;361:2342-52 [[19966341](#)] [10.1056/NEJMoa0906598](#)

10 prolonged oral VKA

Trial	Treatments	Patients	Trials design and methods
6 months vs 1.5 months			
Schulman , 1995 n=NA follow-up: Two years after randomization	6 months treatment with warfarin or dicoumarol adjusted for a target INR between 2.0 - 2.85 versus 1.5 months warfarin or dicoumarol adjusted for a target INR between 2.0 - 2.85	-	Parallel groups open Sweden
3-6 months vs 1.5-3 months			

continued...

Trial	Treatments	Patients	Trials design and methods
Pinede , 2001 n=NA follow-up: 15 months after randomization	Long course of therapy (6 months for proximal DVT and/or PE; 12 weeks for calf DVT) by fluindione adjusted for INR range of 2.0 to 3.0 versus Short oral anticoagulant course (3 months for proximal DVT and/or PE; 6 weeks for isolated calf DVT) by fluindione adjusted for INR range of 2.0 to 3.0	-	Parallel groups open France

References

Schulman, 1995:

Schulman S, Rhedin AS, Lindmarker P, Carlsson A, Lrfars G, Nicol P, Loogna E, Svensson E, Ljungberg B, Walter H A comparison of six weeks with six months of oral anticoagulant therapy after a first episode of venous thromboembolism. Duration of Anticoagulation Trial Study Group. N Engl J Med 1995;332:1661-5 [7760866]

Pinede, 2001:

Pinede L, Ninet J, Duhaut P, Chabaud S, Demolombe-Rague S, Durieu I, Nony P, Sanson C, Boissel JP Comparison of 3 and 6 months of oral anticoagulant therapy after a first episode of proximal deep vein thrombosis or pulmonary embolism and comparison of 6 and 12 weeks of therapy after isolated calf deep vein thrombosis. Circulation 2001;103:2453-60 [11369685]

11 subcutaneous heparin

Trial	Treatments	Patients	Trials design and methods
subcutaneous heparin vs intravenous heparin			
Krahenbuhl , 1979 n=23/25	subcutaneous sodic heparin 30 000 U daily (mean) versus intravenous sodic heparin 30 000 U daily (mean)	-	
Bentley , 1980 n=50/50	subcutaneous calcic heparin 37 000 U daily (mean) versus intravenous sodic heparin 36 800 U daily (mean)	-	
Andersson , 1982 n=72/69	subcutaneous sodic heparin 36 800 U daily (mean) versus intravenous sodic heparin 33 250 U daily (mean)	-	

continued...

Trial	Treatments	Patients	Trials design and methods
Hull , 1986 n=57/58	subcutaneous sodic heparin 32 300 U daily (mean) versus intravenous sodic heparin 29 700 U daily (mean)	-	
Doyle , 1987 n=51/52	subcutaneous calcic heparin 29 200 U daily (mean) versus intravenous calcic heparin 29 600 U daily (mean)	-	
Walker , 1987 n=50/50	subcutaneous calcic heparin 29 375 U daily (mean) versus intravenous calcic heparin 24 384 U daily (mean)	-	
Lopaciuk , 3000 n=48/46	subcutaneous sodic heparin 34 400 U daily (mean) versus intravenous sodic heparin 37 000 U daily (mean)	-	
Pini , 1990 n=138/133	subcutaneous calcic heparin 33 800 U daily (mean) versus intravenous sodic heparin 31 700 U daily (mean)	-	

References

Krahenbuhl, 1979:

Krahenbuhl B, Simon CA, Bouvier CA, Schinas P, Hopf MA, Cochet B [Heparin treatment. Comparison between intravenous and subcutaneous administration] Schweiz Med Wochenschr 1979;109:1322-5 [504973]

Bentley, 1980:

Bentley PG, Kakkar VV, Scully MF, MacGregor IR, Webb P, Chan P, Jones N An objective study of alternative methods of heparin administration. Thromb Res 1980;18:177-87 [7404497]

Andersson, 1982:

Andersson G, Fagrell B, Holmgren K, Johnsson H, Ljungberg B, Nilsson E, Wilhelmsson S, Zetterquist S Subcutaneous administration of heparin. A randomised comparison with intravenous administration of heparin to patients with deep-vein thrombosis. Thromb Res 1982;27:631-9 [7179208]

Hull, 1986:

Hull RD, Raskob GE, Hirsh J, Jay RM, Leclerc JR, Geerts WH, Rosenbloom D, Sackett DL, Anderson C, Harrison L Continuous intravenous heparin compared with intermittent subcutaneous heparin in the initial treatment of proximal-vein thrombosis. N Engl J Med 1986;315:1109-14 [3531862]

Doyle, 1987:

Doyle DJ, Turpie AG, Hirsh J, Best C, Kinch D, Levine MN, Gent M Adjusted subcutaneous heparin or continuous intravenous heparin in patients with acute deep vein thrombosis. A randomized trial. Ann Intern Med 1987;107:441-5 [3307582]

Walker, 1987:

Walker MG, Shaw JW, Thomson GJ, Cumming JG, Thomas ML Subcutaneous calcium heparin versus intravenous sodium heparin in treatment of established acute deep vein thrombosis of the legs: a multicentre prospective randomised trial. Br Med J (Clin Res Ed) 1987;294:1189-92 [3109574]

Lopaciuk, 3000:

Pini, 1990:

Pini M, Pattachini C, Quintavalla R, Poli T, Megha A, Tagliaferri A, Manotti C, Dettori AG Subcutaneous vs intravenous heparin in the treatment of deep venous thrombosis—a randomized clinical trial. Thromb Haemost 1990;64:222-6 [2270531]

12 synthetic oligosaccharide

Trial	Treatments	Patients	Trials design and methods
fondaparinux vs heparin			
MATISSE PE , 2003 n=1103/1110 follow-up: 3 mo	fondaparinux subcutaneously once daily versus continuous intravenous infusion of unfractionated heparin	patients with acute symptomatic pulmonary embolism	Parallel groups open
idraparinux (without heparin) vs heparin/VKA			
VanGogh DVT , 2007 [NCT00067093] n=1452/1452 follow-up: 3 mo (6 mo)	subcutaneous idraparinux (2.5 mg once weekly) versus heparin followed by an adjusted-dose vitamin K antagonist	patients with deep-vein thrombosis	Parallel groups open
VanGogh PE , 2007 [NCT00062803] n=1095/1120 follow-up: 3 mo (6 mo)	subcutaneous idraparinux (2.5 mg once weekly) versus heparin followed by an adjusted-dose vitamin K antagonist	patients with pulmonary embolism	Parallel groups open

References

MATISSE PE, 2003:

Biller HR, Davidson BL, Decousus H, Gallus A, Gent M, Piovella F, Prins MH, Raskob G, van den Berg-Segers AE, Cariou R, Leeuwenkamp O, Lensing AW Subcutaneous fondaparinux versus intravenous unfractionated heparin in the initial treatment of pulmonary embolism. N Engl J Med 2003 Oct 30;349:1695-702 [14585937]

VanGogh DVT, 2007:

VanGogh PE, 2007:

13 synthetic pentasaccharide

Trial	Treatments	Patients	Trials design and methods
idraparinux vs discontinuation			

continued...

Trial	Treatments	Patients	Trials design and methods
VanGogh extension , 2007 [NCT00071279] n=594/621 follow-up: 6 months	once-weekly injections of 2.5 mg of idraparinux for 6 months versus placebo	patients who had completed 6 months of prophylaxis with idraparinux or a vitamin K antagonist and in whom extended anticoagulation was warranted	Parallel groups

References

VanGogh extension, 2007:

14 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.