

# Clinical trials of cholesterol lowering intervention for cardiovascular prevention in patients with prior MI or with CHD

TrialResults-center [www.trialresultscenter.org](http://www.trialresultscenter.org)

## 1 diet

Trial	Treatments	Patients	Trials design and methods
<b>diet vs usual diet</b>			
<b>Kallio , 1979</b> n=188/187 follow-up: 3.0 years	diet (multifactorial intervention programme) versus usual diet	patients below 65 years who had an acute myocardial infarction	Parallel groups open
<b>Los Angeles VA (Dayton) , 1969</b> n=424/422 follow-up: 65279;8.0 y	diet versus usual diet	men in domiciliary care, age>55, with or without CHD	Parallel groups double blind USA
<b>Ornish , 1990</b> n=28/20 follow-up: 1.0 y	low-fat vegetarian diet, stopping smoking, stress management training, and moderate exercise versus usual-care	Patients with angiographically documented coronary artery disease	Parallel groups open USA
<b>Rose , 1965</b> n=28/26 follow-up: 1.2 years	Rgime restreint en graisses + 80 g/j huile de mas versus usual diet	men, <70 years	Parallel groups open
<b>Singh , 1992</b> n=204/202 follow-up: 65279;2.0 years	strict diet versus usual diet	patients with suspected acute myocardial infarction	Parallel groups open
<b>STARS (St Thomas, diet) , 1992</b> n=30/30 follow-up: 3.0 years	dietary advice versus usual diet	patients with angina or past myocardial infarction	open, blind assessment
<b>low fat diet vs mediterranean-style diet</b>			
<b>Tuttle , 2008</b> n=NA follow-up: 24 months	low-fat versus Mediterranean-style diets	First MI survivors	Parallel groups open

## References

### Kallio, 1979:

Kallio V, Hmlinen H, Hakkila J, Luurila OJ Reduction in sudden deaths by a multifactorial intervention programme after acute myocardial infarction. Lancet 1979;2:1091-4 [91836]

### Los Angeles VA (Dayton), 1969:

Rogers MC Sir John Scott Burdon-Sanderson (1828-1905): a pioneer in electrophysiology. Circulation 1969;40:1-2 [4893441]

Dayton S, Pearce ML Diet high in unsaturated fat. A controlled clinical trial. Minn Med 1969;52:1237-42 [4896402]

### Ornish, 1990:

Ornish D, Brown SE, Scherwitz LW, Billings JH, Armstrong WT, Ports TA, McLanahan SM, Kirkeeide RL, Brand RJ, Gould KL Can lifestyle changes reverse coronary heart disease? The Lifestyle Heart Trial. Lancet 1990;336:129-33 [1973470]

### Rose, 1965:

ROSE GA, THOMSON WB, WILLIAMS RT CORN OIL IN TREATMENT OF ISCHAEMIC HEART DISEASE. Br Med J 1965 Jun 12;1:1531-3 [14288105]

### Singh, 1992:

Singh RB, Rastogi SS, Verma R, Laxmi B, Singh R, Ghosh S, Niaz MA Randomised controlled trial of cardioprotective diet in patients with recent acute myocardial infarction: results of one year follow up. BMJ 1992;304:1015-9 [1586782]

### STARS (St Thomas, diet), 1992:

Watts GF, Lewis B, Brunt JN, Lewis ES, Coltart DJ, Smith LD, Mann JI, Swan AV Effects on coronary artery disease of lipid-lowering diet, or diet plus cholestyramine, in the St Thomas' Atherosclerosis Regression Study (STARS) Lancet 1992;339:563-9 [1347091]

### Tuttle, 2008:

Tuttle KR, Shuler LA, Packard DP, Milton JE, Daratha KB, Bibus DM, Short RA Comparison of low-fat versus Mediterranean-style dietary intervention after first myocardial infarction (from The Heart Institute of Spokane Diet Intervention and Evaluation Trial). Am J Cardiol 2008;101:1523-30 [18489927]

## 2 ezetimibe

Trial	Treatments	Patients	Trials design and methods
<b>ezetimibe vs placebo (on top statins)</b>			
<b>IMPROVE-IT , 2014</b> [NCT00202878] n=9067/9077 follow-up: 5.68 years	10 mg/day of ezetimibe and 40 mg/day of simvastatin versus simvastatin 40 mg/day	subjects with stabilized high-risk acute coronary syndrome	Parallel groups double blind 39 countries

## References

### IMPROVE-IT, 2014:

Cannon CP, Blazing MA, Giugliano RP, McCagg A, White JA, Theroux P, Darius H, Lewis BS, Ophuis TO, Jukema JW, De Ferrari GM, Ruzyllo W, De Lucca P, Im K, Bohula EA, Reist C, Wiviott SD, Tershakovec AM, Musliner TA, Braunwald E, Califf RM Ezetimibe Added to Statin Therapy after Acute Coronary Syndromes. N Engl J Med 2015;372:2387-97 [26039521]

## 3 fibrates

Trial	Treatments	Patients	Trials design and methods
<b>bezafibrate vs placebo</b>			
<b>SENDCAP , 1998</b> n=81/83 follow-up: 3.0 years	bezafibrate 400 mg daily versus placebo	type 2 diabetic subjects without a history of clinical cardiovascular	Parallel groups double blind UK

continued...

Trial	Treatments	Patients	Trials design and methods
<b>gemfibrozil vs placebo</b>			
HHS (Frick)(secondary prev subgroup) , 1993 n=311/317 follow-up: 5.0 years	gemfibrozil 600 mg twice daily versus placebo	individuals who exhibited symptoms and signs of possible coronary heart disease	Parallel groups double blind Sweden
LOCAT , 1997 n=197/198 follow-up: 32 months	gemfibrozil 1200 mg/d versus placebo	post-coronary bypass men, who had an HDL cholesterol concentration $\leq 1.1$ mmol/L and LDL cholesterol $\leq 4.5$ mmol/L	Parallel groups double blind Germany
VA-HIT , 1999 [NCT00283335] n=1264/1267 follow-up: 5.1 years	gemfibrozil 1.2g daily versus placebo	men with coronary heart disease, an HDL cholesterol level of 40 mg per deciliter (1.0 mmol per liter) or less, and an LDL cholesterol level of 140 mg per deciliter (3.6 mmol per liter) or less	Parallel groups double blind USA

## References

### SENDCAP, 1998:

Elkeles RS, Diamond JR, Poulter C, Dhanjil S, Nicolaidis AN, Mahmood S, Richmond W, Mather H, Sharp P, Feher MD Cardiovascular outcomes in type 2 diabetes. A double-blind placebo-controlled study of bezafibrate: the St. Mary's, Ealing, Northwick Park Diabetes Cardiovascular Disease Prevention (SENDCAP) Study. *Diabetes Care* 1998;21:641-8 [9571357]

### HHS (Frick)(secondary prev subgroup), 1993:

Frick MH, Heinonen OP, Huttunen JK, Koskinen P, Mnttri M, Manninen V Efficacy of gemfibrozil in dyslipidaemic subjects with suspected heart disease. An ancillary study in the Helsinki Heart Study frame population. *Ann Med* 1993;25:41-5 [8435186]

### LOCAT, 1997:

Frick MH, Syvne M, Nieminen MS, Kauma H, Majahalme S, Virtanen V, Kesniemi YA, Pasternack A, Taskinen MR Prevention of the angiographic progression of coronary and vein-graft atherosclerosis by gemfibrozil after coronary bypass surgery in men with low levels of HDL cholesterol. Lipid Coronary Angiography Trial (LOCAT) Study Group. *Circulation* 1997;96:2137-43 [9337181]

### VA-HIT, 1999:

Rubins HB, Robins SJ, Collins D, Fye CL, Anderson JW, Elam MB, Faas FH, Linares E, Schaefer EJ, Schectman G, Wilt TJ, Wittes J, Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol. Veterans Affairs High-Density Lipoprotein Cholesterol Intervention Trial Study Group. *N Engl J Med* 1999; 341:410-8 [10438259]

Adabag AS, Mithani S, Al Aloul B, Collins D, Bertog S, Bloomfield HE Efficacy of gemfibrozil in the primary prevention of atrial fibrillation in a large randomized controlled trial. *Am Heart J* 2009 May;157:913-8 [19376321]

## 4 hormones

Trial	Treatments	Patients	Trials design and methods
<b>estrogen vs placebo</b>			
CDP estrogen 2.5 , 1975 n=1101/2789 follow-up: 4.7 years	estrogen 2.5 mg daily versus placebo	-	Parallel groups

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>CDP estrogen 5 , 1975</b> n=1119/2788 follow-up: 1.5 years	estrogen 5.0 mg daily versus placebo	-	Parallel groups
<b>Marmorstein , 1962</b> n=285/147 follow-up: 5.0 y	estrogen versus placebo	-	Parallel groups
<b>Stamler , 1963</b> n=156/119 follow-up: 5.0 years	estrogen versus placebo	-	Parallel groups
<b>estrogen or thyroxine vs placebo</b>			
<b>VA drugs (Estrogen or thyroxine) , 1968</b> n=427/143 follow-up: 65279;3.2 years	estrogen or thyroxine versus placebo	-	Parallel groups
<b>thyroxine vs placebo</b>			
<b>CDP tyroxine , 1975</b> n=1083/2715 follow-up: 3.0 years	thyroxine versus placebo	-	Parallel groups

## References

### CDP estrogen 2.5, 1975:

The Coronary Drug Project. Findings leading to discontinuation of the 2.5-mg day estrogen group. The coronary Drug Project Research Group. JAMA 1973;226:652-7 [4356847]

### CDP estrogen 5, 1975:

The Coronary Drug Project. Initial findings leading to modifications of its research protocol. JAMA 1970;214:1303-13 [4320008]

### Marmorstein, 1962:

MARMORSTON J, MOORE FJ, HOPKINS CE, KUZMA OT, WEINER J Clinical studies of long-term estrogen therapy in men with mvocardial infarction. Proc Soc Exp Biol Med 1962;110:400-8 [14470097]

### Stamler, 1963:

SCHWARTZ SO, GREENSPAN I, BROWN ER LEUKEMIA CLUSTER IN NILES, ILL. IMMUNOLOGIC DATA ON FAMILIES OF LEUKEMIC PATIENTS AND OTHERS. JAMA 1963;186:106-8 [14056521]

STAMLER J, PICK R, KATZ LN, PICK A, KAPLAN BM, BERKSON DM, CENTURY D Effectiveness of estrogens for therapy of myocardial infarction in middle-age men. JAMA 1963;183:632-8 [13983325]

### VA drugs (Estrogen or thyroxine), 1968:

### CDP tyroxine, 1975:

The coronary drug project. Findings leading to further modifications of its protocol with respect to dextrothyroxine. The coronary drug project research group. JAMA 1972;220:996-1008 [4337170]

## 5 inhibitor of lipoprotein-associated phospholipase

Trial	Treatments	Patients	Trials design and methods
<b>darapladib vs placebo</b>			
SOLID-TIMI 52 [NCT01000727] n=NA follow-up:	-	-	

## References

### SOLID-TIMI 52, :

O'Donoghue ML, Braunwald E, White HD, Steen DP, Lukas MA, Tarka E, Steg PG, Hochman JS, Bode C, Maggioni AP, Im K, Shannon JB, Davies RY, Murphy SA, Crugnale SE, Wiviott SD, Bonaca MP, Watson DF, Weaver WD, Serruys PW, Cannon CP Effect of Darapladib on Major Coronary Events After an Acute Coronary Syndrome: The SOLID-TIMI 52 Randomized Clinical Trial. JAMA 2014 Aug 31;: [25173516] [10.1001/jama.2014.11061](https://doi.org/10.1001/jama.2014.11061)

## 6 niacin

Trial	Treatments	Patients	Trials design and methods
<b>niacin vs control</b>			
VA drugs , 1968 n=77/143 follow-up: 3.2 years	-	-	Parallel groups double blind
<b>niacin vs placebo</b>			
CDP niacin , 1975 n=1119/2789 follow-up: 6.2 years	niacin 3 mg/d versus placebo	Hommes, de 30 64 ans	Parallel groups double blind

## References

### VA drugs, 1968:

Schoch HK. The US Veterans Administration Cardiology drug lipid study: an interim report Adv Exp Med Biol. 1968;4:405-420

### CDP niacin, 1975:

, Clofibrate and niacin in coronary heart disease. JAMA 1975; 231:360-81 [1088963]

## 7 other cholesterol lowering drugs

Trial	Treatments	Patients	Trials design and methods
<b>colestipol-niacin vs placebo</b>			
CLAS , 1987 n=NA follow-up: 65279;2 years	colestipol + niacin versus placebo	nonsmoking men aged 40 to 59 years with previous coronary bypass surgery	Parallel groups double blind

continued...

Trial	Treatments	Patients	Trials design and methods
<b>various drugs vs placebo</b>			
<b>HARP , 1994</b> [NCT00000461] n=40/39 follow-up: 2.5 years	Various drugs (pravastatin, nicotinic acid, cholestyramine, and gemfibrozil stepwise as needed to reach the specified goal (total cholesterol <or = 4.1 mmol/L, ratio of LDL/high-density-lipoprotein [HDL] cholesterol <or = 2.0) versus placebo	normocholesterolaemic patients with coronary heart disease	Parallel groups open
<b>various drugs vs usual care</b>			
<b>SCRIP , 1994</b> [NCT00000508] n=145/155 follow-up: 4.0 years	multifactor risk reduction (Various drugs) versus usual care	patients with angiographically defined coronary atherosclerosis	Parallel groups open

## References

### CLAS, 1987:

Blankenhorn DH, Johnson RL, Nessim SA, Azen SP, Sanmarco ME, Selzer RH The Cholesterol Lowering Atherosclerosis Study (CLAS): design, methods, and baseline results. *Control Clin Trials* 1987;8:356-87 [[3327654](#)]

Cashin-Hemphill L, Mack WJ, Pogoda JM, Sanmarco ME, Azen SP, Blankenhorn DH Beneficial effects of colestipol-niacin on coronary atherosclerosis. A 4-year follow-up. *JAMA* 1990;264:3013-7 [[2243429](#)]

Blankenhorn DH, Nessim SA, Johnson RL, Sanmarco ME, Azen SP, Cashin-Hemphill L Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts. *JAMA* 1987;257:3233-40 [[3295315](#)]

Blankenhorn DH, Azen SP, Crawford DW, Nessim SA, Sanmarco ME, Selzer RH, Shircore AM, Wickham EC Effects of colestipol-niacin therapy on human femoral atherosclerosis. *Circulation* 1991;83:438-47 [[1991366](#)]

### HARP, 1994:

Sacks FM, Pasternak RC, Gibson CM, Rosner B, Stone PH Effect on coronary atherosclerosis of decrease in plasma cholesterol concentrations in normocholesterolaemic patients. Harvard Atherosclerosis Reversibility Project (HARP) Group. *Lancet* 1994;344:1182-6 [[7934538](#)]

### SCRIP, 1994:

Haskell WL, Alderman EL, Fair JM, Maron DJ, Mackey SF, Superko HR, Williams PT, Johnstone IM, Champagne MA, Krauss RM Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and women with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994;89:975-90 [[8124838](#)]

## 8 Probuocol

Trial	Treatments	Patients	Trials design and methods
<b>Probuocol vs placebo</b>			
<b>McCaughan , 1981</b> n=88/30 follow-up: 1 year	probuocol versus placebo	hypercholesterolemic men	Parallel groups double-blind

continued...

Trial	Treatments	Patients	Trials design and methods
<b>Tardif , 1997</b> n=160/157 follow-up: 0.5 years	probucol 500 mg versus placebo	patients undergoing PTCA	Parallel groups open

## References

### McCaughan, 1981:

McCaughan D The long-term effects of probucol on serum lipid levels. Arch Intern Med 1981;141:1428-32 [7025778]

### Tardif, 1997:

Tardif JC, Ct G, Lesprance J, Bourassa M, Lambert J, Doucet S, Bilodeau L, Nattel S, de Guise P Probuclol and multivitamins in the prevention of restenosis after coronary angioplasty. Multivitamins and Probuclol Study Group. N Engl J Med 1997;337:365-72 [9241125]

## 9 resins

Trial	Treatments	Patients	Trials design and methods
<b>cholestyramine vs control</b>			
<b>STARS (cholestyramine) , 1992</b> n=30/30 follow-up: 3 years	cholestyramine versus diet	patients with angina or past myocardial infarction	
<b>colestipol vs placebo</b>			
<b>Gross , 1973</b> n=23/29 follow-up: 65279;1.0 years	colestipol versus placebo		Parallel groups

## References

### STARS (cholestyramine), 1992:

Watts GF, Lewis B, Brunt JN, Lewis ES, Coltart DJ, Smith LD, Mann JI, Swan AV Effects on coronary artery disease of lipid-lowering diet, or diet plus cholestyramine, in the St Thomas' Atherosclerosis Regression Study (STARS) Lancet 1992;339:563-9 [1347091]

### Gross, 1973:

Gross L, Figueredo R Long-term cholesterol-lowering effect of colestipol resin in humans. J Am Geriatr Soc 1973;21:552-6 [4584170]

## 10 statins

Trial	Treatments	Patients	Trials design and methods
<b>any statin vs no statin</b>			

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>Sakamoto , 2006</b> n=241/245 follow-up: up to 24 months	any available statin versus no statin	Japanese patients with AMI within 96 hours of AMI onset	Parallel groups open Japan
<b>atorvastatin vs placebo</b>			
<b>MIRACL , 2001</b> n=1538/1548 follow-up: 1 and 4 months	Atorvastatin, 80 mg (early initiation) versus Placebo	unstable angina or nonQ-wave acute MI	Parallel groups Double blind Europe, North America, South Africa, and Australasia
<b>macin , 2005</b> n=NA follow-up: 30 days	atorvastatin 40 mg daily for 30 days versus placebo	patients admitted within 48 hours of onset of ACS with CRP levels $\geq$ 1.4 mg/dL	Parallel groups double-blind
<b>fluvastatin vs placebo</b>			
<b>LIPS (sub groups) , 2002</b> n=417/407 follow-up: 1, 4, and 6 months	Fluvastatin, 80 mg versus Placebo	patients with unstable angina and successful first percutaneous coronary intervention	Parallel groups double blind Europe, Canada, and Brazil
<b>FLORIDA , 2002</b> n=265/275 follow-up: 1, 4, and 6 months	Fluvastatin, 80 mg (early initiation) versus Placebo	patients with an AMI and total cholesterol of <6.5 mmol.l	Parallel groups double blind The Netherlands
<b>FLARE , 1999</b> n=409/425 follow-up: 40 weeks	fluvastatin 40 mg twice daily versus placebo	successful coronary balloon angioplasty	Parallel groups double blind
<b>LCAS , 1997</b> n=164/157 follow-up: 2.5 years	fluvastatin 20 mg twice daily versus placebo	men and women aged 35 to 75 years with angiographic CHD and mean low-density lipoprotein (LDL) cholesterol of 115 to 190 mg/dl despite diet	Parallel groups double-blind
<b>Riegger et al. , 1999</b> n=187/178 follow-up: 1.0 years	fluvastatin 40 mg (o.a.d. or b.i.d.) versus placebo	hyperlipidaemic patients with symptomatic, clinically-diagnosed (exercise-ECG) coronary heart disease	Parallel groups double blind
<b>Czech trial ongoing</b> [NCT00171275] n=NA follow-up: 52 weeks	fluvastatin versus placebo	-	Parallel groups double blind
<b>pravastatin vs placebo</b>			
<b>LAMIL , 1997</b> n=36/33 follow-up: 1 and 3 months	Pravastatin, 10-20 mg (starting at D3) versus Placebo	patients suffering an acute myocardial infarction	Parallel groups double blind Belgium
<b>RECIFE , 1999</b> n=30/30 follow-up: 1.5 months	Pravastatin, 40 mg versus Placebo	Patients with acute myocardial infarction or unstable angina and total cholesterol levels at admission $\geq$ 5.2 mmol/L or LDL $\geq$ 3.4 mmol/L	Parallel groups double blind Canada

continued...



<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>PAIS , 2001</b> n=50/49 follow-up: 1 and 3 months	Pravastatin, 40 mg (initiated within 48 hours of hospital admission) versus Placebo	patients with acute coronary syndromes	Parallel groups double blind The Netherlands
<b>PACT , 2004</b> n=1710/1698 follow-up: 1 months	Pravastatin, 20-40 mg within 24 hours of the onset of symptoms in versus Placebo	patients with unstable angina, non-ST-segment elevation myocardial infarction, or ST-segment elevation myocardial infarction within 24 hours of the onset of symptoms	Parallel groups double blind Australia
<b>CARE , 1996</b> n=2081/2078 follow-up: 5 years	pravastatin 40 mg/d versus placebo	men and women with myocardial infarction who had plasma totalcholesterol levels below 240 mg per deciliter (mean,209) and low-density lipoprotein (LDL) cholesterollevels of 115 to 174 mg per deciliter	Parallel groups double blind USA, Canada
<b>LIPID , 1998</b> n=4512/4502 follow-up: 6.1 years	pravastatin 40 mg/d versus placebo	patients with previous myocardial infarction or unstable angina and a baseline plasma cholesterol concentration of 4.0-7.0 mmol/L	Parallel groups double blind Australie et Nouvelle Zlande
<b>simvastatin vs placebo</b>			
<b>4S , 1994</b> n=2221/2223 follow-up: 5.4 years	simvastatin 20 or 40 mg/d, target CT between 3 et 5.2 mmol/l versus placebo	patients with angina pectoris or previous myocardial infarction and serum cholesterol 5.5-8.0 mmol/L on a lipid-lowering diet	Parallel groups double blind Scandinavia
<b>A to Z , 2004</b> n=2265/2232 follow-up: 4 months	Simvastatin, 40-80 mg early initiation versus Placebo	patient with an acute coronary syndrome (ACS)	Parallel groups Double aveugle 41 countries
<b>Ren , 2009</b> n=NA follow-up:	simvastatin (40 mg/d for 4 weeks) versus placebo	patients with unstable angina pectoris	Parallel groups double-blind
<b>CIS , 1997</b> n=129/125 follow-up: 2.3 years	simvastatin 40 mg versus placebo	men with documented coronary artery disease and hypercholesterolaemia	Parallel groups double blind
<b>HPS , 2002</b> n=10269/10267 follow-up: 5 years	simvastatin 40 mg/d versus placebo	adults (aged 40-80 years) with coronary disease, other occlusive arterial disease, or diabete	Factorial plan double blind UK
<b>atorvastatin vs usual care</b>			
<b>Colivicchi , 2002</b> n=40/41 follow-up: 1, 3, and 6 months	Atorvastatin, 80 mg daily early initiation versus Usual care	unstable angina pectoris or non-Q-wave myocardial infarction	Parallel groups open Italy
<b>ESTABLISH , 2004</b> n=35/35 follow-up: 1, 4, and 6 months	Atorvastatin, 20 mg early initiation versus Usual care	patients with ACS undergoing emergency coronary angiography and percutaneous coronary intervention	Parallel groups open Japan

continued...

<b>Trial</b>	<b>Treatments</b>	<b>Patients</b>	<b>Trials design and methods</b>
<b>GREACE , 2002</b> n=800/800 follow-up: 3 years mean	atorvastatin 10-80 mg/d versus usual care	patients with established coronary heart disease	Parallel groups open
<b>lovastatin vs usual care</b>			
<b>CLAPT , 1999</b> n=112/114 follow-up: 2.0 years	lovastatin begun at 20 mg daily and tritrated up to 80 mg daily versus usual care	patients undergoing PTCA	Parallel groups open (blind assesment)
<b>Sahni , 1991</b> n=79/78 follow-up: 2 years	lovastatin 20-40mg/d versus conventional therapy alone	patients undergoing successful PTCA	Parallel groups open
<b>pravastatin vs usual care</b>			
<b>L-CAD , 2000</b> n=70/56 follow-up: 1, 4, and 6 months	Pravastatin, 20-40 mg (strating on average at D6) versus Usual care	patients with acute coronary syndrome	Parallel groups open Germany
<b>PTT , 2002</b> n=79/85 follow-up: 1 and 6 months	Pravastatin, 40 mg versus Usual care	patients who underwent coronary balloon angioplasty of the infarct-related artery during the first month of acute myocardial infarction	Parallel groups open Turkey
<b>OACIS-LIPID , 2008</b> n=176/177 follow-up: 9 months	pravastatin 10 mg/daily versus no pravastatin	patients with AMI who had plasma total cholesterol levels of 200-250 mg/dl and triglyceride levels <300 mg/dl	Parallel groups open
<b>GISSI Prevenzione , 2000</b> n=2138/2133 follow-up: 23 months (mean)	low-dose pravastatin regimen 20 mg daily versus control	recent acute myocardial infarction patients (<= 6 months) with total blood cholesterol >= 200 mg/dl and <250 mg/dl and after a period of 36 months showed plasma cholesterol levels >=200 mg/ dL despite adequate dietary recommendations	Parallel groups open Italy
<b>pitavastatin vs atorvastatin</b>			
<b>JAPAN ACS , 2009</b> [NCT00242944] n=307 follow-up: 8-12 months	pitavastatin 4 mg daily versus atorvastatin 20mg daily	patients with acute coronary syndrome undergoing IVUS-guided percutaneous coronary intervention	Parallel groups open Japan
<b>atorvastatin vs pravastatin</b>			
<b>PROVE IT - TIMI 22 , 2004</b> n=2099/2063 follow-up: 24 mo (18-36 mo)	80 mg of atorvastatin daily (intensive therapy). versus 40 mg of pravastatin daily (standard therapy)	patients who had been hospitalized for an acute coronary syndrome within the preceding 10 days	Parallel groups double blind UK, US, AUstralia, Italy, France, Germany, Spain, Canada

## References

### Sakamoto, 2006:

Sakamoto T, Kojima S, Ogawa H, Shimomura H, Kimura K, Ogata Y, Sakaino N, Kitagawa A Effects of early statin treatment on symptomatic heart failure and ischemic events after acute myocardial infarction in Japanese. Am J Cardiol 2006;97:1165-71 [16616020] [10.1016/j.amjcard.2005.11.031](https://doi.org/10.1016/j.amjcard.2005.11.031)

**MIRACL, 2001:**

Schwartz GG, Olsson AG, Ezekowitz MD, Ganz P, Oliver MF, Waters D, Zeiher A, Chaitman BR, Leslie S, Stern T Effects of atorvastatin on early recurrent ischemic events in acute coronary syndromes: the MIRACL study: a randomized controlled trial. *JAMA* 2001 Apr 4;285:1711-8 [[11277825](#)]

**macin, 2005:**

Macin SM, Perna ER, Faras EF, Franciosi V, Cialzeta JR, Brizuela M, Medina F, Tajer C, Doval H, Badaracco R Atorvastatin has an important acute anti-inflammatory effect in patients with acute coronary syndrome: results of a randomized, double-blind, placebo-controlled study. *Am Heart J* 2005;149:451-7 [[15864233](#)] [10.1016/j.ahj.2004.07.041](#)

**LIPS (sub groups), 2002:**

Serruys PW, de Feyter P, Macaya C, Kokott N, Puel J, Vrolix M, Branzi A, Bertolami MC, Jackson G, Strauss B, Meier B Fluvastatin for prevention of cardiac events following successful first percutaneous coronary intervention: a randomized controlled trial. *JAMA* 2002 Jun 26;287:3215-22 [[12076217](#)]

**FLORIDA, 2002:**

Liem AH, van Boven AJ, Veeger NJ, Withagen AJ, Robles de Medina RM, Tijssen JG, van Veldhuisen DJ Effect of fluvastatin on ischaemia following acute myocardial infarction: a randomized trial. *Eur Heart J* 2002;23:1931-7 [[12473255](#)]

**FLARE, 1999:**

Serruys PW, Foley DP, Jackson G, Bonnier H, Macaya C, Vrolix M, Branzi A, Shepherd J, Suryapranata H, de Feyter PJ, Melkert R, van Es GA, Pfister PJ A randomized placebo-controlled trial of fluvastatin for prevention of restenosis after successful coronary balloon angioplasty; final results of the fluvastatin angiographic restenosis (FLARE) trial. *Eur Heart J* 1999;20:58-69 [[10075142](#)]

**LCAS, 1997:**

Herd JA, Ballantyne CM, Farmer JA, Ferguson JJ 3rd, Jones PH, West MS, Gould KL, Gotto AM Jr Effects of fluvastatin on coronary atherosclerosis in patients with mild to moderate cholesterol elevations (Lipoprotein and Coronary Atherosclerosis Study [LCAS]). *Am J Cardiol* 1997;80:278-86 [[9264419](#)]

**Riegger et al., 1999:**

Riegger G, Abletshauser C, Ludwig M, Schwandt P, Widimsky J, Weidinger G, Welzel D The effect of fluvastatin on cardiac events in patients with symptomatic coronary artery disease during one year of treatment. *Atherosclerosis* 1999;144:263-70 [[10381299](#)]

**Czech trial, :****LAMIL, 1997:**

Kesteloot H, Claeys G, Blanckaert N, Lesaffre E Time course of serum lipids and apolipoproteins after acute myocardial infarction: modification by pravastatin. *Acta Cardiol* 1997;52:107-16 [[9187418](#)]

**RECIFE, 1999:**

Dupuis J, Tardif JC, Cernacek P, Throux P Cholesterol reduction rapidly improves endothelial function after acute coronary syndromes. The RECIFE (reduction of cholesterol in ischemia and function of the endothelium) trial. *Circulation* 1999;99:3227-33 [[10385495](#)]

**PAIS, 2001:**

Den Hartog FR, Van Kalmthout PM, Van Loenhout TT, Schaafsma HJ, Rila H, Verheugt FW Pravastatin in acute ischaemic syndromes: results of a randomised placebo-controlled trial. *Int J Clin Pract* 2001;55:300-4 [[11452676](#)]

**PACT, 2004:**

Thompson PL, Meredith I, Amerena J, et al. Effect of pravastatin compared with placebo initiated within 24 hours of onset of acute myocardial infarction or unstable angina: the Pravastatin in Acute Coronary Treatment (PACT) trial *Am Heart J*. 2004;148:e2

Thompson PL, Meredith I, Amerena J, Campbell TJ, Sloman JG, Harris PJ Effect of pravastatin compared with placebo initiated within 24 hours of onset of acute myocardial infarction or unstable angina: the Pravastatin in Acute Coronary Treatment (PACT) trial. *Am Heart J* 2004;148:e2 [[15215811](#)]

**CARE, 1996:**

Sacks FM, Pfeffer MA, Moye LA, Rouleau JL, Rutherford JD, Cole TG, Brown L, Warnica JW, Arnold JM, Wun CC, Davis BR, Braunwald E, The effect of pravastatin on coronary events after myocardial infarction in patients with average cholesterol levels. Cholesterol and Recurrent Events Trial investigators. *N Engl J Med* 1996; 335:1001-9 [[8801446](#)]

Plehn JF, Davis BR, Sacks FM, Rouleau JL, Pfeffer MA, Bernstein V, Cuddy TE, Moy LA, Piller LB, Rutherford J, Simpson LM, Braunwald E Reduction of stroke incidence after myocardial infarction with pravastatin: the Cholesterol and Recurrent Events (CARE) study. The Care Investigators. *Circulation* 1999;99:216-23 [9892586]

#### **LIPID, 1998:**

, Prevention of cardiovascular events and death with pravastatin in patients with coronary heart disease and a broad range of initial cholesterol levels. The Long-Term Intervention with Pravastatin in Ischaemic Disease (LIPID) Study Group. *N Engl J Med* 1998; 339:1349-57 [9841303]

, Long-term effectiveness and safety of pravastatin in 9014 patients with coronary heart disease and average cholesterol concentrations: the LIPID trial follow-up. *Lancet* 2002; 359:1379-87 [11978335]

Design features and baseline characteristics of the LIPID (Long-Term Intervention with Pravastatin in Ischemic Disease) Study: a randomized trial in patients with previous acute myocardial infarction and/or unstable angina pectoris. *Am J Cardiol* 1995;76:474-9 [7653447]

#### **4S, 1994:**

, Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S) *Lancet* 1994; 344:1383-9 [7968073]

#### **A to Z, 2004:**

de Lemos JA, Blazing MA, Wiviott SD, Lewis EF, Fox KA, White HD, Rouleau JL, Pedersen TR, Gardner LH, Mukherjee R, Ramsey KE, Palmisano J, Bilheimer DW, Pfeffer MA, Califf RM, Braunwald E Early intensive vs a delayed conservative simvastatin strategy in patients with acute coronary syndromes: phase Z of the A to Z trial. *JAMA* 2004 Sep 15;292:1307-16 [15337732]

#### **Ren, 2009:**

Ren HZ, Ma LL, Wang LX Effect of simvastatin on plasma interleukin-6 in patients with unstable angina. *Clin Invest Med* 2009;32:E280-4 [19640331]

#### **CIS, 1997:**

Bestehorn HP, Rensing UF, Roskamm H, Betz P, Benesch L, Schemitat K, Blmchen G, Claus J, Mathes P, Kappenberger L, Wieland H, Neiss A The effect of simvastatin on progression of coronary artery disease. The Multicenter coronary Intervention Study (CIS). *Eur Heart J* 1997;18:226-34 [9043838]

#### **HPS, 2002:**

, MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002; 360:7-22 [12114036]

Armitage J, Collins R Need for large scale randomised evidence about lowering LDL cholesterol in people with diabetes mellitus: MRC/BHF heart protection study and other major trials. *Heart* 2000;84:357-60 [10995396]

MRC/BHF Heart Protection Study of cholesterol-lowering therapy and of antioxidant vitamin supplementation in a wide range of patients at increased risk of coronary heart disease death: early safety and efficacy experience. *Eur Heart J* 1999;20:725-41 [10329064]

#### **Colivicchi, 2002:**

Colivicchi F, Guido V, Tubaro M, Ammirati F, Montefoschi N, Varveri A, Santini M Effects of atorvastatin 80 mg daily early after onset of unstable angina pectoris or non-Q-wave myocardial infarction. *Am J Cardiol* 2002;90:872-4 [12372577]

#### **ESTABLISH, 2004:**

Okazaki S, Yokoyama T, Miyauchi K, Shimada K, Kurata T, Sato H, Daida H Early statin treatment in patients with acute coronary syndrome: demonstration of the beneficial effect on atherosclerotic lesions by serial volumetric intravascular ultrasound analysis during half a year after coronary event: the ESTABLISH Study. *Circulation* 2004;110:1061-8 [15326073]

#### **GREACE, 2002:**

Athyros VG, Papageorgiou AA, Mercouris BR, Athyrou VV, Symeonidis AN, Basayannis EO, Demetriadis DS, Kontopoulos AG Treatment with atorvastatin to the National Cholesterol Educational Program goal versus 'usual' care in secondary coronary heart disease prevention. The GREek Atorvastatin and Coronary-heart-disease Evaluation (GREACE) study. *Curr Med Res Opin* 2002;18:220-8 [12201623]

#### **CLAPT, 1999:**

Kleemann A, Eckert S, von Eckardstein A, Lepper W, Schernikau U, Gleichmann U, Hanrath P, Fleck E, Neiss A, Kerber S, Assmann G, Breithardt and the CLAPT Study Effects of lovastatin on progression of non-dilated and dilated coronary segments and on restenosis in patients after PTCA. The cholesterol lowering atherosclerosis PTCA trial (CLAPT) *Eur Heart J* 1999;20:1393-406 [10487800]

**Sahni, 1991:**

Sahni R, Maniet AR, Voci G, Banka VS Prevention of restenosis by lovastatin after successful coronary angioplasty. Am Heart J 1991;121:1600-8 [2035374]

**L-CAD, 2000:**

Arntz HR, Agrawal R, Wunderlich W, Schnitzer L, Stern R, Fischer F, Schultheiss HP Beneficial effects of pravastatin (+/-colestyramine/niacin) initiated immediately after a coronary event (the randomized Lipid-Coronary Artery Disease [L-CAD] Study). Am J Cardiol 2000;86:1293-8 [11113401]

**PTT, 2002:**

Kayikioglu M, Can L, Kltrsay H, Payzin S, Turkoglu C Early use of pravastatin in patients with acute myocardial infarction undergoing coronary angioplasty. Acta Cardiol 2002;57:295-302 [12222700]

**OACIS-LIPID, 2008:**

Sato H, Kinjo K, Ito H, Hirayama A, Nanto S, Fukunami M, Nishino M, Lim YJ, Kijima Y, Koretsune Y, Nakatani D, Mizuno H, Shimizu M, Hori M Effect of early use of low-dose pravastatin on major adverse cardiac events in patients with acute myocardial infarction: the OACIS-LIPID Study. Circ J 2008;72:17-22 [18159093]

**GISSI Prevenzione, 2000:****JAPAN ACS, 2009:****PROVE IT - TIMI 22, 2004:**

## 11 statins high dose

Trial	Treatments	Patients	Trials design and methods
<b>atorvastatin high dose vs atorvastatin</b>			
<b>TNT , 2005</b> [NCT00327691] n=4995/5006 follow-up: 4.9 years	80 mg of atorvastatin daily versus 10 mg of atorvastatin daily	Chronic coronary artery disease LDL cholesterol <3.4 mmol/L	Parallel groups double blind 14 countries
<b>atorvastatin high dose vs lovastatin</b>			
<b>Vascular basis , 2005</b> n=197/103 follow-up: 1 year	atorvastatin (80 mg) with or without vitamin C and E versus low dose lovastatin (5 mg)	Chronic coronary artery disease	Parallel groups double blind
<b>atorvastatin high dose vs pravastatin</b>			
<b>PROVE-IT , 2004</b> n=2099/2063 follow-up: 2 years	atorvastatin 80 mg daily versus Pravastatin 40 mg	acute myocardial infarction (with or without electrocardiographic evidence of ST-segment elevation) or highrisk unstable angina) in the preceding 10 days	Parallel groups double blind 8 countries
<b>REVERSAL , 2004</b> n=327/327 follow-up: 1.5 years	atorvastatin 80 mg daily versus Pravastatin(40 mg)	Chronic coronary artery disease	Parallel groups double blind
<b>SAGE , 2007</b> n=446/445 follow-up: 1 years	atorvastatin 80 mg daily versus pravastatin(40 mg)	Chronic coronary artery disease	Parallel groups double blind
<b>atorvastatin high dose vs simvastatin</b>			

continued...

Trial	Treatments	Patients	Trials design and methods
<b>IDEAL , 2005</b> [NCT00159835] n=4439/4449 follow-up: 4.8 years	atorvastatin 80mg daily versus simvastatine 20mg/j	Men and women aged 80 years or younger with a history of a definite myocardial infarction and who qualified for statin therapy according to national guidelines	Parallel groups open Denmark, Finland, Iceland, Netherlands, Norway, Sweden
<b>simvastatin high dose vs simvastatin</b>			
<b>SEARCH , 2010</b> [NCT00124072] n=6031/6033 follow-up: 6.7 years (mean)	simvastatin 80 mg daily versus simvastatin 20mg daily	MI survivors	Parallel groups

## References

TNT, 2005:  
 Vascular basis, 2005:  
 PROVE-IT, 2004:  
 REVERSAL, 2004:  
 SAGE, 2007:  
 IDEAL, 2005:  
 SEARCH, 2010:

## 12 surgery

Trial	Treatments	Patients	Trials design and methods
<b>partial ileum bypass surgery vs no surgery</b>			
<b>POSCH , 1990</b> [NCT00000490] n=421/417 follow-up: 9.7 years	partial ileum bypass surgery versus no surgery	survivors to a first myocardial infarction	Parallel groups open

## References

POSCH, 1990:

## 13 About TrialResults-center.org

TrialResults-center is an innovative knowledge database that collects the results of RCTs and provides dynamic interactive systematic reviews and meta-analysis in the field of all major heart and vessels diseases.

The TrialResults-center database provides a unique view of the treatment efficacy based on all data provided directly from clinical trial results, offering a valuable alternative to personal bibliographic search, published meta-analysis, etc. Furthermore, it would allow comparing easily the various concurrent

therapeutic for the same clinical condition.

Rigorous meta-analysis method is used to populate TrialResults-center: widespread search of published and non published trials, study selection using pre-specified criteria, data extraction using standard form.

TrialResults-center is continually updated on a weekly basis. We continually search all new results (whatever their publication channel) and these news results are immediately added to the database with a maximum of 1 week.

TrialResults-center is non-profit and self-funded.